

Table of Contents

Chapter 3j Alaska Arctic Marine Fish Species

| | |
|-----------------------------------|----|
| Structure of Species Account..... | 2 |
| Marbled Eelpout..... | 10 |
| Arctic Eelpout..... | 14 |
| Threespot Eelpout..... | 18 |
| Archer Eelpout..... | 22 |
| Longear Eelpout..... | 26 |
| Scalebelly Eelpout..... | 30 |
| Estuarine Eelpout..... | 34 |

Chapter 3. Alaska Arctic Marine Fish Species Accounts

By Milton S. Love¹, Nancy Elder², Catherine W. Mecklenburg³, Lyman K. Thorsteinson², and T. Anthony Mecklenburg⁴

Abstract

Species accounts provide brief, but thorough descriptions about what is known, and not known, about the natural life histories and functional roles of marine fishes in the Arctic marine ecosystem. Information about human influences on traditional names and resource use and availability is limited, but what information is available provides important insights about marine ecosystem status and condition, seasonal patterns of fish habitat use, and community resilience. This linkage has received limited scientific attention and information is best for marine species occupying inshore and freshwater habitats. Some species, especially the salmonids and coregonids, are important in subsistence fisheries and have traditional values related to sustenance, kinship, and barter. Each account is an autonomous document providing concise information about a species zoogeography, western and Alaska Native taxonomy, life history, niches, and life requirements. Each account is fully referenced with the identification of the most critical literature for Alaska and a more comprehensive listing of referencing from which biological and ecological information was drawn. New-to-science narratives, distributional maps, and vertical profiles, provide quick, reliable sources of information about fish life history and habitat requirements for this segment of the Arctic fauna.

Purpose and Design of Species Accounts

Individual species accounts were prepared for 104 of the 109 confirmed marine fishes for which adequate biological information was available from the U.S. Chukchi and Beaufort Seas. These descriptions are an important source of documentation about Arctic Alaska's marine fish fauna.

Although tailored to address the specific needs of BOEM Alaska OCS Region NEPA analysts, the information presented in each species account also is meant to be useful to other users including state and Federal fisheries managers and scientists, commercial and subsistence resource communities, and Arctic residents. Readers interested in obtaining additional information about the taxonomy and identification of marine Arctic fishes are encouraged to consult the *Fishes of Alaska* (Mecklenburg and others, 2002) and *Pacific Arctic Marine Fishes* (Mecklenburg and others, 2016). By design, the species accounts enhance and complement information presented in the *Fishes of Alaska* with more detailed attention to biological and ecological aspects of each species' natural history and, as necessary, updated information on taxonomy and geographic distribution.

Each species account includes a concise summary of the natural history, population dynamics, functional roles, and traditional and economic values of the marine fish found off Alaska. An initial organizational task was to create a standard format for effective information delivery. The species descriptions by Ehrlich and others (1988) were provided to the USGS by BOEM as an example of a creative template for information transfer. Four pilot species accounts, representing well known to poorly known species, were developed, reviewed, and repeatedly revised for improvements, interagency approval, and selection of the final layout and design. Final decisions about content represented the priority needs of BOEM.

More than 1,200 individual scientific publications relevant to Arctic marine fishes were reviewed in preparation of the species accounts. In each species account, the most relevant literature for each species is cited. A shorter list (about 5–10 articles) identifies key Alaskan information sources that, in our opinion, have had the greatest scientific effect on understanding the species of the Arctic area of the United States.

¹University of California, Santa Barbara.

²U.S. Geological Survey.

³California Academy of Sciences, San Francisco, and Point Stephens Research, Auke Bay, Alaska.

⁴Point Stephens Research, Auke Bay, Alaska.

Limitations of Data

The species accounts reveal many gaps in the biological information needed to conduct vulnerability assessments of the marine fishes of the Beaufort and Chukchi Seas to human interventions. Part of this problem relates to the geographic coverage of existing research and surveys in Alaska as, in many instances, we were required to incorporate the results of investigations conducted outside the region. This raises an important caution because, even though the best available information was used in preparing the species accounts, our reliance on data and information from outside Alaska will introduce uncertainty to EIS expectations. Ideally, and with respect to oil and gas activities, baseline information for fishery resources should be collected from the potentially affected environment to appropriately evaluate the potential effects of oil spills or other possible industrial-related disturbances. However, as has been widely noted (for example, Bluhm and others, 2011), systematic and methodologically comparable data typically are not available from Arctic Alaska marine ecosystems. Evaluating change in populations and communities from natural and anthropogenic stressors is limited by the variable quality and lack of quantitative reports on abundance, distribution, community structure, and demographics for Arctic marine fishes.

In each species account, an attempt was made to incorporate the most reliable baseline information available and offer impressions of information needs. Important ongoing studies sponsored by BOEM, and others, may be addressing some of these needs. The needs assessments for this study considered these efforts to the extent that oral and (or) written communications and preliminary results allowed. The focus of this study was on impressions of the population parameters (Williams and others, 2002) and environmental measurements needed to detect changes in marine fish populations (Reist and others, 2006; Wassmann and others, 2011) and their resilience to a variable and rapidly changing environment (Holland-Bartels and Pierce, 2011). For key marine fish species, examples might include changes in range, community structure, abundance, phenology, behavior, and population growth and survival.

Each species account is designed as a self-contained article; therefore, no references to other accounts are included. Additionally, to reduce complexity in the presentations, only common names were used to identify the major predator and prey species for the marine fish described. Because this document was meant to be a companion document to the *Fishes of Alaska* (Mecklenburg and others, 2002), interested readers are encouraged to consult this book or Page and others (2013) and Mecklenburg and others (2016) for more complete information about the scientific authorities and literature citations associated with the original descriptions of each species. Readers are directed to the references cited in each species account for additional information on the species.

Operational Definitions

In [chapter 1](#), several concepts about the temporal and spatial habitat requirements for Arctic marine fish were introduced. More information is presented in this chapter to explain the vertical distribution and the location of shelf break, as used in this report.

Vertical Distribution

The conceptual design of the species depth profiles (vertical structure by life history stage) was patterned after the “coastal marine life zones” of Allen and Smith (1988). The goal of the profiles is to visualize what is known about a species occurrence and reproductive ecology by depth and location. An idealized characterization of Arctic shelves was designed to visualize these relationships. Additional detail about origins of data was included in the depth profiles to reflect Alaskan records or collections from other Arctic regions. This is important because actual field collections and observations are limited from this region. In many instances, the actual presence of a life stage remains unverified by field sampling. Thus, for many of species, the depth of a fish’s life cycle should be considered untested hypotheses in need of additional testing.

Location of Shelf Break

Early versions of the depth profiles were modified at the request of BOEM with respect to the depiction of the continental shelf break. As a special effect for the Arctic, the species depth profiles were redrawn to depict the change in bathymetry that typically occurs at depths of about 75 m throughout the Chukchi and western Beaufort Seas. This depiction is not an attempt to redefine the oceanographic definition of shelf break. Instead, it highlights the relatively sharp gradient in depths that often occurs near 70- to 80-m contours over much of the region. Although species depth profiles in this report depict an apparent “break” at 75-m, three factors were considered: (1) this is a generalization and the actual shelf break may be geographically close but at a slightly greater depth; (2) shelf edge effects on fish distribution at depths occurring between 75-, 150-, or 200-m are likely negligible due to the gradient and area involved; and (3) the conceptual depictions of depth distributions by life history stage are consistent with accepted oceanographic conventions for continental shelf and slope (despite the magnified view at 75-m) and thus are compatible to the import of biological data obtained elsewhere.

Keystone Species

The concept of keystone species describes the critical role certain organisms are perceived to have in maintaining the structure of biological communities and resilience of ecosystem dynamics (Paine, 1966). Arctic Cod (*Boreogadus saida*) are widely distributed in the Arctic Ocean and by virtue of their abundance and intermediate trophic position between invertebrates and higher-level predators are integral to the movement of nutrients in marine food webs. For this reason, Arctic Cod are considered a keystone species in the Arctic marine (Bradstreet and others, 1986; Walkusz and others, 2011). Arctic Cod are common in United States waters of the Beaufort and Chukchi Seas being considered for energy exploration and development and are an ecological focus of BOEM fishery studies to understand potential effects on the species (Maule and Thorsteinson, 2012).

Outline of Species Accounts

The species accounts are scientifically accurate descriptions of the life histories, populations, habitats, and community values of individual species in the Arctic marine ecosystem. The mix of quantitative and qualitative information presented reflects state-of-the-art knowledge, a faunal assessment of information gaps, and prioritization of priority needs for population and process understanding. Limited information for many Alaskan species required that relevant observations from other geographic locales be included. Each species account attempts to be clear about the geographic origins of data and information, through scientific referencing or special notations in graphics. As an example, *italics* are used in the species accounts to highlight data collections from the Alaska study area. In several instances, species information was so lacking that inferences from a closely related species were required.

The generic species account includes a comprehensive accounting of scientific and cultural information in a standard format. The scientific information addresses multiple disciplinary areas including taxonomy, life history and habitats, ecological relationships including predator-prey interactions and environmental preferences, and population ecology. The population information is critical to evaluations of population status and health, resilience, and vulnerability to natural and anthropogenic changes in the marine environment. Each species account includes a photograph of an adult specimen (or line drawing if an image was not available); distribution maps (horizontal and vertical); and concise descriptions of abundance, life history, and ecology (11 life history categories); major stressors; research needs; and key references. To assist users, a suite of easily recognized icons was developed to provide quick access to specific life history information. In addition, some species attributes

regarding life history, population dynamics, and biological interactions are defined in the Glossary (chapter 7).

Information presented in each species account is outlined and described as:

Taxonomic—Scientific and Common Names

The format of the species accounts was, by design, intended to link the biologic and ecologic information presented in this document directly to the species identification guides contained in the “*Fishes of Alaska*.” This connection was established by adherence to naming conventions as described by Mecklenburg and others, 2002 (p. 25 and 26). The common names of each marine fish are presented first, followed by scientific and family names. Each scientific name includes a reference to the name of the person (author) who formally described and named the species in the ichthyological literature. The bibliographic data for the authors and dates of publication of scientific names can be found in Eschmeyer’s Catalog of Fishes online (<http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>) and are not reported here. In some instances, a *Note* (italicized) has been included to describe exceptional details about existing biological data, morphology, nomenclature, taxonomic status, life history strategy, or occurrence of a species in the United States Chukchi and Beaufort Seas.

Iñupiat Name

The existence of colloquial Iñupiat (Iñupiaq) names for the Arctic’s marine fish fauna by indigenous peoples is an important component of traditional ecological knowledge. Relatively few marine fish species are abundant or susceptible enough to subsistence fisheries to have received special names. For those species having Iñupiat names, this information is reported to assure that a common vocabulary can facilitate future exchanges of ideas and knowledge across disciplinary boundaries. In this manner, colloquial names can provide a cultural link between local marine resources and science supporting sustainability of Arctic communities and ecosystems.

Ecological Role

Fishes play a pivotal role in marine ecosystems as secondary and higher-level consumers in many marine food webs. In many instances, information about predator-prey relationships is so limited that only preliminary, qualitative assessments of the relative role of each species are possible. The ecological niche describes how an organism or population responds to resources and competitors. Importance or significance descriptors do not diminish the fact that all organisms contribute in ways large or small to the provision

of ecosystem goods and services. These descriptors however, may provide useful information about the relative importance of a particular species as an indicator of ecosystem condition and trajectories of change associated with climate change, habitat fragmentation, ecosystem stress, effect of pollutants, or other anthropogenic effects.

Physical Description/Attributes

A brief physical description of the species is summarized from information presented by Mecklenburg and others, (2002) in the *Fishes of Alaska*; the relevant page number is included for quick referral to more comprehensive morphological information. An image of the adult form of each fish is presented with appropriate attribution. High-quality images were selected to highlight the key identifying features of a particular species.

Information about the presence of a swim bladder and antifreeze glycoproteins is included because of its relevance to geo-seismic oil and gas exploration, climate change issues, and evolutionary life history.

Range

The geographic occupancy of the species in United States sectors of Chukchi and Beaufort Seas and adjacent waters is presented in brief narratives and depicted on maps. Known occurrence in the Arctic OCS Planning Areas is highlighted by symbols indicating locations of valid species identifications from properly archived voucher specimens on each map. Although the symbols on the maps may suggest that some of the species are rare in the region, the study of historical collections from the United States and Canadian sectors of the Beaufort Sea, as well as the collections from BOEM surveys in the Beaufort in 2011 and 2012, is still in progress and may reveal that these species are more abundant in deep sectors of the study area than the maps suggest. Definitions of zoogeographic pattern are from the Online Resource 1 (electronic supplemental to Mecklenburg and others, 2011), Mecklenburg and Steinke (2015), and Mecklenburg and others (2016) and relate to ranges of population viability (see [chapter 2](#)).

Depth profiles in each species account graphically summarize existing information about the benthic and reproductive distributions of each marine fish. In both depth profiles, the width of areas depicted confers species information about horizontal (onshore-offshore) patterns of distribution. The italicized captions in the depth profiles highlight species information germane to the study area. Areas in the graphs denoted by the orange coloration represent understanding from data collection within the United States Chukchi and Beaufort Seas; olive colors represent data collection outside the study area. For benthic distributions,

solid lines in the depth profiles represent species for which no specific information is available about its preferred depth range. Solid lines represent a synthesis of understanding that includes information not necessarily specific to the study area. In some instances, only one record of a species occurrence by depth was available and coding in orange was not meaningful. In these cases, an explanatory comment, in italicized font, with a line pointing to the appropriate depth was included in the graph (for example, see the species account for *Megalocottus platycephalus*). Highlighted depths as indicated through “bolded” (dark black) and dashed segments, represent most common depths where the species has been detected, and depth distribution as has been reported throughout the species range, respectively. Areas denoted with diagonal cross-hatching represents depth distribution of juveniles (immature); adult distributions are not cross-hatched and age-related habitat overlaps, are informed by captioning in the figures.

For reproductive distribution, eggs and larvae (pre-juvenile life stages) of marine fishes are represented with respect to depth and distance from the coast. Orange areas in the reproductive distribution profiles represent data collection in the study area. In many instances, information about spawning habitats and egg and larval distributions is summarized from information reported from throughout a species range. In these cases, dark blue represents species distributions in spawning habitats; light blue represents the geographic distributions of eggs and larvae; and light green is used to highlight areas of substantial habitat overlap (for example, see the species account for *Hippoglossus stenolepis*). Distribution patterns of eggs and larvae are symbolized by “dots” and “horizontal dashes,” respectively, in the graphs. As for benthic distribution, solid lines represent species-specific information from data collections from throughout the species entire range. Highlighted (dark black lines) segments of solid lines indicate the most common depths where egg and larvae samples have been collected. Dashed lines represent areas of hypothesized distributions for species for which no information is available about egg or larval occurrence. In these instances the hypothesized distributions are based on known patterns for closely related species; the lack of data is stated in captions above the graph.

Relative Abundance

Relative abundance refers to the contribution a species makes to the total abundance of the fishery community. It is a measure that provides an index of the number of individuals present, but not the actual numbers. Relative abundance terms, such as “common,” “uncommon,” or “rare” often are used to express the general population status of a given species, but are most useful when they are defined by something that is measured or estimated in a manner that makes comparison meaningful.

Depth Range

Benthic distribution refers to the spatial arrangement of a particular species at different depths over continental shelf and slope waters. The life cycle of fishes occurs in multiple dimensions in time and space and generally reflects genetically determined life history or behavior that has evolved to maximize fitness (life time reproductive success, see Gross [1987]). Benthic distribution profiles for each species represent the location of important habitats as they are presently known for juvenile and marine fishes. Reproductive distributions depict important habitats for spawning and early life history development.

Life History, Population Dynamics, and Biological Interactions


Life history theory holds that the schedule and duration of key events in a species' lifetime are shaped by natural selection to produce the largest possible number of surviving offspring. These events, notably juvenile development, age of sexual maturity, first reproduction, number of offspring and level of parental investment, senescence, and death, depend on the abiotic and biotic environment of the organism. Specific information about these traits informs understanding of a species' adaptive capacity including major influences on population abundance. A number of fisheries models use basic length-weight and age-at-size relationships to describe the growth and dynamics of fishery populations (for example, von Bertalanffy and Gompertz, growth models and derivatives [Ricker, 1975]). Ecological models estimate transfer of energy or matter along the trophic chain (Gamito, 1998). The parameters that are estimated in these models are individually important indicators of population condition and may be used with other indicators to derive quantitative information about compensatory responses and resilience. Much of this information, including population parameters, has been compiled in FishBase for the Arctic marine fish (Froese and Pauly, 2012).



Habitats and Life History—Basic information about the life history (for example, body size, reproductive ecology, growth) and ecology (for example, mobility, growth, habitat) of a species and the environmental area inhabited by that species is foundational to

effective resource management. Habitat is the natural environment that influences and is used by a species population. Information about abiotic (that is, temperature, salinity, other physiochemical factors, depth, and substrate types) and biotic (that is, type and abundance of food, presence of other biota) often are used to describe fish habitats and provide insights about a species environmental preferences and habitat associations (for example, water masses). Maximum body size often is reported and can be an

important surrogate of different life history traits (for example, age at maturity, growth, and reproductive output). In population dynamics studies, the relationships between length and weight and size and age form the basis for population growth and production models and quantitative analysis of environmental effects. Length measurements are reported as standard length (SL), total length (TL), and fork length (FL) in fisheries studies.



Behavior (see also Glossary [chapter 7]).—

Behavior is the manner in which a fish operates or functions within its environment (that is, home range, territoriality, and many others) to procure food, orient to specific locations, or relate to other organisms. Knowing how


individuals respond to the environment (physical, chemical, and biological cues) is critical to understanding population processes such as distribution, survival, and reproduction and recruitment and for managing fisheries. Many behaviors are evolutionary adaptations to the physiological and reproductive requirements for a species' survival. For example, migration involves the regular movement of animals between different geographic locations. Migrations can be extensive in terms of time and distance involved (anadromous model) or seasonal (amphidromous and marine models). Each of these models reflects a life strategy adapted for age and growth at sea. Diel relates to daily changes in water column position due to changes in light, temperature, and food supply.

Migratory behaviors are rooted in physiological requirements for food, growth, reproductive, and survival ("scope for growth"). Movement behaviors are more tactical responses to local environmental conditions (for example, variable hydrographic conditions in the nearshore Beaufort Sea). Fish movement can be active or passive and involve large distances in search of suitable habitats and foods. The seasonal nature of migration and movement behaviors are typically related to life history stage, predator-prey distributions, or energetic requirements for growth.

Schooling (that is, social structure of fish of the same species moving in more or less harmonious patterns in the sea) often is related to survival and reproduction. Schooling confers physical benefits to fish movement, safety against predators, search behaviors (for example, foods), population immunology, and reproduction.


The functional feeding morphology of a fish relates to its anatomical adaptations (for example, body size, gape sizes, shape, and body form) to environmental conditions especially food preferences. The adage "function determines morphology and morphology determines way of life" is an important evolutionary concept as it applies to fish feeding behavior, dietary preferences, habitat selection, and trophic stature. Trophic position (within categories of trophic levels) expresses the "tendency of larger (less abundant) fishes feeding on smaller (more abundant) fishes, which themselves feed on zooplankton and all these animals resting upon primary producers" (from Pauly and Watson, 2005). Categories of trophic levels are:

- Trophic level 1 (T1), plants and animals make their own food and are called primary producers;
- Trophic level 2 (T2), herbivores eat plants and are called primary consumers;
- Trophic level 3 (T3), carnivores eat herbivores and are called secondary consumers;
- Trophic level 4 (T4), carnivores eat other carnivores and are called tertiary consumers; and
- Trophic level 5 (T5), apex consumers, which have no predators, are at the top of the food chain.



Populations or Stocks—A population often is defined as a group of organisms of the same species occupying a particular space at a particular time with the potential to breed with each other (Williams and others, 2002). Stocks are subpopulations of a particular species of

fish that result from reproductive isolation and subdivisions within the biological range. The current state of knowledge about local stocks and their genetic population structure is reported. Grossberg and Cunningham (2001) described the combined effects of demographic, behavioral, genetic, oceanographic, climate, and tectonic processes as major determinants of population structure. These mechanisms act across a range of temporal and spatial scales to determine the rates and patterns of dispersal of different life stages of marine fishes. Dispersal, combined with the successful reproduction and survival of immigrants, control the scale and rate of processes that build or erode structure within and among groups of individuals.




Reproduction Mode—Little information is available about the spawning times and locations, mating behaviors (breeders or nonbreeders), and genetic diversity of Arctic marine fishes. What is known is drawn largely from observations from populations studied

outside the United States. For most Arctic marine fish species, there is no information about population or stock structure (for example, age structure, reproductive behavior, sex ratios, age-at-maturity, fecundity, and genetic). These are key population parameters needed for understanding reproductive ecology, population dynamics (for example, growth, survival, and mortality), and assessments of resiliency (response to disturbance).




Food and Feeding—Dietary information is summarized from literature and, unless in italics, is reported from other regions. Fish communities can affect the ecological characteristics of marine ecosystems in

response to productivity and abundance patterns, the mobility and migratory behavior of species, and through food influences in different habitats (for example, Grebmeier and others, 2006b). Trophic Index (T) values are reported from FishBase (Froese and Pauly, 2012). The T values for Arctic marine fishes are largely derived from stomach contents analyses, which have correlated well with stable isotopes of nitrogen in tissues. The fractional values (between 1 and 5) realistically address complexities of consumer feeding behaviors (omnivory and feeding across multiple trophic levels) and predator-prey relationships. For example, the mean T value for Blackline Prickleback (*Acantholumpenus mackayi*) is 3.1 (± 0.31). This mid food web value is indicative of a primary carnivore that feeds across trophic levels, in this case on lower level herbivores.



Biological Interactions.—The effects organisms in a community have on one another. Competition and consumption (predation, herbivory, or cannibalism) are the best known of the major ecological processes affecting resource abundance, community

composition, and ecosystem function. Competition involves interactions between individuals of the same species (intraspecific) or different species (interspecific) in which the fitness of one is lowered by the presence of another. Competition often is related to food and habitat requirements and reproductive behavior. Interspecific competition for foods is greatest for species occupying similar trophic positions in relatively short food chains and for animals living in regions of low biological productivity.



Resilience—In ecology, resilience traditionally refers to the ability of a population or biotic community to sustain or return to its former state after a disturbance. The rate of recovery is a measure of resilience determined by the population processes involved in restoring abundance to healthy, sustainable, or pre-disturbance levels. Four categories of productivity (high, medium, low, and very low) are used to classify reliance in marine fish populations (Musick, 1999). These categories are based on a combination of population parameters for intrinsic rate of growth, growth coefficient, fecundity, age at maturity, and maximum age. Because population parameters were unavailable, resiliency is defined here based on estimated population doubling time where high = <15 months, medium = 1.4–4.4 years, and low = 4.5–14 years.

Traditional, Cultural, and Economic Values

In August 2009, the U.S. Secretary of Commerce approved a Fishery Management Plan for the Arctic Management Area. The plan covers U.S. Arctic waters in the

Chukchi and Beaufort Seas, and acknowledges that changing climate may potentially favor the development of commercial fisheries. However, until adequate fisheries resource assessments are completed, the region remains closed to commercial fishing in federal waters. A small salmon fishery exists in Kotzebue Sound; in 2010, a small commercial fishery for Arctic Ciscoes in the Colville River was terminated.



Traditional and Cultural Importance.—

Several species of nearshore marine fishes are important in subsistence fisheries. The protection of traditional lifestyles and economies, including these subsistence fisheries, is a responsibility of the Federal government. Subsistence relates to resource use patterns (for example, seasonal round) and values (that is, sustenance, kinship, and barter) in coastal communities of northern Alaska.



Commercial Fisheries.—Currently (2016) there are no offshore marine fisheries in the U.S. Chukchi and Beaufort seas. Changing Arctic environmental conditions and shifting distributions of species in response to warming suggest that there may be fisheries in the

future. A precautionary approach by fishery managers has been adopted that requires the collection of reliable baseline information for decision-making and ecosystem management (North Pacific Fishery Management Council [North Pacific Fishery Management Council, 2009; Wilson and Ormseth, 2009]).

Climate Change

Alaska's climate is changing at more than twice the rate of the rest of the United States (Mellilo and others, 2014). Year-to-year and regional variability in air temperatures are evident and the warming trend currently is being moderated by large-scale cooling associated with the Pacific Decadal Oscillation. Even so, climate effects are pronounced and are being seen in changes in sea ice, timing of snowmelt, widespread glacier retreat, and changes in hydrology (runoff) and coastal processes, such as erosion (Markon and others, 2012). The effects of rising ocean temperatures and ocean acidification on marine food webs are of growing regional concern with respect to the condition and trends in marine ecosystems and human community resilience are of concern. Climate changes potentially can affect marine fish in numerous ways, leading to distributional changes, increased or decreased mortality rates, changes in growth rates, and by altering the timing in reproduction (Clow and others, 2011).




Potential Effects of Climate Change.—A pole-ward shift of many fish distributions is possible as is a reduction or extinction of species that are narrowly adapted to Arctic

environments. Generally, the species are expected to increase in abundance if they are currently present in the Bering Sea and decrease if they have very low tolerance for temperatures greater than 1.5–2.0 °C. However, it is hypothesized in current climate projections that temperatures near the ocean floor in the northern Bering Sea will remain cold (<2 °C) due to persistence of winter sea ice (Sigler and others, 2011). Cold-water conditions and other marine ecosystem effects related to seasonal sea ice extent and timing of retreat may effectively block northward migrations and production of exploitable quantities of species, such as pollock and cod, for several decades. Shifts in range and other possible climate-related effects, such as increased predation or competition for food, are identified in the species accounts. Only “loose qualitative generalizations” are presently possible (Reist and others, 2006).

Research Needs

The compilation and review of species information for species in U.S. Arctic waters revealed many gaps in life history understanding and environmental relations. These are evaluated on the basis of a species current fishery and community values and ecological significance in marine ecosystem structure and function. The needs reflect the researcher's perceptions and their understanding that new fishery information is becoming available for the Arctic region and that, although Arctic research is currently a national priority, some aspects of population ecology will take many years of data collection to accurately assess.



Areas for Future Research.—The preparation of individual accounts led to the identification of many information gaps in knowledge about the biology and ecology of marine species including life history, population dynamics, and community associations. Generally,

species life history and ecology gaps are most pronounced with respect to: (1) depth and location of pelagic larvae; (2) depth, location, and timing of young-of-the-year habitats; (3) preferred depth ranges for juveniles and adults; (4) spawning seasons; (5) seasonal and ontogenetic movements; (6) population genetics and dynamics; (7) prey–predator relationships and food web relationships; and (8) environmental health (multiple stressor effects on fitness). Behavioral studies for all life stages are virtually non-existent. New information is being developed and, for the lesser-known species, gaps may be slowly addressed over time. Priority needs, for species having special significance in subsistence fisheries and marine food webs or that may be indicator species are emphasized in the species accounts. One of two categories of identified research need is identified for each species. The meaning of the categories [A] and [B] is as follows:

- **[A]** Many gaps in our understanding of the species life history and ecology remain in Alaska (for example, research areas 1 through 8). These are high profile species in terms of ecological, subsistence, or potential fisheries values. Specific research priorities are briefly discussed.
- **[B]** Most aspects of the species life history and ecology are unknown for Alaska (for example, research areas 1 through 8). Species information will likely accumulate over time and focused studies are not warranted at this time.

References Cited and Bibliography

A thorough review of scientific literature was done in the preparation of the species account. A list of references (References Cited [[chapter 8](#)]) is provided for each species for readers seeking additional information. This list identifies key sources of information that make the greatest contributions to current knowledge (2014) and understanding. The Bibliography section provides a full accounting of all scientific literature cited in each species account. For a small number of species from the family Cottidae, only a Bibliography was possible to provide and this is indicative of the lack of information available. Citations are not always in numerical order in species accounts because new information became available during the production phase of this publication and were incorporated into the species accounts as appropriate.

Marbled Eelpout to Banded Gunnel

Marbled Eelpout (*Lycodes ravidens*)

Taranetz & Andriashev, 1937

Family Zoarcidae

Colloquial Name: *None within U.S. Chukchi and Beaufort Seas.*

Ecological Role: Common but not abundant, the Marbled Eelpout is likely of relatively small ecological importance in the U.S. Chukchi and Beaufort Seas.

Physical Description/Attributes: Elongate cream to tan body with 7–9 brown or reddish brown bands extending onto dorsal and anal fins; blackish dorsal fin margin at ends of the bands; head, nape, and body bands becoming marbled in appearance in adults. For specific diagnostic characteristics, see *Fishes of Alaska* (Mecklenburg and others, 2002, p. 711) [1]. Swim bladder: Absent [1]. Antifreeze glycoproteins in blood serum: Unknown.

Range: *U.S. Chukchi and Beaufort Seas. Northernmost record in U.S. Chukchi Sea is at 71°27'N, 158°02'W, and in U.S. Beaufort Sea is at 71°13.5'N, 152°47.9'W; easternmost record in U.S. Beaufort Sea is north of Arey Island at 70°36'N, 143°55'W* [2]. Elsewhere in Alaska, eastern Bering Sea to Bristol Bay and one record north of Near Islands, western Aleutian Islands [1]. Worldwide, in western Bering Sea and Commander Islands to Okhotsk Sea [1] and in East Siberian Sea [2].

Relative Abundance: *Common in U.S. Chukchi and Beaufort Seas* [2, 4]. Patchily abundant (common) from eastern Sakhalin Island, Russia through Bering Sea and to *Chukchi Sea and Beaufort Seas* [2, 4–9].



Marbled Eelpout (*Lycodes ravidens*), 102 mm, Chukchi Sea, 2007. Photograph by C.W. Mecklenburg, Point Stephens Research.




Geographic distribution of Marbled Eelpout (*Lycodes ravidens*) within Arctic Outer Continental Shelf Planning Areas [3] based on review of published literature and specimens from historical and recent collections [1, 2, 4].


Depth Range: 8–467 m, rarely deeper than 150 m [4]. Taken in U.S. Chukchi Sea between 55–59 m [9]. A maximum depth of 525 m was reported for fish wintering off shelf in Sea of Okhotsk [10]. Generally, eelpout spawning and larvae occur at same depths that adults inhabit [1].

Lycodes ravidens
Marbled Eelpout

Benthic distribution



Reproductive distribution



Benthic and reproductive distribution of Marbled Eelpout (*Lycodes ravidens*).



Habitats and Life History

Eggs—Size: Unknown. Time to hatching: Unknown. Habitat: Benthic [1].


Larvae—Size at hatching: Unknown. Size at juvenile transformation: Unknown. Days to juvenile transformation: Unknown. Habitat: Benthic [1].

Juveniles—Age and size: Unknown. Habitat: Benthic [1].

Adults—Age and size at first maturity: Unknown. Maximum age: 7 years in western Bering Sea [7]. Maximum size: To 86 cm TL [7]. Habitat: Benthic [1].

Substrate—Sandy mud and mud [1, 11].

Physical/chemical—Temperature: -1.7 – 7.9 °C [7, 11, 12]. Salinity: Marine, for example, 32.58 parts per thousand in the U.S. Chukchi Sea [9] and occasionally in brackish waters [4].



Behavior

Diel—Unknown.

Seasonal—Migrates to deeper water in Sea of Okhotsk in winter [10].

Reproductive—Unknown.

Schooling—Unknown.

Feeding—Unknown.



Populations or Stocks

There have been no studies.



Reproduction

Mode—Separate sexes, oviparous.

Spawning season—Likely in autumn or autumn-winter in western Bering Sea [7].


Fecundity—Unknown.



Food and Feeding

Food items—Benthic and epibenthic prey, such as gammarid amphipods, euphausiids, shrimps, polychaetes, and clams [7].


Trophic level—3.6 [13].



Biological Interactions

Predators—Unknown.

Competitors—Likely other benthic-dwelling fishes, including flatfishes, sculpins, snailfishes, and other eelpouts.



Resilience

Medium, minimum population doubling time: 1.4–4.4 years (Fecundity=3,116) [14].




Traditional and Cultural Importance

None reported.




Commercial Fisheries

Currently, Marbled Eelpout are not commercially harvested.



Potential Effects of Climate Change

The Marbled Eelpout is a predominantly Boreal Pacific species. Increases in abundance and interspecific competition are possible outcomes of climate warming.



Areas for Future Research [B]

Little is known about the biology and ecology of this species from the region. Research needs include:

- (1) depth and location of pelagic larvae, (2) depth, location, and timing of young-of-the-year benthic recruitment, (3) preferred depth ranges for juveniles and adults, (4) spawning season, (5) seasonal and ontogenetic movements, (6) population studies, (7) prey, and (8) predators.

References Cited

- Andriashev, A.P., 1954, Fishes of the northern seas of the U.S.S.R.—Keys to the fauna of the U.S.S.R.: Academy of Sciences of the U.S.S.R., Zoological Institute, no. 53, 566 p. [In Russian, translation by Israel Program for Scientific Translation, Jerusalem, 1964, 617 p., available from U.S. Department of Commerce, Springfield, Virginia.] [11]
- Balanov, A.A., Badaev, O.Z., Napazakov, V.V., and Chuchukalo, V.I., 2006, Distribution and some biological features of *Lycodes ravidens* (Zoarcidae) in the western part of the Bering Sea: *Journal of Ichthyology*, v. 46, no. 2, p. 148–155. [7]

Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, *Fishes of Alaska*: Bethesda, Maryland, American Fisheries Society, 1,116 p. [1]

Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: *Marine Biodiversity*, v. 41, no. 1, p. 109–140, Online Resource 1. [2]

Mecklenburg, C.W., Stein, D.L., Sheiko, B.A., Chernova, N.V., Mecklenburg, T.A., and Holladay, B.A., 2007, Russian–American long-term census of the Arctic—Benthic fishes trawled in the Chukchi Sea and Bering Strait, August 2004: *Northwestern Naturalist*, v. 88, no. 3, p. 168–187. [9]

Bibliography

1. Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, *Fishes of Alaska*: Bethesda, Maryland, American Fisheries Society, 1,116 p.
2. Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: *Marine Biodiversity*, v. 41, no. 1, p. 109–140, Online Resource 1.
3. Minerals Management Service, 2008, Beaufort Sea and Chukchi Sea planning areas—Oil and Gas Lease Sales 209, 212, 217, and 221: U.S. Department of the Interior, Minerals Management Service Alaska OCS Region, OCS EIS/EA, MMS 2008-0055, 538 p.
4. Mecklenburg, C.W., Mecklenburg, T.A., Sheiko, B.A., and Steinke, D., 2016, Pacific Arctic marine fishes: Akureyri, Iceland, Conservation of Arctic Flora and Fauna, Monitoring Series Report No. 23, 406 p., accessed May 10, 2016, at <http://caff.is/monitoring-series/370-pacific-arcticmarine-fishes>.
5. Barber, W.E., Smith, R.L., Vallarino, M., and Meyer, R.M., 1997, Demersal fish assemblages of the northeastern Chukchi Sea, Alaska: *Fishery Bulletin*, v. 95, no. 2, p. 195–209.
6. Kim, S.T., and Shepeleva, O.N., 2001, The structure of shelf ichthyocenoses of northeastern Sakhalin and Terpeniya Bay: *Journal of Ichthyology*, v. 41, no. 9, p. 711–722.
7. Balanov, A.A., Badaev, O.Z., Napazakov, V.V., and Chuchukalo, V.I., 2006, Distribution and some biological features of *Lycodes raridens* (Zoarcoidea) in the western part of the Bering Sea: *Journal of Ichthyology*, v. 46, no. 2, p. 148–155.
8. Hoff, G.R., 2006, Biodiversity as an index of regime shift in the eastern Bering Sea: *Fishery Bulletin*, v. 104, no. 2, p. 226–237.
9. Mecklenburg, C.W., Stein, D.L., Sheiko, B.A., Chernova, N.V., Mecklenburg, T.A., and Holladay, B.A., 2007, Russian–American long-term census of the Arctic—Benthic fishes trawled in the Chukchi Sea and Bering Strait, August 2004: *Northwestern Naturalist*, v. 88, no. 3, p. 168–187.
10. Kim, S.T., 2001, Winter migrations of shelf fish to the continental slope: *Journal of Ichthyology*, v. 41, no. 8, p. 564–574.
11. Andriashev, A.P., 1954, *Fishes of the northern seas of the U.S.S.R.—Keys to the fauna of the U.S.S.R.*: Academy of Sciences of the U.S.S.R., Zoological Institute, no. 53, 566 p. [In Russian, translation by Israel Program for Scientific Translation, Jerusalem, 1964, 617 p., available from U.S. Department of Commerce, Springfield, Virginia.]
12. Mueter, F.J., University of Alaska-Fairbanks, written commun., 2010.
13. Mueter, F.J., and Litzow, M.A., 2008, Sea ice retreat alters the biogeography of the Bering Sea continental shelf: *Ecological Applications*, v. 18, no. 2, p. 309–320.
14. Froese, R., and Pauly, D., eds., 2012, *FishBase—Global information system on fishes*: FishBase database, accessed July 8, 2012, at <http://www.fishbase.org>.

Arctic Eelpout (*Lycodes reticulatus*)

Reinhardt, 1835

Family Zoarcidae

Note: Morphological differences between *L. reticulatus* (Reinhardt, 1835) and *L. rossi* are not clear, making identifications and geographic ranges uncertain [1].

Colloquial Name: None within U.S. Chukchi and Beaufort Seas.

Ecological Role: Because of the lack of taxonomic clarity, uncommon occurrence, and paucity of life history information, the ecological role of this species was not evaluated.

Physical Description/Attributes: Brown body with dark bands that are reticulate in large individuals; light spots on upper side of head. For specific diagnostic characteristics, see *Fishes of Alaska* (Mecklenburg and others, 2002, p. 713) [2]. Swim bladder: Absent [2]. Antifreeze glycoproteins in blood serum: Unknown.

Range: U.S. Beaufort Sea [3]. Elsewhere, Arctic Canada to Greenland and east to Barents, Kara, and Laptev Seas [2].

Relative Abundance: Apparently common in U.S. Beaufort Sea. Abundance estimates are unreliable because of lack of sampling in the offshore waters, and the species is likely common, at least locally, considering the 2012 archived voucher specimens from at least five stations [3]. Common in Barents Sea [5].



Arctic Eelpout (*Lycodes reticulatus*), 306 mm, Beaufort Sea, 2011. Photograph by C.W. Mecklenburg, Point Stephens Research.




Geographic distribution of Arctic Eelpout (*Lycodes reticulatus*) within Arctic Outer Continental Shelf Planning Areas [4], based on review of published literature and specimens from historical and recent collections [1-3].


Depth Range: Taken offshore at 6–930 m in U.S. Beaufort Sea [3]. 20–930 m, usually 380 m or shallower [1, 2]. Generally, eelpout spawning and larvae occur at same depths that adults inhabit [2].

Lycodes reticulatus
Arctic Eelpout


Benthic distribution



Reproductive distribution



Benthic and reproductive distribution of Arctic Eelpout (*Lycodes reticulatus*).



Habitats and Life History

Eggs—Size: Unknown. Time to hatching: Unknown. Habitat: Benthic [2].


Larvae—Size at hatching: Unknown. Size at juvenile transformation: Unknown. Days to juvenile transformation: Unknown. Habitat: Benthic [2].

Juveniles—Age and size: Unknown. Habitat: Benthic [2].

Adults—Age and size at first maturity: Unknown. Maximum age: 19 years [6]. Maximum size: 76.0 cm [2]. Habitat: Benthic, most often on outer shelf [1].

Substrate—Sand to mud [2].

Physical/chemical—Temperature: -1.5–4.5 °C [7]. Salinity: Marine [2].



Behavior

Diel—Unknown.

Seasonal—Unknown.

Reproductive—Unknown.

Schooling—Unknown.

Feeding—Unknown.



Populations or Stocks

There have been no studies.



Reproduction

Mode—Separate sexes. Oviparous [8].

Spawning season—Unknown.


Fecundity—Unknown.



Food and Feeding

Food items—A wide variety of benthic invertebrates (amphipods, bivalves, brittle stars, gastropods, echinoderms, crustaceans, and polychaetes) and fishes and pelagic crustaceans (euphausiids) [6, 9–11].


Trophic level— 3.5 ± 0.53 standard error [12].



Biological Interactions

Predators—Bearded seals in northeastern Canada [13].

Competitors—Likely other benthic microcarnivores including some sculpins, flatfishes, snailfishes, and other eelpouts.



Resilience

Medium, minimum population doubling time: 1.4–4.4 years (Preliminary *K* or Fecundity) [12].



Traditional and Cultural Importance

None reported.



Commercial Fisheries

Currently, Arctic Eelpout are not commercially harvested.



Potential Effects of Climate Change

Arctic Eelpouts have fairly wide depth and temperature tolerances. Thus, it is difficult to speculate about the effects of climate warming. Changes in temperature, species composition of fish assemblages, and productivity can be expected to affect distribution and abundance patterns



Areas for Future Research [B]

Little is known about this species biology and ecology from the region.

Research needs include:

(1) Depth and location of pelagic larvae; (2) depth, location, and timing of young-of-the-year benthic recruitment; (3) preferred depth ranges for juveniles and adults; (4) spawning season; (5) seasonal and ontogenetic movements; (6) population studies; (7) prey; and (8) predators.

References Cited

Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, *Fishes of Alaska*: Bethesda, Maryland, American Fisheries Society, 1,116 p. [2]

Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: *Marine Biodiversity*, v. 41, no. 1, p. 109–140, Online Resource 1. [1]

Bibliography

1. Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: *Marine Biodiversity*, v. 41, no. 1, p. 109–140, Online Resource 1.
2. Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, *Fishes of Alaska*: Bethesda, Maryland, American Fisheries Society, 1,116 p.
3. Mecklenburg, C.W., Mecklenburg, T.A., Sheiko, B.A., and Steinke, D., 2016, Pacific Arctic marine fishes: Akureyri, Iceland, Conservation of Arctic Flora and Fauna, Monitoring Series Report No. 23, 406 p., accessed May 10, 2016, at <http://caff.is/monitoring-series/370-pacific-arcticmarine-fishes>.
4. Minerals Management Service, 2008, Beaufort Sea and Chukchi Sea planning areas—Oil and Gas Lease Sales 209, 212, 217, and 221: U.S. Department of the Interior, Minerals Management Service Alaska OCS Region, OCS EIS/EA, MMS 2008-0055, 538 p.
5. Wienerroither, R., Johannesen, E., Langøy, H., Børve Eriksen, K., de Lange Wenneck, T., Høines, Å., Bjelland, O., Aglen, A., Prokhorova, T., Murashko, P., Prozorkevich, D., Konstantin, Byrkjedal, I., Langhelle Drevetnyak, and G., Smirnov, O., 2011, Atlas of the Barents Sea fishes: IMR/PINRO Joint Report Series 1-2011, ISSN 1502-8828, 274 p.
6. von Dorrien, C.F., 1993, Ecology and respiration of selected Arctic benthic fish species: *Berichte zur Polarforschung*, v. 125, 118 p. [In German]
7. Møller, P.R., and Jørgensen, O.A., 2000, Distribution and abundance of eelpouts (Pisces, Zoarcidae) off West Greenland: *Sarsia*, v. 85, no. 1, p. 23–48.
8. Love, M.S., 2011, *Certainly more than you wanted to know about the fishes of the Pacific Coast*: Santa Barbara, California, Really Big Press, 649 p.
9. Chambers, C.A., and Dick, T.A., 2005, Trophic structure of one deep-sea fish community in the eastern Canadian Arctic—Application of food, parasites and multivariate analysis: *Environmental Biology of Fishes*, v. 74, nos. 3–4, p. 365–378.
10. Torres, P., Rodriguez-Marin, E., and Loureiro, I., 2000, Preliminary results from feeding analysis for the most abundant demersal fishes in Flemish Cap during summer (1993–2000): Northwest Atlantic Fisheries Organization, Serial No. N4302, NAFO SCR Doc. 00/60.
11. Román, E., González, C., and Ceballos, E., 2004, Food and feeding of most abundant fish species in Flemish Cap: Northwest Atlantic Fisheries Organization, Serial No. N5018, NAFO SCR Doc. 04/58.
12. Froese, R., and Pauly, D., eds., 2012, *FishBase—Global information system on fishes*: FishBase database, accessed July 8, 2012, at <http://www.fishbase.org>.
13. Dunbar, M.J., and Hildebrand, H.H., 1952, Contributions to the study of the fishes of Ungava Bay: *Journal of the Fisheries Research Board of Canada*, v. 9, no. 2, p. 83–126.

Threespot Eelpout (*Lycodes rossi*)

Malmgren, 1865

Family Zoarcidae

Note: Except for geographic range data, all information is from areas outside of the study area.

Note on taxonomy: Morphological differences between *L. reticulatus* (Reinhardt, 1835) and *L. rossi* are not clear, making identifications and geographic ranges uncertain [1].

Colloquial Name: None within U.S. Chukchi and Beaufort Seas.

Ecological Role: Uncertain. The life history and ecology of this species and its roles in regional ecosystems and food webs are poorly understood. Issues regarding the taxonomy of this species need to be resolved.

Physical Description/Attributes: Brown, with dark brown bands on body and dorsal fin; light band across top of head connecting gill openings, often broken into spots. For specific diagnostic characteristics, see *Fishes of Alaska* (Mecklenburg and others, 2002, p. 713) [1]. Swim bladder: Absent [1]. Antifreeze glycoproteins in blood serum: Unknown.

Range: U.S. Beaufort Sea, at least between 152° and 155°W [2]. Elsewhere, from Greenland and Norwegian Seas to Kara Sea, including Iceland, Svalbard (Norway), and Barents Sea; and Canadian Beaufort Sea eastward to Dease Strait [1, 2].

Relative Abundance: Rare in U.S. Beaufort Sea [2, 4]. Common in Barents Sea [5].




Threespot Eelpout (*Lycodes rossi*), Beaufort Sea, 2008. Photograph by E. Akuna, National Oceanic and Atmospheric Administration, Alaska Fisheries Science Center.




Geographic distribution of Threespot Eelpout (*Lycodes rossi*), within Arctic Outer Continental Shelf Planning Areas [3] based on review of published literature and specimens from historical and recent collections [1, 2, 4].

Depth Range: Adults at 42–365 m, usually deeper than 130 m [1, 2, 6]. Juveniles as shallow as 9 m [1, 6]. Generally, eelpout spawning and larvae occur at same depths that adults inhabit [1].

Lycodes rossii
Threespot Eelpout



Benthic and reproductive distribution of Threespot Eelpout (*Lycodes rossii*).



Habitats and Life History

Eggs—Size: 3–4 mm [5]. Time to hatching: Unknown. Habitat: Benthic [1].


Larvae—Size at hatching: Unknown. Size at juvenile transformation: Unknown. Days to juvenile transformation: Unknown. Habitat: Benthic [1].

Juveniles—Age and size: Unknown. Habitat: Benthic [1].

Adults—Age and size at first maturity: Unknown. Maximum age: 10 years [5]. Maximum size: 38 cm TL [7]. Habitat: Benthic, most often on outer shelf and upper slope [2].

Substrate—Muddy [1, 8].

Physical/chemical—Temperature: -1.5–1.1 °C [8]. Found mainly at minus temperatures [6, 8]. Salinity: 27.6–35.0 ppt [8]; prefers high salinity, usually not found in less than 34 ppt [6].



Behavior

Diel—Unknown.

Seasonal—Unknown.

Reproductive—Unknown.


Schooling—Unknown.

Feeding—Unknown.



Populations or Stocks

There have been no studies.



Reproduction

Mode—Separate sexes. Oviparous [9].

Spawning season—Probably winter or early spring in Barents Sea [5].


Fecundity—390 eggs [5].



Food and Feeding

Food items—Polychaetes, copepods, amphipods, cumaceans [6, 8].


Trophic level— 3.49 ± 0.53 standard error [10].



Biological Interactions

Predators—Bearded seals in Ungava Bay, Canada [11].

Competitors—Likely other benthic microcarnivores including some sculpins, flatfishes, snailfishes, and other eelpouts.



Resilience

Medium, minimum population doubling time: 1.4–4.4 years (Preliminary *K* or Fecundity) [10].



Traditional and Cultural Importance

None reported.



Commercial Fisheries

Currently, Threespot Eelpout are not commercially harvested.



Potential Effects of Climate Change

Because Threespot Eelpout are mainly a deep-water, slope-dwelling Arctic species and little is known about climate change at slope depths in the Arctic, potential effects on this species cannot be estimated.



Areas for Future Research [B]

Little is known about this species biology and ecology from the region. Research needs include: (1) depth and location of pelagic larvae, (2) depth, location, and timing of young-of-the-year benthic recruitment, (3) preferred depth ranges for juveniles and adults, (4) spawning season, (5) seasonal and ontogenetic movements, (6) population studies, (7) prey, and (8) predators.

References Cited


- Andriashev, A.P., 1954, Fishes of the northern seas of the U.S.S.R.—Keys to the fauna of the U.S.S.R.: Academy of Sciences of the U.S.S.R., Zoological Institute, no. 53, 566 p. [In Russian, translation by Israel Program for Scientific Translation, Jerusalem, 1964, 617 p., available from U.S. Department of Commerce, Springfield, Virginia.] [6]
- McAllister, D.E., Anderson, M.E., and Hunter, J.G., 1981, Deep-water eelpouts, Zoarcidae, from Arctic Canada and Alaska: Canadian Journal of Fisheries and Aquatic Sciences, v. 38, no. 7, p. 821–839. [8]
- Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, Fishes of Alaska: Bethesda, Maryland, American Fisheries Society, 1,116 p. [1]
- Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: Marine Biodiversity, v. 41, no. 1, p. 109–140, Online Resource 1. [2]

Bibliography

1. Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, Fishes of Alaska: Bethesda, Maryland, American Fisheries Society, 1,116 p.
2. Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: Marine Biodiversity, v. 41, no. 1, p. 109–140, Online Resource 1.
3. Minerals Management Service, 2008, Beaufort Sea and Chukchi Sea planning areas—Oil and Gas Lease Sales 209, 212, 217, and 221: U.S. Department of the Interior, Minerals Management Service Alaska OCS Region, OCS EIS/EA, MMS 2008-0055, 538 p.
4. Mecklenburg, C.W., Mecklenburg, T.A., Sheiko, B.A., and Steinke, D., 2016, Pacific Arctic marine fishes: Akureyri, Iceland, Conservation of Arctic Flora and Fauna, Monitoring Series Report No. 23, 406 p., accessed May 10, 2016, at <http://caff.is/monitoring-series/370-pacific-arcticmarine-fishes>.
5. Wienerroither, R., Johannesen, E., Langøy, H., Børve Eriksen, K., de Lange Wenneck, T., Høines, Å., Bjelland, O., Aglen, A., Prokhorova, T., Murashko, P., Prozorkevich, D., Konstantin, Byrkjedal, I., Langhelle Drevetnyak, and G., Smirnov, O., 2011, Atlas of the Barents Sea fishes: IMR/PINRO Joint Report Series 1-2011, ISSN 1502-8828, 274 p.
6. Andriashev, A.P., 1954, Fishes of the northern seas of the U.S.S.R.—Keys to the fauna of the U.S.S.R.: Academy of Sciences of the U.S.S.R., Zoological Institute, no. 53, 566 p. [In Russian, translation by Israel Program for Scientific Translation, Jerusalem, 1964, 617 p., available from U.S. Department of Commerce, Springfield, Virginia.]
7. Coad, B.W., and Reist, J.D., 2004, Annotated list of the Arctic marine fishes of Canada: Canadian Manuscript Report of Fisheries and Aquatic Sciences, Fisheries and Oceans Canada, no. 2674, 112 p.
8. McAllister, D.E., Anderson, M.E., and Hunter, J.G., 1981, Deep-water eelpouts, Zoarcidae, from Arctic Canada and Alaska: Canadian Journal of Fisheries and Aquatic Sciences, v. 38, no. 7, p. 821–839.
9. Love, M.S., 2011, Certainly more than you wanted to know about the fishes of the Pacific Coast: Santa Barbara, California, Really Big Press, 649 p.
10. Froese, R., and Pauly, D., eds., 2012, FishBase—Global information system on fishes: FishBase database, accessed July 8, 2012, at <http://www.fishbase.org>.
11. Dunbar, M.J., and Hildebrand, H.H., 1952, Contributions to the study of the fishes of Ungava Bay: Journal of the Fisheries Research Board of Canada, v. 9, no. 2, p. 83–126.

Archer Eelpout (*Lycodes sagittarius*)


McAllister, 1976

Family Zoarcidae**Colloquial Name:** *None within U.S. Chukchi and Beaufort Seas.***Ecological Role:** Likely to be absent over shelf habitats of the U.S. Beaufort Sea. Survey data suggest that this species rarely occurs in slope and deep water habitats. The ecological role of the species is probably minimal.**Physical Description/Attributes:** Elongate, narrow body colored uniformly dark brown; peritoneum black. For specific diagnostic characteristics see *Fishes of Alaska* (Mecklenburg and others, 2002, p. 717) [1]; specimens with white bars and fewer vertebrae illustrated therein [1] and in previous publications, (for example, McAllister and others, 1981) [2], are now known to belong to *L. marisalbi* [3]. Swim bladder: Absent [1]. Antifreeze glycoproteins in blood serum: Unknown.**Range:** *U.S. Beaufort Sea, and Beaufort Sea of western Canada and Kara and Laptev Seas [5].***Relative Abundance:** *Apprently common in Beaufort Sea off Alaska [5].*Archer Eelpout (*Lycodes sagittarius*), 180 mm TL or less from Franklin Bay, Northwest Territories, Canada, (from Mecklenburg and others 2002, p. 717).


Geographic distribution of Archer Eelpout (*Lycodes sagittarius*), within Arctic Outer Continental Shelf Planning Areas [4] based on review of published literature and specimens from historical and recent collections [3, 5].

Depth Range: 357–600 m in U.S. Beaufort Sea [2, 3] otherwise depths of 120 to at least 1,934 m [5]. Generally, eelpout spawning and larvae occur at same depths that adults inhabit [1].

Lycodes sagittarius
Archer Eelpout



Benthic and reproductive distribution of Archer Eelpout (*Lycodes sagittarius*).



Habitats and Life History

Eggs—Size: Unknown. Time to hatching: Unknown. Habitat: Benthic [1].


Larvae—Size at hatching: Unknown. Size at juvenile transformation: Unknown. Days to juvenile transformation: Unknown. Habitat: Benthic [1].

Juveniles—Age and size: Unknown. Habitat: Benthic [1].

Adults—Age and size at first maturity: Unknown. Maximum age: Unknown. Maximum size: To 27.8 cm TL [2]. Habitat: Benthic [1].

Substrate—Mud [2].

Physical/chemical—Temperature: Range unknown. Taken at -0.9°C [2]. Salinity: Marine, taken at 34 ppt. [2].



Behavior

Diel—Unknown.

Seasonal—Unknown.

Reproductive—Unknown.

Schooling—Unknown.

Feeding—Unknown.



Populations or Stocks

There have been no studies.



Reproduction

Mode—Separate sexes, oviparous.

Spawning season—Likely late summer or early autumn [2].


Fecundity—Unknown.



Food and Feeding

Food items—Annelids, bivalves, gastropods, and crustaceans [2].


Trophic level—3.22 ±0.40 standard error [6].



Biological Interactions

Predators—Unknown.

Competitors—Likely other benthic microcarnivorous fishes, including snailfishes, sculpins, flatfishes, and other eelpouts.



Resilience

Medium, minimum population doubling time: 1.4–4.4 years (Preliminary *K* or Fecundity) [6].



Traditional and Cultural Importance

None reported.




Commercial Fisheries

Currently, Archer Eelpout are not commercially harvested.



Potential Effects of Climate Change

Unknown. The Archer Eelpout is an endemic species occurring rarely in deep water areas of the U.S. Beaufort Sea [3]. Lack of information about this species and understanding of its role in this benthic ecosystem precludes an informed assessment of potential climatic effects.



Areas for Future Research [B]

Little is known about the biology and ecology of this species from the region. Research needs include:

- (1) depth and location of pelagic larvae, (2) depth, location, and timing of young-of-the-year benthic recruitment, (3) preferred depth ranges for juveniles and adults, (4) spawning season, (5) seasonal and ontogenetic movements, (6) population studies, (7) prey, and (8) predators.

Bibliography

1. Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, *Fishes of Alaska*: Bethesda, Maryland, American Fisheries Society, 1,116 p.
2. McAllister, D.E., Anderson, M.E., and Hunter, J.G., 1981, Deep-water eelpouts, Zoarcidae, from Arctic Canada and Alaska: *Canadian Journal of Fisheries and Aquatic Sciences*, v. 38, no. 7, p. 821–839.
3. Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: *Marine Biodiversity*, v. 41, no. 1, p. 109–140, Online Resource 1.
4. Minerals Management Service, 2008, Beaufort Sea and Chukchi Sea planning areas—Oil and Gas Lease Sales 209, 212, 217, and 221: U.S. Department of the Interior, Minerals Management Service Alaska OCS Region, OCS EIS/EA, MMS 2008-0055, 538 p.
5. Mecklenburg, C.W., Mecklenburg, T.A., Sheiko, B.A., and Steinke, D., 2016, Pacific Arctic marine fishes: Akureyri, Iceland, Conservation of Arctic Flora and Fauna, Monitoring Series Report No. 23, 406 p., accessed May 10, 2016, at <http://caff.is/monitoring-series/370-pacific-arcticmarine-fishes>.
6. Froese, R., and Pauly, D., eds., 2012, FishBase—Global information system on fishes: FishBase database, accessed July 8, 2012, at <http://www.fishbase.org>.

Longear Eelpout (*Lycodes seminudus*)

Reinhardt, 1837

Family Zoarcidae

Note: Except for geographic range data, all information is from areas outside of the study area.

Colloquial Name: None within U.S. Chukchi and Beaufort Seas.

Ecological Role: Because of its apparent rarity in shallow U.S. Arctic waters, this species is likely of little ecological importance in the U.S. Chukchi and Beaufort Sea.


Physical Description/Attributes: Elongate, dark gray-brown body, sometimes with short, pale bands above lateral line and extending onto dorsal fin. For specific diagnostic characteristics, see *Fishes of Alaska* (Mecklenburg and others, 2002, p. 707) [1]. Swim bladder: Absent [1]. Antifreeze glycoproteins in blood serum: Unknown.

Range: Continental slope of U.S. Chukchi and Beaufort Seas [2]. Elsewhere, in Baffin Bay off Canada and northwest Greenland, northeast Greenland, off northern Iceland to Faroe Islands, Svalbard Archipelago, northern Barents Sea, and Kara Sea [2, 3].

Relative Abundance: Rare in shelf waters of U.S. Chukchi and Beaufort Seas, but potentially more common in deeper waters over the continental slope [2, 5]. Common off Greenland [2].



Longear Eelpout (*Lycodes seminudus*), 238 mm TL, Chukchi Borderland, 2009. Photograph by C.W. Mecklenburg, Point Stephens Research.




Geographic distribution of Longear Eelpout (*Lycodes seminudus*) within Arctic Outer Continental Shelf Planning Areas [4] based on review of published literature and specimens from historical and recent collections [1, 2, 5].


Depth Range: 50–1,400 m, mostly 200–600 m [2, 6]. In general, eelpout spawning and larvae occur at same depths that adults inhabit [1].

Lycodes seminudus
Longear Eelpout


Benthic distribution



Reproductive distribution



Benthic and reproductive distribution of Longear Eelpout (*Lycodes seminudus*).



Habitats and Life History

Eggs—Time to hatching: Unknown. Size: Unknown. Habitat: Benthic [1].


Larvae—Size at hatching: Unknown. Size at juvenile transformation: Unknown. Days to juvenile transformation: Unknown. Habitat: Benthic [1].

Juveniles—Age and size: Unknown. Habitat: Benthic [1].

Adults—Age and size at first maturity: Unknown. Maximum age: Unknown. Maximum size: 56 cm TL [3]. Habitat: Benthic [1].

Substrate—Mud or mud-clay [7].

Physical/chemical—Temperature: -1.7–4.9 °C [5]. In Russia, almost exclusively at less than 0 °C [6]. Salinity: Marine, high salinity [6].



Behavior

Diel—Unknown.

Seasonal—Unknown.

Reproductive—Unknown.

Schooling—Unknown.

Feeding—Unknown.



Populations or Stocks

There have been no studies.



Reproduction

Mode—Separate sexes, oviparous.


Spawning season—Unknown.

Fecundity—Unknown.




Food and Feeding

Food items—Amphipods, shrimps, isopods, polychaetes [6].
Trophic level—3.45 standard error 0.44 [8].



Biological Interactions

Predators—Unknown.
Competitors—Likely other benthic microcarnivores, including flatfishes, snailfishes, and other eelpouts.



Resilience

Low, minimum population doubling time: 4.5–14 years (Preliminary *K* or Fecundity) [8].



Traditional and Cultural Importance

None reported.




Commercial Fisheries

Currently, Longear Eelpout are not commercially harvested.



Potential Effects of Climate Change

Unknown. The Longear Eelpout is an endemic Arctic species occurring most commonly in deep water habitats over the slope [2]. Not enough is known about this species to predict climate change effects.



Areas for Future Research [B]

Little is known about biology and ecology of this species from the region. Research needs include: (1) depth and location of pelagic larvae, (2) depth, location, and timing of young-of-the-year benthic recruitment, (3) preferred depth ranges for juveniles and adults, (4) spawning season, (5) seasonal and ontogenetic movements, (6) population studies, (7) prey, and (8) predators.

References Cited

- Andriashev, A.P., 1954, Fishes of the northern seas of the U.S.S.R.—Keys to the fauna of the U.S.S.R.: Academy of Sciences of the U.S.S.R., Zoological Institute, no. 53, 566 p. [In Russian, translation by Israel Program for Scientific Translation, Jerusalem, 1964, 617 p., available from U.S. Department of Commerce, Springfield, Virginia.] [6]
- Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, Fishes of Alaska: Bethesda, Maryland, American Fisheries Society, 1,116 p. [1]
- Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: Marine Biodiversity, v. 41, no. 1, p. 109–140, Online Resource 1. [2]

Bibliography

1. Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, Fishes of Alaska: Bethesda, Maryland, American Fisheries Society, 1,116 p.
2. Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: Marine Biodiversity, v. 41, no. 1, p. 109–140, Online Resource 1.
3. Møller, P.R., and Jørgensen, O.A., 2000, Distribution and abundance of eelpouts (Pisces, Zoarcidae) off West Greenland: Sarsia, v. 85, no. 1, p. 23–48.
4. Minerals Management Service, 2008, Beaufort Sea and Chukchi Sea planning areas—Oil and Gas Lease Sales 209, 212, 217, and 221: U.S. Department of the Interior, Minerals Management Service Alaska OCS Region, OCS EIS/EA, MMS 2008-0055, 538 p.
5. Mecklenburg, C.W., Mecklenburg, T.A., Sheiko, B.A., and Steinke, D., 2016, Pacific Arctic marine fishes: Akureyri, Iceland, Conservation of Arctic Flora and Fauna, Monitoring Series Report No. 23, 406 p., accessed May 10, 2016, at <http://caff.is/monitoring-series/370-pacific-arcticmarine-fishes>.
6. Andriashev, A.P., 1954, Fishes of the northern seas of the U.S.S.R.—Keys to the fauna of the U.S.S.R.: Academy of Sciences of the U.S.S.R., Zoological Institute, no. 53, 566 p. [In Russian, translation by Israel Program for Scientific Translation, Jerusalem, 1964, 617 p., available from U.S. Department of Commerce, Springfield, Virginia.]
7. McAllister, D.E., Anderson, M.E., and Hunter, J.G., 1981, Deep-water eelpouts, Zoarcidae, from Arctic Canada and Alaska: Canadian Journal of Fisheries and Aquatic Sciences, v. 38, no. 7, p. 821–839.
8. Froese, R., and Pauly, D., eds., 2012, FishBase—Global information system on fishes: FishBase database, accessed July 8, 2012, at <http://www.fishbase.org>.

Scalebelly Eelpout (*Lycodes squamiventer*)

Jensen, 1904

Family Zoarcidae

Note: Except for geographic range data, all information is from areas outside of the study area.

Colloquial Name: None within U.S. Chukchi and Beaufort Seas.

Ecological Role: This species generally does not occur on the continental shelves in the Arctic Ocean but likely has some ecological significance in the relatively unexplored, deep waters of the U.S. Chukchi and Beaufort Seas.

Physical Description/Attributes: Light to dark grayish brown body without bands or other marks. For specific diagnostic characteristics, see *Fishes of Alaska* (Mecklenburg and others, 2002, p. 722)[1]. Swim bladder: Absent [1]. Antifreeze glycoproteins in blood serum: Unknown.

Range: U.S. Beaufort Sea [2]. Elsewhere, Canadian Beaufort Sea, Davis Strait off western Greenland; Greenland Sea, Norwegian Sea, Kara Sea, and off Faeroese-Shetland slope, Barents Sea [1, 3].

Relative Abundance: Common in U.S. Beaufort Sea and western Canada Beaufort Sea. Although known from only a few records, this species could be more common on the slope as has been reported elsewhere in the Arctic [2]. Common in Norwegian Sea and around Håkon Mosby Mud Volcano in Barents Sea [5].




Scalebelly Eelpout (*Lycodes squamiventer*), 358 mm, Beaufort Sea, 2012. Photograph by C.W. Mecklenburg, Point Stephens Research.




Geographic distribution of Scalebelly Eelpout (*Lycodes squamiventer*) within Arctic Outer Continental Shelf Planning Areas [4] based on review of literature and specimens from historical and recent collections [1-3].

Depth Range: Benthic, 160–1,808 m [2], typically from 1,273–1,546 m in Norwegian Sea [5]. Reported but not confirmed as shallow as 160 m [6]. *Records from U.S. Beaufort Sea are depths of 357–500 m* [2]. In general, eelpout spawning and larvae occur at the same depths that adults inhabit [7].

Lycodes squamiventer
Scalebelly Eelpout



Benthic and reproductive distribution of Scalebelly Eelpout (*Lycodes squamiventer*).



Habitats and Life History


Eggs—Size: 3.5 mm [5]. Time to hatching: Unknown. Habitat: Benthic [1, 5].

Larvae—Size at hatching: Unknown. Size at juvenile transformation: Unknown. Days to juvenile transformation: Unknown. Habitat: Benthic [1].

Juveniles—Age and size: Unknown. Habitat: Benthic [1].

Adults—Age and size at first maturity: Age unknown. Females at 14 cm, males at 17.9 cm [6]. Maximum age: 21 years [5]. Maximum size: 37 cm TL [2]. Habitat: Benthic, in deep waters on the slope [3, 5, 6]. Substrate—Muddy bottoms [1].

Physical/chemical—Temperature: -1.2–0.6 °C [5]. Salinity: Marine.



Behavior

Diel—Unknown.

Seasonal—Unknown.

Reproductive—Unknown.

Schooling—Unknown.

Feeding—Unknown.



Populations or Stocks

There have been no studies.



Reproduction

Mode—Separate sexes. Oviparous [8].

Spawning season—Autumn, but ripe females have been found in June [5, 6].


Fecundity—50–60 eggs [5, 6].



Food and Feeding

Food items—Various benthic species such as pogonophores, gastropods, amphipods, polychaetes, copepods, ophiuroids, bivalves, and crustaceans [5, 6].


Trophic level— 3.4 ± 0.4 standard error [9].



Biological Interactions

Predators—Unknown.

Competitors—Likely other benthic microcarnivores including some sculpins, flatfishes, snailfishes, and other eelpouts.



Resilience

Medium, minimum population doubling time: 1.4–4.4 years (Preliminary *K* or Fecundity) [9].



Traditional and Cultural Importance

None reported.




Commercial Fisheries

Currently, Scalebelly Eelpout are not commercially harvested.



Potential Effects of Climate Change

Unknown. The Scalebelly Eelpout is a predominantly Arctic, slope and deep-water species. Not enough information is available to evaluate potential climatic effects.



Areas for Future Research [B]

Little is known about the biology and ecology of this species from the region. Research needs include:

- (1) depth and location of pelagic larvae, (2) depth, location, and timing of young-of-the-year benthic recruitment, (3) preferred depth ranges for juveniles and adults, (4) spawning season, (5) seasonal and ontogenetic movements, (6) population studies, (7) prey, and (8) predators.

References Cited

- Hildebrandt, N., Bergmann, M., and Knust, R., 2011, Longevity and growth efficiency of two deep-dwelling Arctic zoarcids and comparison with eight other zoarcid species from different climatic regions: *Polar Biology*, v. 34, no. 10, p. 1,523–1,533. [5]
- Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, *Fishes of Alaska*: Bethesda, Maryland, American Fisheries Society, 1,116 p. [1]
- Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: *Marine Biodiversity*, v. 41, no. 1, p. 109–140, Online Resource 1. [3]
- Wienerroither, R., Johannesen, E., Langøy, H., Børve Eriksen, K., de Lange Wenneck, T., Høines, Å., Bjelland, O., Aglen, A., Prokhorova, T., Murashko, P., Prozorkevich, D., Konstantin, Byrkjedal, I., Langhelle Drevetnyak, and G., Smirnov, O., 2011, *Atlas of the Barents Sea fishes: IMR/PINRO Joint Report Series 1-2011*, ISSN 1502-8828, 274 p. [6]

Bibliography

1. Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, *Fishes of Alaska*: Bethesda, Maryland, American Fisheries Society, 1,116 p.
2. Mecklenburg, C.W., Mecklenburg, T.A., Sheiko, B.A., and Steinke, D., 2016, *Pacific Arctic marine fishes: Akureyri, Iceland, Conservation of Arctic Flora and Fauna, Monitoring Series Report No. 23*, 406 p., accessed May 10, 2016, at <http://caff.is/monitoring-series/370-pacific-arcticmarine-fishes>.
3. Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: *Marine Biodiversity*, v. 41, no. 1, p. 109–140, Online Resource 1.
4. Minerals Management Service, 2008, *Beaufort Sea and Chukchi Sea planning areas—Oil and Gas Lease Sales 209, 212, 217, and 221*: U.S. Department of the Interior, Minerals Management Service Alaska OCS Region, OCS EIS/EA, MMS 2008-0055, 538 p.
5. Hildebrandt, N., Bergmann, M., and Knust, R., 2011, Longevity and growth efficiency of two deep-dwelling Arctic zoarcids and comparison with eight other zoarcid species from different climatic regions: *Polar Biology*, v. 34, no. 10, p. 1,523–1,533.
6. Wienerroither, R., Johannesen, E., Langøy, H., Børve Eriksen, K., de Lange Wenneck, T., Høines, Å., Bjelland, O., Aglen, A., Prokhorova, T., Murashko, P., Prozorkevich, D., Konstantin, Byrkjedal, I., Langhelle Drevetnyak, and G., Smirnov, O., 2011, *Atlas of the Barents Sea fishes: IMR/PINRO Joint Report Series 1-2011*, ISSN 1502-8828, 274 p.
7. Andriashev, A.P., 1954, *Fishes of the northern seas of the U.S.S.R.—Keys to the fauna of the U.S.S.R.*: Academy of Sciences of the U.S.S.R., Zoological Institute, no. 53, 566 p. [In Russian, translation by Israel Program for Scientific Translation, Jerusalem, 1964, 617 p., available from U.S. Department of Commerce, Springfield, Virginia.]
8. Love, M.S., 2011, *Certainly more than you wanted to know about the fishes of the Pacific Coast*: Santa Barbara, California, Really Big Press, 649 p.
9. Froese, R., and Pauly, D., eds., 2012, *FishBase—Global information system on fishes*: FishBase database, accessed July 8, 2012, at <http://www.fishbase.org>.

Estuarine Eelpout (*Lycodes turneri*)

Bean, 1879

Family Zoarcidae

Colloquial Name: *None within U.S. Chukchi and Beaufort Seas.*

Ecological Role: Unknown. Could be important in coastal food webs as this eelpout could be prey for piscivorous fishes and birds of the nearshore community.

Physical Description/Attributes: Elongate, body with 10–12 bands. Color in adults is purple with bluish white bands bordered with purplish olive, or umber with creamy white bands and dark umber borders. Juveniles are creamy white with blackish bordered brown bands. For specific diagnostic characteristics, see *Fishes of Alaska* (Mecklenburg and others, 2002, p. 709) [1]. Swim bladder: Absent [1]. Antifreeze glycoproteins in blood serum: Unknown.

Range: *U.S. Chukchi Sea and western U.S. Beaufort Sea in vicinity of Point Barrow.* Elsewhere in Alaska, southward in eastern Bering Sea to Bristol Bay. Worldwide, southward in western Bering Sea to Cape Olyutorskiy, Russia [2].

Relative Abundance: *Uncommon in U.S. Chukchi and Beaufort Seas* [2, 5, 6]. *A record from the eastern U.S. Beaufort Sea off Point Franklin was recently shown to be in error because of a mistake in a museum catalog; the fish were actually captured in the Chukchi Sea* [1].



Estuarine Eelpout (*Lycodes turneri*), juvenile, 145 mm TL, Chukchi Sea, 2009. Photograph by C.W. Mecklenburg, Point Stephens Research. Fish shown is a juvenile.




Base modified from USGS and other digital data. U.S.-Russia Maritime Boundary follows the EEZ/200-mile limit line, western edge. Coordinate reference system: projection, Lambert Azimuthal Equal Area; latitude of origin, 75.0°; horizontal datum, North American Datum of 1983.

Geographic distribution of Estuarine Eelpout (*Lycodes turneri*) within Arctic Outer Continental Shelf Planning Areas [3] based on review of literature and specimens from historical and recent collections [1, 2, 4].


Depth Range: 1–125 m, typically less than 50 m [2]. Generally, eelpout spawning and larvae occur at same depths that adults inhabit [1].

Lycodes turneri
Estuarine Eelpout


Benthic distribution



Reproductive distribution



Benthic and reproductive distribution of Estuarine Eelpout (*Lycodes turneri*).



Habitats and Life History

Eggs—Size: Unknown. Time to hatching: Unknown. Habitat: Benthic [1].


Larvae—Size at hatching: Unknown. Size at juvenile transformation: Unknown. Days to juvenile transformation: Unknown. Habitat: Benthic [1].

Juveniles—Age and size: Unknown. Habitat: Benthic [1].

Adults—Age and size at first maturity: Unknown. Maximum age: Unknown. Maximum size: 64 cm TL [7]. Habitat: Benthic [1], limited to inner and mid-shelf [5].

Substrate—Soft bottoms [1].

Physical/chemical—Temperature: Not reported. Salinity: Taken mostly in or near estuaries [4].



Behavior

Diel—Unknown.

Seasonal—Unknown.

Reproductive—Unknown.

Schooling—Unknown.

Feeding—Unknown.



Populations or Stocks

There have been no studies.



Reproduction

Mode—Unknown.


Spawning season—Unknown.

Fecundity—Unknown.




Food and Feeding

Food items—Unknown.
Trophic level—3.36 standard error 0.44 [8]



Biological Interactions

Predators—Unknown.
Competitors—Unknown.



Resilience

Medium, minimum population doubling time: 1.4–4.4 years (Preliminary *K* or Fecundity) [8].




Traditional and Cultural Importance

Reported in the late 1800s to be an important food fish at Saint Michael, a village on the coast of Norton Sound in the Alaskan Bering Sea [9].



Commercial Fisheries

Currently, Estuarine Eelpout are not commercially harvested.



Potential Effects of Climate Change

Unknown. Range expansions are possible with expansion of brackish water conditions.



Areas for Future Research [B] Little is known about the biology and ecology of this species from the region.

Research needs include: (1) depth and location of pelagic larvae, (2) depth, location, and timing of young-of-the-year benthic recruitment, (3) preferred depth ranges for juveniles and adults, (4) spawning season, (5) seasonal and ontogenetic movements, (6) population studies, (7) prey, and (8) predators.

References Cited

- Allen, M.J., and Smith, G.B., 1988, Atlas and zoogeography of common fishes in the Bering Sea and northeastern Pacific: National Oceanic and Atmospheric Administration Technical Report NMFS 66, 151 p. [5].
- Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, Fishes of Alaska: Bethesda, Maryland, American Fisheries Society, 1,116 p. [1]
- Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: Marine Biodiversity, v. 41, no. 1, p. 109–140, Online Resource 1. [2]

Bibliography

1. Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, Fishes of Alaska: Bethesda, Maryland, American Fisheries Society, 1,116 p.
2. Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: Marine Biodiversity, v. 41, no. 1, p. 109–140, Online Resource 1.
3. Minerals Management Service, 2008, Beaufort Sea and Chukchi Sea planning areas—Oil and Gas Lease Sales 209, 212, 217, and 221: U.S. Department of the Interior, Minerals Management Service Alaska OCS Region, OCS EIS/EA, MMS 2008-0055, 538 p.
4. Mecklenburg, C.W., Mecklenburg, T.A., Sheiko, B.A., and Steinke, D., 2016, Pacific Arctic marine fishes: Akureyri, Iceland, Conservation of Arctic Flora and Fauna, Monitoring Series Report No. 23, 406 p., accessed May 10, 2016, at <http://caff.is/monitoring-series/370-pacific-arcticmarine-fishes>.
5. Allen, M.J., and Smith, G.B., 1988, Atlas and zoogeography of common fishes in the Bering Sea and northeastern Pacific: National Oceanic and Atmospheric Administration Technical Report NMFS 66, 151 p.
6. Barber, W.E., Smith, R.L., Vallarino, M., and Meyer, R.M., 1997, Demersal fish assemblages of the northeastern Chukchi Sea, Alaska: Fishery Bulletin, v. 95, no. 2, p. 195–209.
7. Andriashev, A.P., 1954, Fishes of the northern seas of the U.S.S.R.—Keys to the fauna of the U.S.S.R.: Academy of Sciences of the U.S.S.R., Zoological Institute, no. 53, 566 p. [In Russian, translation by Israel Program for Scientific Translation, Jerusalem, 1964, 617 p., available from U.S. Department of Commerce, Springfield, Virginia.]
8. Froese, R., and Pauly, D., eds., 2012, FishBase—Global information system on fishes: FishBase database, accessed July 8, 2012, at <http://www.fishbase.org>.
9. Bean, T.H., 1887, The fishery resources and fishing-grounds of Alaska, in Goode, G.B., ed., The fisheries and fishery industries of the United States, Section III: United States Commission of Fish and Fisheries, p. 81–115.