

**ARCTIC FISH HABITAT USE INVESTIGATIONS:  
NEARSHORE STUDIES IN THE ALASKAN BEAUFORT SEA,  
SUMMER 1990**

by

**Lyman K. Thorsteinson, Laurie E. Jarvela, and David A. Hale**

**National Oceanic and Atmospheric Administration  
National Ocean Service  
Office of Ocean Resources Conservation and Assessment  
Grace Hall, Suite 300  
4230 University Drive  
Anchorage, Alaska 99508-4626**

**Annual Report**

**September 1991**

## NOTICES

This document has been prepared by the Alaska Office of the National Ocean Service, Office of Ocean Resources Conservation and Assessment, National Oceanic and Atmospheric Administration (NOAA), U.S. Department of Commerce. The interpretation of data and opinions expressed in this document are those of the authors and do not necessarily reflect the views and policies of the Department of Commerce.

NOAA does not approve, recommend, or endorse any proprietary material mentioned in this publication. No reference shall be made to NOAA or this publication in any advertising or sales promotion which would indicate or imply that NOAA approves, or recommends, or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose or intent to cause directly or indirectly the advertised product to be used or purchased because of this publication.

## ACKNOWLEDGMENTS

Many individuals contributed to this project. We are especially thankful to Mr. Robert Meyer, Minerals Management Service (MMS), Anchorage, for his guidance and continued support of the arctic fisheries studies. Dr. Richard Brill, National Marine Fisheries Service (NMFS), Honolulu, provided information that guided our preparations for ultrasonic telemetry studies. Dr. Charles O'Clair, NMFS, Auke Bay, graciously loaned us telemetry instruments. Mr. Randy Bailey, U.S. Fish & Wildlife Service, Anchorage, coordinated our fieldwork with the ongoing FWS studies in the Arctic National Wildlife Range. The FWS field crew at Camden Bay provided fish for tracking, as well as timely outboard motor troubleshooting expertise. Ms. Barbara Mahoney, NMFS, Anchorage, and Ms. Jean Hanson, EPA, Anchorage, assisted in the fieldwork, as did Ms. Shelly Clouse, Lt. Matt Eagleton, Ltcd. Pat Harmon, and John Dormody of our office. Mr. David Friis reviewed the draft document. We would also like to acknowledge the assistance of the late Rae Baxter, who brought taxonomic expertise to the identification of some problematic specimens. Finally, we thank Tony and Kitty Mecklenburg of Point Stephens Press for their graphics and editorial support.

This study was funded by the Minerals Management Service, U.S. Department of the Interior, through an Interagency Agreement with the National Oceanic and Atmospheric Administration, as part of the MMS Alaska Environmental Studies Program.

## ABSTRACT

During summer 1990, NOAA conducted research to evaluate temporal and spatial patterns of fish habitat utilization in the coastal Beaufort Sea. Two distinct approaches were followed to address various issues that have been identified with respect to OCS development in the Arctic. One approach involved systematic fish sampling across geographically large areas and offshore habitats where many fish occur. The second method involved intensive study of the movement behaviors of several individuals equipped with ultrasonic tags. Large numbers of small fish were captured in the offshore sampling. Their distribution and relative abundance was then related to various oceanographic properties. Fish telemetry involved fewer and much larger anadromous fish. Individual movement responses were evaluated with respect to telemetered temperature data and other habitat attributes measured during the course of tracking.

More than 17,000 fish representing 12 families were collected in 77 townet sets. Arctic cod, capelin, and Arctic cisco were the numeric dominants of the catch. Relative abundance varied by time and location of sampling and all species appeared to be patchily distributed. Strong east winds and low river discharges resulted in the prevalence of marine conditions throughout the offshore study area. The abundance of numerous small pelagic fish in the offshore catches suggests their potential availability as an abundant food resource for anadromous species using outer portions of the coastal water band and adjacent marine environments.

Young-of-the-year (YOY) Arctic ciscoes were captured between August 2 and September 5 at stations ranging from 0.5 km to more than 12 km from the coast. The lack of environmental contrast between marine and brackish habitat conditions precluded definitive analysis regarding habitat preferences (i.e., mean CPUE brackish = 7.45 fish, mean CPUE marine = 7.23); however, the presence of juvenile Arctic ciscoes outside their suspected modal temperature range suggests the possibility of acclimation. The highest densities were reported in catches where temperatures ranged from 4 to 6°C and salinities from 27 to 29 ppt. Instantaneous increases in mean body weight per day (G) was estimated at 4.95% during early August and 2.54% during the later part of the sampling period. The occurrence of juvenile Arctic ciscoes in catches off West Dock in mid-August is indicative of the movement of small fish in coastal currents being deflected offshore by the causeway structure during east winds. Young fish were abundant in the marine conditions detected in Stump Island Lagoon at this time, suggesting that strong currents at the western end of the causeway, and not thermoregulatory behaviors, may be responsible for their presence in the lagoon. Passive and directed migration rates of 15 and 18 km/d, respectively, are presented for YOY fish.

YOY Arctic cod were abundant throughout the eastern and central portions of Camden Bay in early September. Area and volume swept methods were used to calculate population and biomass estimates for the young cod sampled on September 2 and 3. A total of 83.6 million cod (95% CI = 52.7 to 114.5 million fish) were estimated in the upper 2 m of the bay. The estimated population weight was 1.6 million kg (95% CI = 1.275 to 1.98 million kg). The wet-weight biomass of YOY cod was estimated to be 0.8-0.9 g/m<sup>2</sup>.

During the period July 30–August 11 we conducted an ultrasonic telemetry study of char and Arctic cisco in Camden Bay. Movement patterns and rates, as well as temperature occupancy and preferences, were monitored. Six large char and one cisco were tracked over a total of 120 km. Most fish sought the shoreline, but the directedness of their movements was quite variable. The chars' average gross ground speed was 55.8 cm/s (range 48.2–74.2 cm/s), or 1.04 L/s (range 0.96–1.37 L/s). Net speeds of char ranged from 9.5 to 63.2 cm/s. No significant differences in alongshore and offshore speeds were evident. The mean temperature occupied by the char was 3.6°C (range –0.5 to 6.5°C, n = 493). The frequency distribution of temperature was multimodal. The char used the entire range of observed temperatures in the study area. Temperature preference observations indicated that the fish selected the warmer halves of the available ranges in the water column. However, no consistent avoidance of cold water was evident in the horizontal dimension so the former observations may be more reflective of depth selection. The tagged Arctic cisco displayed very directed movement toward the vicinity of its capture site. Its gross ground speed was 48.9 cm/s, or 1.14 L/s, while its net ground speed was 44.1 cm/s. The mean temperature occupied by the cisco was 2.8°C (range 1.5–3.5°C, n = 51). The temperature frequency histogram was unimodal. The results are discussed in the context of similar observations on chars and other salmonids.

## TABLE OF CONTENTS

	<i>Page</i>
ACKNOWLEDGMENTS .....	iii
ABSTRACT .....	v
LIST OF FIGURES .....	xi
LIST OF TABLES .....	xv
I. BACKGROUND .....	1
Key Species .....	1
Brackish Water Habitats .....	1
Offshore Habitats .....	2
Thermal Requirements .....	3
Thermal Influences .....	4
Arctic Cisco Migration .....	6
The Issues .....	7
II. ACTIVE SAMPLING .....	9
Introduction .....	9
Objectives .....	10
Study Area .....	10
Methods .....	12
Fish Sampling .....	12
Physical Data .....	15
Fish Sample Processing .....	16
Analysis of Fisheries Data .....	16
Arctic cisco .....	17
Arctic cod .....	18
Analysis of Physical Data .....	18
Results .....	19
Distribution of Effort .....	19
Physical Oceanography .....	19
General Catch Summary .....	33
Species Characterizations .....	37
Arctic cisco .....	37
Size and growth .....	38
Time and area .....	42
Migration .....	46
Habitat .....	46

## TABLE OF CONTENTS (continued)

	<i>Page</i>
Arctic cod .....	53
Length frequencies .....	53
Habitat .....	54
Standing stock estimates .....	54
Cell block method .....	55
Zone method .....	63
Capelin .....	67
Length frequencies .....	67
Habitat .....	68
Ninespine stickleback .....	69
Length frequencies .....	69
Habitat .....	70
Kelp snailfish .....	70
Length frequencies .....	70
Habitat .....	71
Other Species .....	71
Discussion .....	71
Conclusions .....	77
<b>III. FISH TELEMETRY .....</b>	<b>79</b>
Introduction .....	79
Objectives and Approach .....	80
Study Area .....	82
Geomorphology and Geology .....	82
Meteorology .....	82
Hydrography and Oceanography .....	84
Methods and Materials .....	85
Fieldwork .....	85
Tracking apparatus .....	85
Navigation .....	86
Oceanographic Data Collection .....	87
Tagging and Tracking Techniques .....	87
Data Analysis .....	89
Telemetry Data .....	89
Oceanographic Data .....	91
Results .....	91
Fish Tracks .....	91
Movement Patterns .....	107

## TABLE OF CONTENTS (continued)

Page

Movement Rates .....	107
Temperature Occupancy .....	109
Temperature Preference .....	112
Discussion .....	113
Habitat Attributes .....	113
Methodology .....	114
Northern Dolly Varden Char .....	115
Movement Patterns .....	115
Movement Rates .....	116
Temperature Occupation .....	118
Temperature Preference .....	118
Swimming Depth .....	119
Arctic Cisco .....	120
Movement Patterns .....	120
Movement Rates .....	121
Temperature Occupation and Preference .....	121
Conclusions .....	121
Recommendations for Further Work .....	122
Instrumentation .....	122
Data .....	123
IV. REFERENCES CITED .....	125
APPENDIX A. CTD Station Data .....	135
APPENDIX B. Townet Catch Per Unit Effort .....	143
APPENDIX C. Fish Densities .....	146
APPENDIX D. Catch Summary .....	149
APPENDIX E. Townet and CTD Station Locations .....	152
APPENDIX F. Ground Speed Data .....	154



## LIST OF FIGURES

<i>Figure</i>		<i>Page</i>
1.	Study area, Beaufort Sea coast, Alaska .....	11
2.	Townet sample station locations in the vicinity of Prudhoe Bay .....	13
3.	CTD/Townet stations .....	14
4.	Distribution of townet sampling effort by date and location .....	22
5.	Distribution of fishing effort by date and location .....	22
6.	Percent fishing effort by depths sampled on the coastal transects .....	23
7.	Percent distribution of sampling effort by distance from the coast .....	23
8.	Townet station depths along coastal transects in the Beaufort Sea .....	24
9.	Location of nearshore sampling locations with distance from the coast .....	24
10.	One-meter temperatures versus one-meter salinities .....	25
11.	Time series of speed, direction, and E-W and N-S wind components in Camden Bay .....	26
12.	Time series of speed, direction, and E-W and N-S wind components in Beaufort Lagoon .....	27
13.	Vertical sections of temperature and salinity from the Endicott transect, August 8, 1990 .....	29
14.	Vertical sections of temperature and salinity from the West Dock transect, August 21-22, 1990 .....	30
15.	Vertical sections of temperature and salinity from the Arey Island transect, September 2, 1990 .....	31
16.	Vertical sections of temperature and salinity from the Collinson Point transect on September 2 and September 3, 1990 .....	32
17.	Vertical sections of temperature and salinity from the Brownlow Point transect, September 3, 1990 .....	34
18.	Total fish densities versus mean temperatures of the upper 2 meters .....	36

**LIST OF FIGURES (continued)**

<i>Figure</i>	<i>Page</i>
19. Total fish densities versus mean salinities of the upper 2 meters .....	36
20. Arctic cisco CPUE by sampling transect in 1990 .....	39
21. Arctic cisco length frequencies, all stations .....	40
22. Mean size of juvenile Arctic ciscoes for early, middle, and late periods in 1990 .....	42
23. Length frequencies of YOY Arctic ciscoes sampled at West Dock, August 21-22, 1990 .....	43
24. Ninety-five percent confidence intervals for Arctic cisco CPUE by catch district .....	44
25. Mean daily rates and mean daily directions of a passive migration .....	47
26. Mean daily rates and mean daily directions of a directed migration .....	48
27. YOY Arctic cisco catch versus mean temperatures of the upper 2 meters .....	49
28. YOY Arctic cisco catch versus mean salinities of the upper 2 meters .....	49
29. Cluster plot of YOY Arctic cisco CPUE with mean temperature and salinity conditions of the upper 2 meters .....	50
30. Bivariate normal surface plot of temperature (upper 2 m) and Arctic cisco density .....	54
31. Arctic cod length frequencies, all stations .....	55
32. Arctic cod catch versus mean temperatures of the upper 2 meters .....	56
33. Arctic cod catch versus mean salinities of the upper 2 meters .....	56
34. Length frequency distribution of Arctic cod in Camden Bay, September 2-3, 1990 .....	57
35. Cell blocks in Camden Bay .....	58
36. Zones in Camden Bay .....	64
37. Capelin length frequencies, all stations .....	68

## LIST OF FIGURES (continued)

<i>Figure</i>		<i>Page</i>
38.	Ninespine stickleback length frequencies, all stations .....	69
39.	Snailfish length frequencies, all stations .....	70
40.	Study area for acoustic telemetry, Camden Bay, Beaufort Sea, Alaska, 1990 ..	83
41.	Track of Fish 1 .....	93
42.	Temperature vs. elapsed time, Fish 1 .....	94
43.	Track of Fish 2 .....	95
44.	Temperature vs. elapsed time, Fish 2 .....	96
45.	Track of Fish 3 .....	97
46.	Temperature vs. elapsed time, Fish 3 .....	98
47.	Track of Fish 4 .....	99
48.	Temperature vs. elapsed time, Fish 4 .....	100
49.	Track of Fish 5 .....	101
50.	Temperature vs. elapsed time, Fish 5 .....	102
51.	Track of Fish 6 .....	103
52.	Temperature vs. elapsed time, Fish 6 .....	104
53.	Track of Fish 7 .....	105
54.	Temperature vs. elapsed time, Fish 7 .....	106
55.	Directedness indices and compass bearings of fish .....	107
56.	Standardized ground speed vs. elapsed time .....	109
57.	Raw ground speed vs. temperature, Fish 2 .....	110
58.	Standardized ground speed vs. temperature .....	110
59.	Temperature occupancy, all char .....	111

**LIST OF FIGURES (continued)**

<i>Figure</i>		<i>Page</i>
60.	Temperature occupancy, Arctic cisco .....	111
61.	Temperature selection observations .....	112
62.	Track of Fish 2 relative to horizontal temperature field as indicated by temperatures transmitted from the fish .....	119

## LIST OF TABLES

<i>Table</i>	<i>Page</i>
1. Scientific and common names of fish collected by townetting in 1990, coastal Beaufort Sea, Alaska .....	35
2. Species composition and relative abundance of the 1990 townet catch .....	35
3. Arctic cisco catch summary by transect or other coastal area sampled by townet in 1990 .....	38
4. Length measurements of juvenile Arctic ciscoes collected by townet in 1990 ..	41
5. Mean CPUE and density of YOY Arctic cisco during four sampling periods in August and September 1990 .....	45
6. Summary of cluster analysis including stations, sampling dates, and mean indices of relative abundance and environmental conditions for each cluster ..	51
7. Grid cell information for YOY Arctic cod in Camden Bay study area, September 2-3, 1990 .....	60
8. Estimated population of YOY Arctic cod in Camden Bay during September 2-3, 1990, using the cell block method .....	61
9. Variance estimates of $\overline{CPUE}_{ik}$ in Simpson Cove, Camden Bay, Arey Island, and Brownlow Point subareas .....	62
10. Zone and station information used to compute population parameters for YOY Arctic cod in Camden Bay, September 2-3, 1990 .....	65
11. CPUE and population weight estimates for YOY Arctic cod in Camden Bay, September 2-3, 1990 .....	66
12. Summary of results from ultrasonic tracking of northern Dolly Varden char and Arctic cisco in Beaufort Sea coastal waters, 1990 .....	92
13. Ground speed statistics of northern Dolly Varden char and Arctic cisco based on hourly averaged data .....	108
14. Gross ground speeds of salmonids .....	116

## PART I. BACKGROUND

### Key Species

The exploration and development of Arctic oil reserves has prompted numerous fishery investigations in the Alaskan sector of the Beaufort Sea. The study of anadromous<sup>1</sup> fish populations has received special attention because of their importance in rural economies and coastal ecosystems, and the possibility that industrial developments associated with petroleum activities could affect these populations. Additionally, the management of several species is of international concern. In these instances, the anadromous life cycle includes long-distance migrations, extended rearing at sites far removed from natal origins, and overwintering in transboundary rivers.

Arctic cisco (*Coregonus autumnalis*), Arctic char (*Salvelinus alpinus*), least cisco (*C. sardinella*), and broad whitefish (*C. nasus*) have been the focus of most coastal fish studies in the Alaskan Beaufort Sea. These anadromous fish tend to be the largest and most conspicuous members of the nearshore fish community in the arctic. Actually, several of the less-studied marine species are numerically much more abundant. Anadromous fish stocks are characterized by low recruitment rates and other density-dependent mechanisms that allow their populations to survive arctic conditions (Craig 1989a). Availability and quality of overwintering habitat are viewed by many as the greatest determinant of population size.

### Brackish Water Habitats

A common feature of arctic anadromy is the annual migration of fish from freshwater overwintering areas to summer feeding habitats. Summer habitats are located within the narrow band of relatively warm, low salinity water that develops during summer along the Beaufort Sea coast. The brackish water band is an ephemeral and dynamic habitat that, typically, is continuous early in the summer and discontinuous later. Discontinuity coincides with declines in runoff and the mixing of coastal and oceanic water masses. The width of the brackish water band differs with location and has been estimated to extend from 1 to 10 km from the coast (Craig 1989a). Wind stress, horizontal pressure gradients, and tides are the driving forces of the coastal circulation (Colonell and Niedoroda 1990a,b). Such geophysical forcing occurs across a wide range of temporal and spatial scales and its expression may be periodic (tides) or less regular (upwelling). For instance, the hydrographic properties at any given time in summer at Prudhoe Bay result not only from prevailing meteorological and river discharge conditions, but also from conditions several hours to several weeks prior (Hale et al. 1989). A similar generalization is pertinent to any location along the coast.

Maughan (1990) likened the brackish water band to a "constantly changing mosaic of salinity and temperature patches driven by wind and water inflow and bathymetric conditions." Superimposed on the physical environment he envisioned heterogeneous distributions of predators and their prey. This variability in hydrographic regimes is evident at mesoscales (10's - 100 km) from satellite observations of the nearshore environment during

---

<sup>1</sup>Anadromous is synonymous with amphidromous in this paper (see McDowall 1987).

the summer. A qualitative analysis of thermal images taken during 13 days in 1988 revealed a general trend for warmest waters to be located nearest the coast and coldest waters to be found nearest the ice pack (Thorsteinson et al. 1990). The heterogeneity of the coastal band was confirmed by the "patchwork" distribution of temperature regimes noted in the thermal images.

## **Offshore Habitats**

We have argued previously that the existing arctic fish database is largely one-dimensional in space, reflecting the inshore concentration of existing research efforts (Thorsteinson et al. 1990). This inshore emphasis has led to generalizations about use of coastal waters by fish in poorly sampled areas. For instance, Colonell and Gallaway (1990) state that research has demonstrated the anadromous species are generally restricted in distribution, from the coastline out to a depth of 2 m, where cold (0°C), marine bottom water occasionally upwells near or to the surface. Such a restricted range is probably more true for some species than others. In fact, considering the annual variability in ocean conditions, it may reflect depth-imposed gear limitations as much as availability of suitable fish habitat. As a consequence, the onshore-offshore features of movements and migrations, habitat use, and general ecology of several species are unknown.

Few studies have addressed fish use of habitats in outer portions of the coastal brackish band or in the marine habitats beyond. Craig (1984) concluded that the wide variations in temperature and salinity distributions across the coastal habitats defied their easy description by any physical attribute such as depth or distance from shore. Instead, he reasoned that the categorization of habitats by water mass characteristics was probably the most meaningful approach within the context of existing information.

In 1988, NOAA conducted offshore purse seining to augment the meager offshore data and to provide an "offshore dimension" to an ongoing Arctic char stock identification project (Everett and Wilmot 1990). Fish sampling was conducted in three hydrographically defined habitat types (Thorsteinson et al. 1990). It was hypothesized that Arctic char occupy the entirety of the coastal brackish water habitat, but do not habitually venture into the colder, more saline waters. An inferential analysis was performed to describe frequency of occurrence patterns indicative of habitat associations in coastal, transitional, or marine habitats. Anadromous species were associated with the coastal habitat while marine species were distributed throughout the habitats sampled. The structural occurrence of coastal waters overlaying other water masses was thought to provide the physical pathway enabling offshore dispersals by anadromous fish. It was suggested from qualitative observations of food habits that the offshore movements were a distributional response to density of prey.

Arctic char and Arctic cisco were the only anadromous species captured in the offshore seining. The greatest number of Arctic char was captured approximately 5 km off Bullen Point in eastern Stefansson Sound. These char were generally small (FL 240 mm) and may have been moving with Sagavanirktok River water that had been transported to the east under west wind conditions. Gallaway (1990) indicates that there is considerable evidence that small fish, at least, travel in the direction of drift and that fish displaying such movement behavior have a bioenergetic advantage. Dempson and Kristofferson (1987)

characterized the ocean movements of Arctic char as a feeding migration influenced by the spatial and temporal distribution of a major prey, the capelin (*Mallotus villosus*).

Rose and Leggett (1989) hypothesized that the geographic distribution of Atlantic cod (*Gadus morhua*) in the northeastern Gulf of St. Lawrence is determined by geophysically forced oceanographic conditions. Within these limits, cod distribution is further delimited by interacting physical (e.g., water temperature) and biotic (prey) conditions. They argued that: (1) cod distribution is largely controlled by water temperature; (2) most species frequent a much narrower "modal" temperature range; and (3) highly mobile migratory species reduce their exposures to environmental extremes by moving in synchrony with preferred conditions. Correlation analysis (univariate methods) between cod catch and sea temperature, and cod catch and prey (capelin) density variables did not adequately explain the observed patterns of cod distribution or catch. Regression analysis, on the other hand, indicated that under mesoscale conditions of high prey density and favorable temperatures cod distributions closely matched those of capelin throughout their distributional range (-0.5 to 8.5°C). When prey density was low, the distribution of cod was unaffected; they remained within their modal range (0-5°C). In the latter instance, the infrequent occurrence of cod at temperatures outside their modal range was thought to reflect a distributional response to prey density. In conclusion the authors note that the interaction of sea temperature and prey density to regulate cod distribution may apply to all cod populations.

### Thermal Requirements

In the study above, "modal" and "preferred" temperature ranges were distinguished by reference to observations of free-ranging and experimentally held fish, respectively. Houghton et al. (1990) conducted a habitat use analysis based on a 4-year arctic fish database containing some 180,000 fyke-net catch records from Prudhoe Bay. This analysis provides preliminary information about the temperature-dependent range limits of the four most common species of anadromous fish in the Beaufort Sea. The analysis was limited to early (mid-June) and midsummer (mid-August) periods. As presented below, modal range reflects temperatures associated with peak utilization as indicated by daily CPUE.

<i>Species</i>	<i>Geographic Range (°C)</i>	<i>Modal Range (°C)</i>
Arctic cisco	0->12	6-12
Arctic char	0-12	4-10
Least cisco	0-12	8->12
Broad whitefish	2->12	8->12

The thermal ranges described above undoubtedly underestimate the true distributional limits of the fish during the open-water season. Craig (1984) reported additional information on the temperature ranges of the anadromous fish from a synthesis of many research projects; however, his ranges are in good agreement with those presented above. The Houghton et al. (1990) analysis was truncated in mid-August to reflect coastal usage patterns during summer periods of greatest fish dispersal. The result is that at least 1 month of open-water season is excluded from the analysis. Water temperatures below 0°C are not uncommon in the coastal zone during summer months and are especially evident just prior



to freeze-up. After August 15, the nearshore tends to experience steadily decreasing temperatures and increasing salinities. Anadromous fish are able to withstand such conditions for at least brief periods (Colonell and Gallaway 1990). As an example, Johnson (1980) reports Arctic char in Creswell Bay, Canada, withstanding subzero temperatures ( $-1.4$  to  $1.44^{\circ}\text{C}$ ) at depths to 15 m.

Selected oceanographic data from Camden Bay in "warm" and "cold" years illustrate the annual variability in temperature conditions of anadromous fish habitats. Hale (1991) indicates that water temperatures during the 1989 season were above average. A maximum temperature of  $11.4^{\circ}\text{C}$  was reported on August 8 at a Fish and Wildlife Service sampling station (CB06) located 2.6 km (9-m water depth) off Carter Creek in Camden Bay. Closer inshore at station CB02 (located approximately 0.7 km off Carter Creek, 3-m water depth), a maximum temperature of  $14.2^{\circ}\text{C}$  was observed. At shallow-water stations located in Simpson Cove and in the lagoons east of Barter Island, water temperatures in excess of  $14^{\circ}\text{C}$  were observed. Similar temperatures from these stations on August 8, 1988, were  $2.2^{\circ}\text{C}$  at CB06 and  $2.2^{\circ}\text{C}$  at CB02 (Hale 1990).

Temperature preference data are lacking for most of the arctic anadromous fishes. Fechhelm and Gallaway (1984) reported temperature preferences of small Arctic cisco (83–136 mm) acclimated to several temperature and salinity regimes representative of the brackish water zone. Their experiments showed a preference by Arctic cisco for temperatures generally greater than  $10^{\circ}\text{C}$  over the range of salinities studied (5–30 ppt). In fact, the test fish selected the warmest available temperatures ( $14$ – $15^{\circ}\text{C}$ ). Such conditions approach the upper limits of the thermal spectrum that might be encountered in the coastal Beaufort Sea (Fechhelm et al. 1982). These authors noted that the apparent preference for warmer water explains the Arctic cisco's utilization of a relatively narrow coastal corridor (20–50 m wide) during summer. The physiological benefits conferred by such habitat use, especially on growth and migration, have been discussed elsewhere (Fechhelm et al. 1982; Craig 1984, 1989a; Gallaway 1990; Thorsteinson et al. 1990; English, in press).

Similar experimental results for other anadromous species are not available. The large body of field data that has been obtained from the Prudhoe Bay area over the past 15 years indicates varying tolerances of the major anadromous species (cf., Ross 1988). Besides Arctic cisco, only Arctic char are expected to disperse farther than 1 km offshore (Craig 1984; Fruge et al. 1989). Circumstantial evidence suggests Arctic char may prefer slightly cooler temperature conditions than the Arctic cisco. Berg and Berg (1989) studied the at-sea growth of anadromous Arctic char in northern Norway. Optimal growth in char was observed during June when temperatures ranged from  $5$  to  $10^{\circ}\text{C}$ . The authors suggested that ocean temperatures in excess of  $14^{\circ}\text{C}$  limit the southern distribution of the species. Elsewhere, in arctic lakes, optimum growth in Arctic char has been observed at temperatures between  $12$  and  $16^{\circ}\text{C}$  (Johnson 1980).

## **Environmental Influences**

Water temperature may be the greatest physical determinant of fish distribution (e.g. Rose and Leggett 1989). In the Arctic, the brackish water nearest the coast is typically characterized as hydrographically most favorable for anadromous fish (e.g., Craig 1984).

Houghton et al. (1990) and others (see Fechhelm et al., in press) report a general trend for greater use of warm (4–12°C), low- to moderate-salinity (0–20 ppt) water. Other interacting factors such as currents, food habits, prey availability, and size and condition of the fish, influence dispersal patterns (cf. Dempson and Kristofferson 1987). Colonell and Gallaway (1990) subjectively list (in order of importance) the most important ingredients of anadromous fish habitat quality: (1) abundance of prey, (2) temperature, (3) salinity, and (4) “predictability” or stability of the other three factors. Temperatures at or below 5°C seem to represent a lower limit for habitat use by arctic anadromous fish (i.e., reduced growth rates and return migrations begin below this temperature) (Colonell and Gallaway 1990).

Temperature–salinity challenge experiments conducted on small Arctic ciscoes support the generalized trend of reliance on brackish water habitats by anadromous fish (Fechhelm et al., in press). The size range of Arctic cisco in these bioassays was 120 to 170 mm (FL). Young-of-the-year (YOY) fish were not studied. Prior to experimental testing the ciscoes were held at ambient conditions approximating those of their capture location for a minimum of 3 d. The experimental protocol required acclimation rates of 2°C and 5 ppt changes/d to one of three temperature (12, 7, and 3°C) and salinity (0, 15, and 32 ppt) regimes. The ciscoes were held at acclimation conditions for at least 14 d before starting 96-h bioassays involving instantaneous exposures to various temperature and salinity combinations.

Acclimation appeared to enhance the tolerances of experimental fish to elevated salinity (Fechhelm et al., in press). Osmoregulatory failure and lethal effects were, however, observed at salinities of 25 ppt or greater. The observed deleterious effects of prolonged exposures to cold water on osmoregulation in Arctic cisco may be similar for other anadromous species. This result would indicate avoidance, or very limited use, of marine areas in the wild. Other experimental evidence supports this conclusion. The few bioenergetic studies from which data are available indicate minor effects on growth rate at salinities of 0–15 ppt and significant reductions in growth and increased mortality at higher salinities (e.g., English in press).

Colonell and Gallaway (1990) suggest that salinity has little effect on the distributional response of anadromous fish until some threshold level is reached. Salinity tolerance evidently varies by species and with size of fish. Among the anadromous species, small fish tend to be less tolerant than larger fish. Conversely, immature Atlantic cod are able to endure temperatures well below those tolerated by older fish (Rose and Leggett 1989). Many factors such as size, age, and sexual condition of the fish influence its tolerance and ability to acclimate to changing salinity. These relationships remain undescribed for most arctic fish.

In situ growth experiments conducted in 1988 (English, in press) reinforce the concept of anadromous fish preference for brackish water habitats. Arctic cisco and broad whitefish were held for 6 wk in mesh enclosures at Endicott and Niakuk study areas. Water depths at these sites averaged about 1.5 m. Mean weekly temperature and salinity conditions were similar at each site, ranging from about 7°C and 11 ppt (early season) to 2°C and 25 ppt (late season). Test fish ranged in length from 120 to 180 mm (FL) and in weight from 15 to 45 g. Differences in growth and survival were noted between enclosure sites and species. Highest growth rates were observed during late July to mid-August. In 1988 these corresponded to

periods of highest ambient temperatures (about 7°C) and low to intermediate salinities (>15 ppt). Broad whitefish held at the Niakuk site suffered higher mortalities and lower growth rates than fish rearing near Endicott. While temperature and salinity conditions were partially responsible, the Niakuk site was more exposed, resulting in increased mortalities to fish during inclement weather. Broad whitefish fared poorly at each location, suggesting poor habitat conditions for this species at these sites in 1988. The growth rate observed in Arctic cisco was 6 times higher than that of broad whitefish. This was thought to reflect the greater flexibility in habitat use by the Arctic cisco.

### **Arctic Cisco Migration**

The lack of discovery of spawning or spawned-out Arctic cisco in the Colville River and delta led Gallaway et al. (1983) to hypothesize that Arctic cisco in the Alaskan Beaufort Sea are of Mackenzie River origin. Prevailing east-to-west longshore currents provide the dispersal mechanism for age 0 fish from the Mackenzie estuary to reach Alaskan overwintering sites. Gallaway et al. (1983) proposed that the Arctic cisco migration involves a passive transport of fish in wind-driven currents. A passive migration is supported in part by comparisons of summer wind conditions with (1) commercial catch statistics from the Colville Delta (Fechhelm and Fissel 1988); and (2) relative abundance estimates of YOY fish in Prudhoe Bay (Fechhelm and Griffiths 1990). Average northeasterly wind conditions (5 m/s), and current speeds approximating 3 to 4 percent of the winds (15 cm/s), imply a 35 d (13 km/d) migration period for YOY fish travelling between Mackenzie and Colville river habitats (Gallaway et al 1983). The average rate of westward movement by Arctic cisco of 13.6 km/d observed by Moulton (1989) is consistent with this estimate.

Neill et al. (1983) developed a biased random walk model to evaluate the movement patterns of Arctic cisco around the West Dock causeway. Under various scenarios, high changes in modeled catch rates could not be explained by swimming rate alone. It was suggested that either the fish were being entrained in strong currents in the causeway area or that catchability rates (and not fish density) were being affected (delayed or obstructed) by hydrographic changes. The "pulsing" observed in the model's results was attributed to distributional responses by the fish to shifting winds and currents and altered nearshore conditions.

One constraint of the model was that it was unable to address adequately the width-of-path effect of the Arctic cisco migration (Neill et al. 1983). No information (physical or biological) was available regarding this aspect of migration and it was therefore assumed that the fish either (1) do not venture farther offshore than 0.5 km north or northwest of West Dock (which extended 3.9 km from the coast at this time); or (2) venture offshore but only into waters possessing conditions equivalent to those found closer inshore. The first attempt to obtain information on the offshore distribution and abundance of age 0 Arctic cisco in Prudhoe Bay and to study their migration in the causeway areas was undertaken by Houghton and Whitmus (1988). Unfortunately, their sampling objectives were confounded because few recruits moved into the study area in 1988 (Schimdt et al., in press). Variable and westerly wind conditions in 1988 may have been responsible agents for the lack of YOY Arctic cisco.

Only three Arctic cisco were captured in NOAA's seining in 1988: one off Bullen Pt. (FL = 375 mm); one in Camden Bay (84 mm); and one near Thetis Island (146 mm). The juvenile fish sampled in Camden Bay on August 21, 1988 (station C04081, 5.5-m depth), was captured approximately 0.75 km off the coast. This catch provides preliminary evidence that a wider migration corridor is being utilized than has been suggested elsewhere.

More recently, modeling (random walk with advection) of the movement of young Arctic cisco in Prudhoe Bay was performed by Bryan (1990) as part of the Endicott Development Fish Monitoring Program. The study focused on the migration of the fish past the Endicott Causeway from August 20 to September 1. Initial fish density distributions were selected on the basis of actual fyke-net data and the temperature preference data of Fechhelm and Gallaway (1984). Oceanographic data inputs (current vectors, etc.) from other sources were indicated by the author but were not clearly specified. Fine-scale (1 km) movement patterns were apparently biased such that fish were generally moving toward more optimal water quality conditions. The algorithms used were successful predictors of the relative magnitudes and general trends in observed catches. A major conclusion was that currents, and not thermoregulatory behavior, had the greatest effect on YOY Arctic cisco distribution. The effects of temperature and salinity different from those modeled require further investigation.

The above is not meant to downplay the role of currents in anadromous fish migrations and movements. Previously, we argued that the development of realistic predictive models would require a more detailed history of fish movements along known environmental gradients than is presently available (Thorsteinson et al. 1990). Catch data from passive gears with long set periods are not amenable to correlation with physical phenomena of brief temporal and spatial scales, such as the passage of fronts, eddies, and wind-driven events, which are prevalent in North Slope coastal waters. Such short-lived phenomena are important because they influence the local distributions, movements, and other daily activities of fish and their prey. Colonell and Gallaway (1990) argue that because fish distributions are not static, sampling with active gear will not necessarily provide data that would define preference. It will, however, with adequate sampling intensity delimit the "modal" range described above and provide more precise data on fish abundance.

## **The Issues**

Solid-fill gravel causeways have proven to be a cost-effective technology for onshore transport of (1) seawater via pipelines for waterflooding, and (2) produced crude oil to the Trans-Alaska Pipeline (Standard Alaska Production Company 1989). However, environmental issues have been identified concerning causeway construction and subsequent effects on coastal oceanography. For instance, Hale et al. (1989) describe the major effects of West Dock on local circulation. They include offshore deflection of alongshore flow (including river plumes) and the blockage of alongshore movement and mixing of water mass properties. As a result, other marine processes are influenced as well, including coastal hydrographic conditions, upwelling, and vertical mixing. These authors conclude that under either easterly or westerly winds, West Dock can deflect alongshore surface currents and plumes northward around the tip at least 4 km offshore.

Other authors indicate that the marine environment has been little affected by the construction of causeways. Colonell and Gallaway (1990) also provide an assessment of marine environmental impacts of West Dock causeway. Colonell and Niedoroda (1990a) maintain (from a physical oceanographic perspective) that Stump Island Lagoon is a minor appendage of Simpson Lagoon that "never has played a major role in establishing the hydrographic conditions there, even in the eastern part of Gwydyr Bay." This conclusion is based on the cross-sectional area of the barrier island channels in eastern Simpson Lagoon.

The relative time and spatial dimensions of the marine areas affected by causeway obstruction are dependent on the prevailing oceanographic and meteorological conditions at any given time. The extents of coastal areas affected by changes in water quality are therefore difficult to determine. Colonell and Gallaway (1990) assert that the geographic extent of salinity increase attributed to the Waterflood extension of West Dock is of the order of 800 ha but that this area is not easily determined. Ross (1988) estimated that water quality and circulation patterns along as much as 60 km of coastline have been altered by the West Dock and Endicott causeways.

Prediction of biological effects on anadromous fish populations of the North Slope associated with oil and gas development in or near Prudhoe Bay has proven to be an arduous and controversial process. It has been hypothesized that in the marine environment causeways and waterflood projects may interfere with fish migrations, habitat quality, and overall fish use of coastal waters. Since estimates of population size are unreliable for most species, parameters such as growth and condition have been evaluated for several years in the Prudhoe Bay area. The sensitivity of the condition analysis is questionable as it tends only to include fish in good condition. With increasing industrialization of the North Slope, the resolution of these and other fishery issues (e.g., oil spills, availability of fresh water, and gravel excavation) will be required on regional as well as local scales.

Information is lacking concerning the environmental tolerances and responses of most anadromous fish species to hydrographic conditions they may confront in the coastal Beaufort Sea. This, in part, has limited our ability to accurately assess the probable effects of certain development activities. The coastal brackish band serves as an avenue for the alongshore movement of anadromous fish between feeding, spawning, and wintering locations. Because anadromous fish must acquire so much of their annual energy supply during the brief open-water season (Craig 1989a), the availability and quality of feeding habitats are of primary concern. Access to these habitats is critical during summer months. Blockage, or delays, of fish migration may result from physical barriers imposed by the causeways. The potential effects of causeways on YOY Arctic cisco migration have been of concern (MBC 1990). There are two issues: (1) blockage or delay of migration in the causeway areas, and (2) the transport and fate of young fish entrained in coastal currents that have been deflected offshore.

## **PART II. ACTIVE SAMPLING**

### **INTRODUCTION**

The goal of NOAA's Arctic Fish Habitat Use Investigations is to ensure the protection of fishery populations and their habitats from potential adverse effects of OCS development in the Alaskan Beaufort Sea. The research has focused on the identification and characterization of important marine habitats and temporal-spatial features of their use by arctic fishes. Anadromous species are of special management concern. Their populations tend to be small in size, possessing low recruitment rates and other traits common to K-strategists (Craig 1989a). Such populations (i.e., characterized by great longevity coupled with low adult mortality and recruitment) differ from truly stable populations where recruitment and mortality rates are nearly equal. Several species are valued subsistence resources and others are targeted in small commercial fisheries located in the lower Colville River. In addition, the management of several species is of international concern because of the transboundary nature of migration routes and overwintering areas.

In an ecological sense, the aim of our research is to gain an understanding of the dispersal mechanisms and processes responsible for observed patterns of fish distribution and abundance across the Beaufort Sea shelf. This has required oceanographic sampling in habitats occupying outer portions of the brackish water band and beyond. This is significant because it provides an offshore component to the existing arctic fish database. Fishery information is essentially lacking from marine environments of the Beaufort Sea (Craig 1984). The offshore sampling provides an opportunity to incorporate different sampling methods into the research design than have been practical closer inshore. We chose an active sampling gear approach because resultant indices of abundance provide quantitative estimates of standing stock densities that can be directly related to ambient environmental conditions (e.g., temperature, salinity, winds and currents, and food availability).

Exploratory oil drilling in Camden Bay, to the northeast of Barter Island, and northwest of Harrison Bay, are indicators of impending industrial expansion in the Beaufort Sea. With few exceptions, fishery resource information from these areas is sparse or nonexistent. The possible opening of the Arctic National Wildlife Refuge (ANWR) to oil and gas exploration has prompted coastal fishery surveys in the eastern Beaufort Sea (e.g., Fruge et al. 1989). Unlike the open lagoon systems characteristic of Stefansson Sound, exposed nearshore waters and limited exchange lagoons predominate in the east. Such geomorphological differences are important because the structural gradients defined by freshwater inflows, wind-mixing, upwelling, and other physical processes ultimately define the useable boundaries of available habitat for marine and anadromous fish. They may also establish important mechanisms which may enhance or impede migrations, dispersals, or other uses of the coastal zone by fish.

Four species of anadromous fish have been of primary concern in fishery investigations in the Alaskan Beaufort Sea. These are the Arctic cisco, Arctic char, least cisco, and broad whitefish. The research has consisted of various assessments of the potential impacts to fish populations associated with oil and gas development in or near Prudhoe Bay. In particular,

the effects of solid-fill causeways on coastal oceanography and anadromous fish habitats have been a special environmental concern. It has been hypothesized that causeways and water-flood projects may interfere with fish migrations, habitat quality, and overall fish use of coastal waters. The range of possible impacts varies by species and appears to be largely a function of their tolerance to coastal temperature and salinity conditions. With increasing industrialization of the North Slope the resolution of these, and other issues, will be required on regional as well as local scales.

## OBJECTIVES

A primary focus of the 1990 sampling was the coastal migration of juvenile Arctic cisco (*Coregonus autumnalis*) in Alaskan waters. Gallaway et al. (1983) had proposed a wind-aided, passive transport of Mackenzie River cisco to the Colville River delta. Evidence is mounting to support the "fish from Canada" hypothesis (Fechhelm and Fissel 1988; Fechhelm and Griffiths 1990) and it has been suggested that the migration is primarily a shoreline-directed nearshore phenomenon. There is, however, little reliable offshore catch data available to support the latter contention and it is not yet possible to delineate the migratory corridor itself.

Although our sampling targeted on juvenile Arctic ciscoes, various other species were collected in the offshore fishing. A secondary objective was, therefore, to describe the general features of habitat use by these fish and, if possible, to evaluate growth and condition parameters for the dominant species involved.

The specific objectives of the 1990 fieldwork were (1) to determine the spatial-temporal distributions, relative abundances, and habitat associations of dominant fish species in the nearshore Beaufort Sea; and (2) to delineate the migratory corridor of juvenile Arctic cisco at several locations in the Alaskan Beaufort Sea.

## STUDY AREA

The study area included the coastal waters lying between Prudhoe Bay and Barter Island (Fig. 1).

Most operational activities were based at Camden Bay. Camden Bay is a relatively deep embayment in the western portion of the ANWR. It is located between Barter Island, in the east, and the Canning River delta, to the west. The natural setting of Camden Bay (see Fruge et al. 1989) provides easy access to coastal and marine fish habitats, while at the same time providing close proximity to safe anchorages at Simpson Cove and Barter Island.

Camden Bay is typical of many of the exposed nearshore habitats characteristic of the eastern Beaufort Sea. With respect to oil and gas development, some exploratory drilling in the western bay has already occurred. Further development can be expected. If oil leasing occurs in the ANWR, Camden Bay is a likely site for port expansion and other shore-based support. With respect to our study objectives, the deep nature of the bay, particularly on the




Figure 1.—Study area, Beaufort Sea coast, Alaska.



eastern shore, allows towed gears to be deployed quite close inshore. This is an important operational feature of the migration and dispersal components of the research.

The research vessel was regularly provisioned and serviced at West Dock in Prudhoe Bay. Since these activities required regular travel between Camden and Prudhoe bays, several additional sampling stations were established in the causeway areas (Fig. 2). These stations coincided with the sampling locations of Houghton and Whitmus (1988). Mid- to late August sampling periods were given highest priority as they would encompass periods of peak YOY Arctic cisco migration through Prudhoe Bay.

## METHODS

### Fish Sampling

All oceanographic research was conducted from NOAA vessel *1273*, an 11-m aluminum craft with a draft of 1.3 m. This boat is currently configured as a drum seiner. It has a maximum speed of 8 knots, a range of 730 km, and an endurance of 7–10 d, and can accommodate four persons. Other attributes of *1273* for working in the shallow Beaufort Sea have been described by Thorsteinson et al. (1990).

Five transects were selected for fish sampling in 1990 (Fig. 3). From west to east, they were located (1) west of West Dock, (2) on the east side of the Endicott Causeway, (3) off Brownlow Point, (4) off Collinson Point, and (5) off Arey Lagoon to the west of Barter Island. The transect placements reflect entry and exit locations of migratory fish passing through Camden and Prudhoe bays. The Collinson Point transect increased the spatial resolution of sampling in Camden Bay (mid-point transect) and added a necessary onshore–offshore dimension to our monitoring of the Arctic cisco migration through the center of the bay.

Our study plan also included fish sampling near Carter Creek in Camden Bay. It was anticipated that this sampling would coincide with FWS coastal sampling in the area. Our expectation was that synoptic fishing efforts would facilitate the description of a realistic relationship between catches associated with active and passive gear types.

A single station (MS01) was positioned near Flaxman Island and the Mary Sachs Entrance to eastern Stefansson Sound. This station was strategically located to examine its role as a potential “choke” point for migrant cisco en route to Prudhoe Bay.

In most instances the locations of stations on each transect were stratified by their relative distance from the mainland coast. Inner transect stations, or “base stations,” were generally defined by the shallowest working limitations of the primary sampling gear. Two exceptions included the location of base stations on Arey Island and Brownlow Point transects. Along these transects, the bases were located 3.5 km from the coast. In Camden Bay, once the base station had been defined, other transect stations were located approximately 1.5, 4.5, 7.0, and 9.0 km from the base (and 11.0 km on the Collinson Pt. transect).




Figure 2.—Townnet sample station locations in the vicinity of Prudhoe Bay.




Figure 3.—CTD/Townet stations.

A description of the Prudhoe Bay transects and stations has been provided by Houghton and Whitmus (1988) and is briefly described here. The Endicott transect stations were located at 0, 1, 3, 5, 7, 10, 15, 20, 25, and 30 km from a base lying 1.3 km from the shoreline in Foggy Island Bay. At West Dock the base station was 0.6 km from the east end of Stump Island and the causeway. This translates to a distance of approximately 3 km from the coast. Stations were sampled at about 0, 2, 4, and 6 km from this base. An additional station (WD25), located about 11 km offshore, was added to the transect.

The stations in Simpson Cove (CP10) and Mary Sachs Entrance (MS01) were located almost 2 km from the mainland coast. In each instance, this position roughly demarked midway points in the respective lagoons. The Carter Creek stations were originally selected as a function of depth (i.e., 5 and 20 m, respectively).

Fish sampling was accomplished with an 11.0-m-long townet. The mouth of the net was  $6.8 \times 1.8$  m tapering to a  $0.3 \times 0.3$  m cod end bag. Net panels were of variable mesh size as follows: front panel, 50.8 mm; 2nd panel, 38.1 mm; 3rd panel, 19.1 mm; and bag, 7.9 mm. At each station replicate sampling consisted of two 10-minute hauls to the east. The net was towed at an approximate speed of 0.5 m/s.

A Magnavox MX 5400 GPS Satellite Navigation System was utilized in the field to locate the townet sampling stations. The accuracy of the GPS is better than  $\pm 100$  m. However, because the GPS system was not fully operational in 1990, satellite coverage was intermittent during the field season. During periods when fewer than 3 satellites were available for GPS fixes, a Furuno Model FR805D 48 nm radar was used to determine position information. The accuracy of radar fixes is assumed to be 0.5 km.

### Physical Data

Thermohaline structure was recorded at each transect station prior to the conduct of replicate townet sampling with a portable conductivity-temperature-depth instrument (Applied Microsystems STD-12) having an internal recording capability. The instrument's depth, temperature, and salinity sensors are accurate within 0.10 m, 0.03°C, and 0.2 ppt, respectively. The CTD stores data in an internal memory and records eight samples of temperature, conductivity, and pressure per second. During oceanographic casts, the CTD's sensors were held just below the sea surface and allowed to equilibrate, then the instrument was slowly lowered ( $\bar{x} = 6.89$  cm/s,  $SD = 3.95$ ,  $n = 39$ ) to the sea floor. Upon completion of the cast, CTD data were downloaded to a Zenith laptop portable computer, reviewed for accuracy, and stored on minidiskettes for processing.

Summer 1990 wind data from the eastern Beaufort Sea were obtained from the Fish and Wildlife Service. These data were acquired from weather stations maintained at Simpson Cove and Beaufort Lagoon. Wind speed, direction, temperature, and barometric pressure were recorded every 0.5 h. The Simpson Cove record was truncated due to sensor malfunctioning and data were available only for the period July 20-31, 1990. The Beaufort Lagoon record was longer, extending from July 21 to September 10, 1990.

Other physical measurements were taken in concert with oceanographic casts. Water depths were determined with a Ray Jefferson Color Telescan 2000 sounder. Wind speed (from the vessel's anemometer) and direction (estimated) were recorded during townet operations.

### Fish Sample Processing

Townet catches were sorted to species, counted, and individually measured. All catches were enumerated in the field. A volumetric approach (known quantities/unit volume) was utilized to estimate total numbers in several instances of very large catches. Fork length (nearest mm) and wet weight measurements (0.1 g) were taken immediately following fish capture. An OHAUS (2610 g) Dial-O-Gram was used to measure fish weights as sea conditions allowed. All station, measurement, and environmental data were recorded on field forms for later data entry.

Some fish were kept for additional processing and taxonomic verification. Specimens were either (1) frozen on dry ice in seawater, or (2) preserved in a solution of 10% formalin in seawater, with preservation technique dependent on the work to be conducted ashore. A total of 290 frozen samples (121 Arctic cisco; 69 capelin; and 100 Arctic cod, *Boreogadus saida*) are currently being maintained at  $-40^{\circ}\text{C}$  in Anchorage. Forty-two (42) specimens collected for taxonomic purposes have been identified (6 species), transferred to an alcohol solution, and archived in the Alaska Office.

### Analysis of Fisheries Data

The relative abundance of the catch at each station was described in units of fishing effort and fish density. Catch per unit of effort (CPUE) was expressed as total number of fish/10-minute set, and, in special cases, as kilograms of fish/kilometer. Fish densities were estimated by computing the volume ( $\text{m}^3$ ) of water fished per 10-minute set (distance towed [m]  $\times$  area of townet mouth [ $\text{m}^2$ ]) and standardizing the catch to numbers/1,000  $\text{m}^3$ .

Length frequency analyses were performed to determine size and approximate age composition of the species sampled. Fork length frequencies were plotted for the major species captured by total numbers captured or subsets of this as appropriate to the analysis being performed.

The catch (C) was summarized using standard descriptive and graphical techniques available on Statgraphics Version 4.0 software. The coefficient of variation (CV) was calculated (as in Zar 1984) to describe the relative variability of the station catches and to serve as an indicator of spatial patchiness in selected species' distributions. The classification of CV values used by Grossman et al. (1990) was adapted to indicate patchiness by station, between transects, and with relative distance from shore as:

CV $\leq$ 25%	Even distribution;
25% < CV $\leq$ 50%	Relatively even distribution;
50% < CV $\leq$ 75%	Moderately patchy; and
CV > 75%	Highly patchy.

After initial testing for normalcy, the catch data were transformed  $[\ln(x + 1)]$  for conduct of parametric tests. Green (1979) indicates that species abundance data are nearly normally distributed and recommends the logarithmic transformation. The significance level for all statistical tests was 0.05 ( $\alpha = 0.05$ ). Relative abundance (mean CPUE and density estimates) was associated with environmental parameters (e.g., mean temperature and salinity conditions of the upper 2 m) using multivariate frequency plots and univariate correlation analyses for the major species studied. The frequency presentations would demonstrate modal temperature and salinity preferences of the fish species captured offshore.

## Arctic Cisco

**Time and area.**—Logistical constraints and environmental factors precluded routine use of *a priori* random sampling design for statistical testing. Thus, statistical partitioning of the data into time, area, and size class categories was performed after the data had been collected. Emphasis was placed on relative abundance and distributional attributes of the juvenile catch. Relationships examined by one- or two-way ANOVA included variations in (1) mean fish size stratified by relative distance from the coast; (2) mean stratum fish size with respect to time; (3) mean stratum CPUE in relation to sampling time; and (4) mean station CPUE and transect CPUE with respect to time, distance, and habitat (defined by hydrographic properties averaged over the upper 2 meters). Means and their 95% confidence intervals (CI) were calculated using standard descriptive techniques. A non-hierarchical clustering procedure (Hair et al. 1987) was used to investigate salinity and temperature influences on juvenile distribution and abundance  $[\ln(\text{density} + 1)]$ . Analysis of variance and the Tukey multiple comparison tests were employed to further examine habitat relationships within the cluster groups.

**Size and growth.**—Changes in mean fork length were examined over time to evaluate seasonal growth rate in young-of-the-year fish (<100 mm). Relative rates of increase in mean length (%) and instantaneous rates of weight increase (G) were calculated after Ricker (1975). In order to compute G it was necessary to know the b coefficient (slope) from the weight-length relationship for Arctic ciscoes. The coefficient described for Arctic ciscoes in 1990 by the FWS Fishery Assistance Office in Fairbanks was used in this analysis where:

$$b = 3.22 \text{ for July 10 through July 24; and} \\ b = 3.19 \text{ for August 28 through September 12.}$$

Because growth in the young fish is exponential ( $W = aL^b$ ) the instantaneous increase in weight is described by the expression:

$$G = 100 [b (\ln l_2 - \ln l_1)/t]$$

where G is the instantaneous rate in weight increase (%) during the time period t, in this case on a per day basis. For these calculations  $l_1$  and  $l_2$  represent mean fish length at the beginning and end, respectively, of time period t.

**Migration.**—The migration rate of young-of-the-year fish was examined as a function of winds and nearshore currents using vector analysis techniques. Both passive and directed wind-aided migrations were hypothesized. A passive migration was premised on the assumption that the direction of fish movement is always with the current, whereas in the

directed migration the swimming direction is always to the west. The sustained, or voluntary, swimming speed of the fish was assumed to be 1 body length per second.

It was further assumed that the Beaufort Lagoon wind record was representative of winds along the eastern Beaufort Sea in 1990. This record was relied on exclusively to estimate nearshore currents (see Physical Oceanography in Results section) because of its length—extending from July 21 to September 10. By comparison, the Camden Bay record was too short (July 20–31) to be effectively incorporated into the migrational study. Mean daily nearshore current components ( $u_c$  and  $v_c$ ) were calculated as 4% of the mean daily wind speed ( $-0.04 \times u$  and  $v$  where negative sign corrects wind direction in the oceanographic sense, i.e. an east wind has a positive  $u$  component but the current it produces is in the negative  $u$  direction).

The mean daily speeds (in km/d) and directions of passive and directed migration were computed as follows:

Passive: It is assumed that the fish moves with the wind-driven current. Therefore, a fish's

u component ( $u_f$ ) is  $S \sin (Ac)$ ;

v component ( $v_f$ ) is  $S \cos (Ac)$ ;

Total u component ( $U_p$ ) is  $u_c + u_f$ ;

Total v component ( $V_p$ ) is  $v_c + v_f$ ;

Daily migration rate ( $M_p$ ) is  $\text{SQRT}(U_p^2 + V_p^2)$ ; and

Mean daily direction of migration is  $\text{ARCTAN}(U_p/V_p)$ ;

where  $S$  is the fish's mean daily swimming speed and  $Ac$  is the mean daily current direction.

Directed: It is assumed that the fish always moves alongshore to the west. Therefore, a fish's

u component ( $u_f$ ) is  $S$ ;

v component ( $v_f$ ) is 0;

Total u component ( $U_d$ ) is  $u_c - u_f$ ;

Total v component ( $V_d$ ) is  $v_c$ ;

Daily migration rate ( $M_d$ ) is  $\text{SQRT}(U_d^2 + V_d^2)$ ; and

Mean daily direction of migration is  $\text{ARCTAN}(U_d/V_d)$ .

**Onshore-offshore comparisons.**—An exploratory analysis was undertaken to examine catch comparisons between active and passive gear types, the latter being more common to the arctic coast. In order to determine whether the inshore (fyke-net) catches differed greatly from those offshore (towntnet), a simple linear regression analysis was performed relating CPUE and instantaneous density distributions (expanded to hourly catch rates) reported in the towntnet catches. Fyke-net catches (catch per day) were acquired from FWS (Fairbanks) and LGL Research Associates (Anchorage) for selected dates on which towntnetting had been conducted. Fyke-net catches were adjusted to an hourly basis assuming the 24-h catches reflected a constant migration rate.

A bivariate response surface analysis (Menke 1973; Grant 1986) was used to further investigate dynamic relationships between relative abundance and thermal conditions at selected onshore and offshore fishing sites. Fish density was used as an index of relative abundance. Temperature was selected as the independent variable that elicits the catch

response because of its suspected great influence on distribution and abundance. Depth-averaged temperatures for the upper 2 m were used as a proxy for environmental temperature at the time of capture.

The response surface plot represents the three-dimensional surface for the function describing the relationship between density and water temperature. A matrix of  $Z$  values (normal deviate) describes the response of the density-temperature relationship. Parameters of the normal distribution were estimated using standard descriptive techniques (i.e., means and their standard deviations). The quantitative response to the bivariate normal function was obtained by converting the calculated  $Z$  to a random variate from the specified distribution of interest using the formula:

$$X = u + \sigma Z$$

where  $X$  is the normal random deviate from the specified normal distribution of interest,  $u$  is estimated by the mean, and  $\sigma$  is estimated by the standard deviation of the mean.

## Arctic Cod

**Population and biomass.**—Area swept methods (Bakkala and Smith 1978) were used to describe Arctic cod population abundance and biomass parameters in Camden Bay. With respect to relative abundance portrayals, these methods were slightly modified to reflect "volume swept" approximations. There are several reasons for this. First, in our sampling, we are more confident about the effectiveness of the gear in the vertical, rather than the horizontal, dimension. Second, the expression of fish density as a standardized unit volume is viewed as the most quantitative index of CPUE. Finally, unlike many demersal fish surveys where most of the fish being sampled occur on a horizontal plane along the bottom, Arctic cod apparently occur throughout the water column, and volume swept methods more adequately address this vertical distribution.

Ninety-five percent (95%) confidence intervals were computed for population standing stock and wet-weight biomass as described by Bakkala and Smith (1978). Sample sizes were based on the computation of effective degrees of freedom as estimated by the Cochran (1963) approximation.

## Analysis of Physical Data

CTD data were quality controlled and processed by conventional methods. Sensor readings were converted to engineering units and sorted by pressure into 0.1-m bins. Data in each bin were then averaged to derive the temperature and conductivity for that pressure. Empty bins were filled by a value derived from interpolating between closest non-empty bins. Salinity and density were calculated and stored along with temperature as a function of depth. Data recorded during the lowering of the instrument were primarily used for analyses in order to minimize turbulence-induced effects on the sensors due to the passage of the instrument through the water. Plots of temperature, salinity, and density ( $\sigma_t$ ) versus depth and plots of temperature vs. salinity were examined to detect erroneous data and to categorize stations in terms of thermohaline structure and properties. Due to the frequency of suspected erroneous readings at the sea surface, 1-meter data are used in the following



discussions to represent conditions at the sea surface (see Thorsteinson et al. 1990). Selected profiles were used to construct contoured vertical temperature and salinity sections to show the temporal and spatial distribution of these parameters.

Data from the meteorological stations at Camden Bay and Beaufort Lagoon were in electronic format and had been converted to engineering units; however, the data had not received any preliminary processing. Each station sampled wind speed every 5 minutes and, every 0.5 hour, recorded values for average wind speed (mph), maximum wind speed, temperature (°C), and barometric pressure (in-Hg). Of these we were interested in resolving the wind vector into its alongshore and cross-shore components. This was accomplished by rotating the coordinate system such that the alongshore axis (u component) lay parallel to the coastline while the cross-shore axis (v component) lay perpendicular to the coast. An observer standing at the origin of the rotated coordinate system, looking offshore with his shoulders parallel to the coastline, would have the positive u axis on his right and the positive v axis directly in front of him.

Initial processing of the wind data consisted of several steps. First, the wind speeds were converted from mph to m/s. Next, wind angles were computed for the rotated coordinate system according to the following:

$$A_i = B_i - C_i - D_i \text{ where:}$$

$A_i$  = Wind angle in the rotated coordinate system.

$B_i$  = Wind angle as recorded by the instrument.

$C_i$  = Angular difference between the instrument's north and magnetic north (MN). This value is positive if MN is rotated clockwise from the instrument's north and negative if counterclockwise. In Camden Bay, the instrument's north was 27° E of MN while in Beaufort Lagoon it was 22° E.

$D_i$  = Angular difference between MN and north of the rotated coordinate system. The sign convention is the same as above.

At the location of the meteorological station in Camden Bay, the coastline is oriented approximately parallel to true east-west; thus, the rotated coordinate system coincides with true directions and  $D_i$  is just the magnetic variation measured in Camden Bay (32° E). The Camden Bay correction angle is therefore:

$$A_i = B_i - (-27^\circ) - (-32^\circ) = B_i + 59^\circ.$$

At Beaufort Lagoon, the coastline is oriented approximately 133°–313° true or 98°–278° magnetic. The rotated coordinate system is 8° clockwise from MN and the correction angle for Beaufort Lagoon is:

$$A_i = B_i - (-22^\circ) - (8^\circ) = B_i + 14^\circ.$$

The alongshore (u) and cross-shore (v) components were then computed as:

$$u = S \sin (A_i), \text{ and}$$

$$v = S \cos (A_i),$$

where S is the wind speed in m/s and  $A_i$  is the wind angle computed above. Mean daily values for u and v were used to compute the mean daily wind speed and direction using descriptive and circular statistics methods described by Zar (1984).

In the shallow, nearshore regions of the Beaufort Sea, currents generally run in the direction of the wind at 3 to 5 percent of the wind speed. Therefore, in this report, current speeds are estimated at 4 percent of the wind speed and are converted to units of cm/s.

## RESULTS

### Distribution of Effort

Fish sampling was conducted at 28 stations on 13 days beginning on August 2 and ending on September 5, 1990 (Fig. 4). On-station activities involved completion of a CTD cast prior to the conduct of replicate townet sampling. Only one townet sample was completed at WD22 on September 5; replicate sampling was achieved at all other stations. While fishing, the net was towed at an average speed of 0.5 m/s over a mean distance of 317.67 m. An average volume of 3,873.61 m<sup>3</sup> was strained during each tow.

Seventy-seven townet sets were successfully completed in 1990. The temporal sequence of the 1990 fishing season is depicted in Figure 5. Each point in the figure reflects a townet set, thus the cumulative effort is shown. Fish tracking experiments and poor weather conditions resulted in numerous disruptions to the sampling schedule and these are apparent in the graph. The Collinson Pt. and West Dock transects received the heaviest sampling effort in 1990. The fishing effort was distributed, in decreasing order, as follows: Collinson Pt., 23 sets; West Dock, 18 sets; Endicott and Arey Island, 10 sets each; and Brownlow Pt., 8 sets. Four sets (each) were completed off Carter Creek and at Mary Sachs Entrance to eastern Stefansson Sound. While all of the stations were sampled at least once during the summer, the coverage was inadequate both in time and space for rigorous analysis of fish migration through the study area. Depth and distance characteristics of the sampling are shown in Figures 6-9. The townet was fished in depths ranging from 2.0 to 25.8 m and from 0.32 to 15.25 km from the mainland coast. In this instance, the 15.25 km reflects the distance from the southern shore of Simpson Cove to station CP15. If Collinson Pt. is assumed to be the mainland, the distance is about 12 km to station CP15.

### Physical Oceanography

Hydrographic data (CTD) were collected at 39 stations in association with the townet sampling. A solitary CTD cast was completed at the Endicott base station (EC10) on September 4; wind and sea conditions prohibited further sampling on this date. Vertical temperature and salinity plots for the CTD casts are provided in Appendix A. Sea surface and bottom water temperatures ranged between 1.03 and 6.05°C ( $\bar{x}$  = 3.6, SD = 1.2) and 0.11 and 6.05°C ( $\bar{x}$  = 2.8, SD = 1.3), respectively, during the sampling period. The mean water temperature of the upper 2 m was 3.6°C (SD = 1.2) for all stations combined. Surface salinities ranged from 23.5 to 32.5 ppt ( $\bar{x}$  = 29.8, SD = 2.5) and bottom salinities between 23.5 and 32.3 ppt ( $\bar{x}$  = 31.1, SD = 1.5). The average salinity of the upper 2 m was 30.2 ppt (SD = 2.0).

The salinity ranges described above are indicative of a primarily marine environment. Plots of the 1-meter and bottom temperature and salinity values obtained during the fish




Figure 4.—Distribution of townet sampling effort by date and location.




Figure 5.—Distribution of fishing effort by date and location.




Figure 6.—Percent fishing effort by depths sampled on the coastal transects.




Figure 7.—Percent distribution of sampling effort by distance from the coast.




Figure 8.—Townet station depths along coastal transects in the Beaufort Sea.





Figure 9.—Location of nearshore sampling locations with distance from the coast.



**Figure 10.**—One-meter temperatures versus one-meter salinities.

sampling confirm this condition (Fig. 10). The apparent warming of the marine waters was probably due to insolation. The temperatures and salinities of surface waters are also indicative of a relatively homogeneous environment for fish. Observed temperature and salinity differences between surface and 2-m depth intervals were small. Temperature differences ranged from  $-0.34$  to  $1.22^{\circ}\text{C}$  ( $\bar{x} = 0.11^{\circ}\text{C}$ ). Salinity differences were more pronounced ( $\bar{x} = -0.43$  ppt; range  $6.20$  to  $-6.21$  ppt). However, large salinity gradients were observed at very few stations; the majority of data indicate a well-mixed upper layer.

Prior to August, brackish influences may have been greater in the nearshore areas sampled. If coastal water was present early on, our data suggests that it had been washed out quickly by the prevailing east wind conditions. Easterly wind conditions were dominant in Camden Bay during the latter part of July (Fig. 11). Meteorological data collected at Beaufort Lagoon reflect a dominance of northerly and southwesterly winds in this sector of the eastern Beaufort Sea; conditions that would promote the transport or presence of marine waters in nearshore areas (Fig. 12). (It should be remembered that the time series shown in Figure 12 are in the rotated coordinate system).

Daily mean time series of wind speed and direction of the wind records were computed for Camden Bay and Beaufort Lagoon (Figs. 11 and 12). Mean vector winds from the east prevailed for 8 of the 12 d for which wind data were collected in Camden Bay. The easterly winds had daily average speeds of 3.31 to 8.22 m/s. Mean daily winds for July 20–31 were from a northwesterly direction with daily mean speeds of 0.56 m/s and 5.08 m/s, respectively. The remaining 2 d had mean winds from the north with mean speeds of 2.0 m/s (July 22) and

TIME SERIES OF SPEED, DIRECTION, AND E-W AND N-S WIND COMPONENTS  
CAMDEN BAY STATION MET STATION: CBAY DELTA T(MIN) 30.





Figure 11.—Time series of speed, direction, and E-W and N-S wind components in Camden Bay.

TIME SERIES OF SPEED, DIRECTION, AND E-W AND N-S WIND COMPONENTS  
BEAUFORT LAGOON STATION MET STATION: BFLG DELTA T(MIN) 30.



-27-

Figure 12.—Time series of speed, direction, and E-W and N-S wind components in Beaufort Lagoon.



3.53 m/s (July 29). The hydrographic profiles for CTD stations occupied in Camden Bay during late July (Appendix A) indicate nearshore salinities exceeding 27 ppt.

The wind record from Beaufort Lagoon shows that during July and August the strongest winds were from the northeast: the mean daily wind speed was 5.51 m/s from the northeast on July 28; 8.62 m/s on August 5; 10.52 m/s on August 15; and 5.97 m/s on August 25. Hydrographic data from CP11 (equals CP01 in Appendix A) shows nearly isohaline conditions of around 30 ppt just offshore of Collinson Point.


A sustained period of northwesterly winds occurred between July 21 and 26. The winds were strong (7.22–7.84 m/s) to moderate (3.03–3.90 m/s) and probably sufficient to depress marine conditions offshore (i.e., piling of coastal water against the coast). The strongest westerlies occurred in early September (8.85–9.10 m/s) after fishing had been terminated. By this time little coastal water remained in the nearshore and marine conditions were observed at all but the shallowest of stations sampled. For instance, on September 3 it appears that Katakaturuk River water had been transported into the eastern portion of Simpson Cove (CP10).

During periods of lower wind speeds the data become much noisier with respect to direction of wind. This switching of winds is apparent in the mean daily wind speed and direction records for Camden Bay and Beaufort Lagoon. The variability is much more noticeable in the plots of the raw data depicting alongshore components of the wind and nearshore currents at 0.5-h intervals. The u components have not been corrected in the oceanographic sense and therefore the high frequency of positive values is indicative of the predominance of east winds. Such conditions would promote the transport of juvenile Arctic cisco under the Mackenzie River hypothesis.

A summary of the shipboard observations of wind speed and direction at the time of sampling ( $n = 77$ ) indicated that easterly winds were reported 52% of the time; westerly winds 21%; northerly 14%; and no winds the remaining proportion of the time. These observations provide additional circumstantial evidence of prevailing east to northeasterly wind conditions that dominated the 1990 open-water season.

In several instances, the hydrographic data were collected over short enough time spans that they could be construed as synoptic. This allowed the construction of vertical temperature and salinity sections for selected times and areas during the 1990 field season. Brief descriptions of each follow:


**Endicott Causeway.**—Hydrographic data were collected on August 9 for this section (Fig. 13). The distribution of temperatures indicates the lack of a strong thermal gradient as temperatures varied by only 2°C over the 12-km transect. Colder marine water was observed at approximately 8 km from the coast. Transitional waters (< 30 ppt with temperatures of 4 to 6°C) were found inshore of 8 km. The base station, EC10, was isothermal and isohaline. Nearshore waters were nearly isothermal to a distance of 6 km offshore. At this distance there was slight stratification with indications of marine waters 8 km offshore. The salinity section shows similar trends with transitional waters lying



**Figure 13.**—Vertical sections of temperature (A) and salinity (B) from the Endicott transect, August 8, 1990.



within 8 km of the coast and salinities greater than 30 ppt beyond. The mean daily winds at Beaufort Lagoon were from the west at a speed of 4.34 m/s.

**West Dock Causeway.**—This section was reconstructed from hydrographic measurements taken over a 22-h time period on August 21 and 22 (Fig. 14). The section indicates the encroachment or upwelling of marine waters along the bottom to within about 2 km of the coast with salinities in excess of 31 ppt. It is interesting that the waters inshore of station WD21 appear to have salinities in excess of 31 ppt. Warmer transitional waters (5°C and greater and 25 ppt) have been displaced to at least 5 km offshore. The presence of brackish waters extends beyond WD25 and below the depth of fish sampling; however, salinities are high offshore representing transitional waters. The indication is that this represents the offshore deflection of the Sagavanirktok River plume. The hydrographic properties suggest east wind conditions.




**Figure 14.**—Vertical sections of temperature (A) and salinity (B) from the West Dock transect, August 21-22, 1990.

**Arey Island.**—The transect was sampled on September 2 (Fig. 15). The temperature and salinity sections indicate the presence of purely marine waters with temperatures slightly greater than 2.5°C less than 1 km from the coast. Salinities greater than 30 ppt were found nearest the shore, and greater than 32 ppt beyond 4 km of the coast. Mean daily winds from the northwest at 3.51 m/s were reported at Beaufort Lagoon on this date.



**Figure 15.**—Vertical sections of temperature (A) and salinity (B) from the Arey Island transect, September 2, 1990.



**Figure 16.**—Vertical sections of temperature (A) and salinity (B) from the Collinson Point transect on September 2 (CP11–15) and September 3 (CP10), 1990.

**Collinson Point.**—The transect was sampled late in the evening of September 2 and early morning of September 3 (Fig. 16). Marine conditions were found at all exposed sites. Transitional waters with surface temperatures of 4°C and salinities of 26 ppt were observed within Simpson Cove (CP10). A strong salinity gradient was evident at this station with 31 ppt, < 4°C water, near the bottom of the water column. Offshore waters were primarily

marine with temperatures greater than 4°C extending just beyond 1 km of Collinson Pt. and decreasing to less than 3°C about 8.5 km offshore. Sea temperatures < 3°C were found at all depths exceeding 5 m. Salinities were > 30 ppt at all stations seaward of Collinson Pt. Winds were from the northwest during the sampling period with mean daily speeds of 3.51 and 3.65 m/s, respectively. However, due to the lateness of the season, and extent of marine conditions, the effects of west winds on all but the nearshore were negligible.

**Brownlow Point.**—Hydrographic data were obtained on September 3 (Fig. 17). The sections look very similar to what was reported off Arey Island. Marine conditions were found at all stations, and, under the northwesterly winds reported for this date, Canning River influences are not expected or evident in the data.



### General Catch Summary

A total of 17,087 fish were captured by the 1990 tow-netting effort. Eleven species from eight families were represented in the catch (Table 1). Fish were captured at almost all of the sampling stations. Only nine water hauls (zero catches) occurred, and of these, several resulted from sets discontinued due to weather or mechanical difficulties. The mean number of fish caught per set was 222 (SD = 487.6, range 0–3,451). This corresponds to a mean fish density of 67 fish/1,000 m<sup>3</sup> (SD = 153.5, range 0–898.5). The species composition of the catch and corresponding mean CPUE and density indices for all but the rarely encountered species are shown in Table 2.

The largest catches were generally reported within 10 km of the coast in depths shallower than 15 m. This may reflect the distribution of the sampling effort as much as relative abundance of the fish species in nearshore habitats. Perhaps more revealing is the presentation of total catch (all species combined) according to the mean environmental conditions of the upper (2 m) water column. The majority of the catch was sampled in marine waters with temperatures of 3 to 5°C and salinities less than 31 ppt. Density distributions followed a similar pattern (Figs. 18 and 19).

The offshore sampling began midway through the open-water season. By this time, brackish water influences on nearshore areas, particularly exposed coastal environments such as Camden Bay, were not evident. Under such habitat conditions, marine species would be expected to dominate the catch. Not surprisingly, Arctic cod, capelin, and juvenile kelp snailfish (*Liparis tunicatus*) composed the bulk (almost 96%) of the fish collected offshore. The catches of juvenile Arctic cisco in marine habitats were more unexpected. Their occurrence in offshore waters is examined in more detail in the next section.

The catch results have been summarized in several tables appended to this report. Appendix B provides a summary of CPUE (catch per set) for the major species by station and date. Appendix C provides a similar summary for density information. Appendix D provides a summary of the replicate sampling by station, date and location of sampling. Standard descriptive statistics including CVs by species and total catch for each station are also provided in Appendix D. Appendix E lists the geographical coordinates of the offshore stations.



**Figure 17.**—Vertical sections of temperature (A) and salinity (B) from the Brownlow Point transect, September 3, 1990.

**Table 1.**—Scientific and common names of fish collected by townetting in 1990, coastal Beaufort Sea, Alaska.

Family	Species	Common name
Salmonidae	<i>Coregonus autumnalis</i>	Arctic cisco
	<i>Salvelinus alpinus</i>	Arctic char
Gasterosteidae	<i>Pungitius pungitius</i>	Ninespine stickleback
Osmeridae	<i>Mallotus villosus</i>	Capelin
Gadidae	<i>Boreogadus saida</i>	Arctic cod
Stichaeidae	<i>Lumpenus medius</i>	Stout eelblenny
	<i>Lumpenus fabricii</i>	Slender eelblenny
Liparidae	<i>Liparis tunicatus</i>	Kelp snailfish
Cottidae	<i>Gymnocephalus tricuspis</i>	Arctic staghorn sculpin
	<i>Myoxocephalus quadricornis</i>	Fourhorn sculpin
Pleuronectidae	<i>Liopsetta glacialis</i>	Arctic flounder

**Table 2.**—Species composition and relative abundance of the 1990 townet catch.

Species	Number caught	Percent composition	Mean CPUE	Mean density
Arctic cod	14,945	87.46	194.09	57.94
Capelin	906	5.32	4.77	4.15
Arctic cisco	560	3.27	7.29	1.83
Snailfish	521	3.05	6.77	2.48
Stickleback	97	0.57	1.26	0.33
Sculpins	36	0.21	0.47	0.12
Eelblennies	12	0.07	—	—
Arctic flounder	5	0.03	—	—
Unidentified species	4	0.02	—	—
Arctic char	1	0.01	—	—

Catch size varied widely at the 19 stations where juvenile Arctic ciscoes were sampled; CV values ranged from 0% to 142% with a mean of 67.4%. The relative abundance of juveniles was typically low and characterized by a high degree of patchiness throughout the season. This was especially true of early August (sampling through August 9) when a mean CV of 124.5% (n = 5 stations) was indicated by the catch. A moderately patchy distribution was observed throughout the remainder of the season (mean CV = 55.9%, n = 14). Similarly contagious distributions were indicated on finer temporal and spatial scales. For instance, a mean CV of 66.2% (n = 6) was calculated from West Dock catches on August 21 and 23. However, if the outer transect stations WD24 and WD25 are treated as outliers, the catch






Figure 18.—Total fish densities versus mean temperatures of the upper 2 meters.




Figure 19.—Total fish densities versus mean salinities of the upper 2 meters.

rates were much less variable (mean CV = 29%, n = 4) and the limited data suggest the migration was centered within 9 km of the coast.

Although Arctic cod were the most abundant species captured in 1990, their relative abundance was quite variable with respect to location and time of sampling (mean CV = 70.3%, range 0–142%, n = 28 stations). The townet appeared to select for small individuals (FL < 100 mm). Gear avoidance and size-related differences in the vertical distribution of the Arctic cod may be responsible for much of the observed variability. Although several instances of highly uniform fish distribution were reported (e.g. station CP01 on August 2) the overall trend was for fluctuating abundance and spatial patchiness throughout most of the summer. However, a trend of increasing abundance of YOY cod was observed late in the season. Prior to August 28, the mean CV for Arctic cod sampled at 11 stations was 89.5%. During the remainder of the season cod were sampled at 17 stations and the mean was 57.9%.

Arctic cod were especially abundant in Camden Bay during early September. On September 2 and 3 the species was present at 14 of 15 stations sampled. Station CV's ranged from 2 to 142% with an overall mean of 48.4%. In most instances, their abundance was observed to decrease with increasing distance from the coast. Other distributional patterns were evident. Arctic cod were abundant in catches taken off Arey Island and Collinson Point. Conversely, they were nearly absent in sampling conducted off the Canning River delta (Brownlow Pt. transect). With the exception of sampling at station CP10, marine conditions were encountered throughout the bay. The mean transect CV's were (from east to west) 52.6%, 26%, and 102%, for Arey Island, Collinson Pt., and Brownlow Pt., respectively.

Capelin and snailfish were also numeric dominants of the catch. Both species were captured at inshore and offshore stations, although there appeared to be a greater tendency for capelin to be distributed closer to the coast. The mean CV's were 108.8% for capelin (range 16–142%, n = 11 stations) and 86.9% for snailfish (range 5–142%, n = 17). These values are indicative of the highly contagious distributions observed for each species in 1990.

### **Species Characterizations**

Juvenile Arctic cisco and Arctic cod were dominant components of the catch. Both fishes have been identified as "key" species of the southeastern Beaufort Sea and thus much of the focus of this report has been placed on them. Other dominant species included capelin, ninespine stickleback, and juvenile snailfish. Separate accounts are provided for each of these species. The remaining species in the catch were viewed as incidental and are treated together; these species were either apparently unable to avoid the gear or were collected at shallow stations where the net accidentally touched bottom.

#### ***Arctic Cisco***

A total of 560 Arctic cisco were captured in 32 of the 77 sets completed in 1990. The earliest date of capture was reported on August 2 near Carter Creek (CC01) in Camden Bay and the latest on September 5 in Stump Island Lagoon (WD21) in Prudhoe Bay.

Juvenile ciscoes were sampled at stations on four of the five transects, in eastern Stefansson Sound (MS01), and near Carter Creek in Camden Bay. No fish were captured off Brownlow Pt. in early September; the inner station on this transect is located approximately 3 km offshore and small numbers of juveniles were likely to have been present inshore of station BP01. Juvenile ciscoes were captured at all stations on the West Dock transect on August 21 and 22. Mean CPUE and density by transect, and other areas fished, are shown in Table 3.

Young Arctic ciscoes were captured at stations located from 0.32 km to more than 12 km off the mainland coast. The catch ranged from 0 to 154 fish per set ( $\bar{x} = 7.29$ ) with corresponding densities of 0–32.3 ciscoes/1,000 m<sup>3</sup> ( $\bar{x} = 1.83$ ). Most fish were collected within 9 km of the coast and relative abundance was generally highest within the first 5 km offshore. A trend of decreasing catch size with relative distance offshore was observed. The juveniles were sampled in water depths of 2 to 10 m and were most abundant at stations of less than 6.5 m.


**Table 3.**—Arctic cisco catch summary by transect or other coastal area sampled by townet in 1990.

Sampling area	CPUE (number of fish/set)			Density (number of fish/1,000 m <sup>3</sup> )		
	Mean	SD	Range	Mean	SD	Range
West Dock Causeway	23.0	38.4	0–154	5.9	8.6	0–32.6
Endicott Causeway	1.1	1.9	0–6	0.3	0.5	0–1.5
Mary Sachs Entrance	20.8	26.4	0–55	4.0	5.1	0–10.6
Brownlow Point	0.0	—	—	0.0	—	—
Collinson Point	1.8	3.3	0–12	0.5	0.9	0–3.9
Carter Creek	0.8	1.5	0–3	0.1	0.3	0–0.6
Arey Island	0.9	2.2	0–7	0.5	1.4	0–4.5

The distribution of catch by transect fished is shown in Figure 20. More than 90% of the total catch was reported from the causeway areas. Juvenile ciscoes were especially abundant at West Dock during the second half of August and this may suggest the timing of peak migration through Prudhoe Bay.


### Size and Growth

Length frequency distributions indicate that the majority of Arctic ciscoes sampled were young-of-the-year fish (Fig. 21). The mean length was 77.2 mm (SD = 9.45, range 51–123 mm). Only one cisco was longer than 100 mm. This fish was captured near the Endicott Causeway (EC12) on August 9 and measured 123 mm long. This size is indicative of an age 1 fish. A summary of length measurements by station and date is provided in Table 4.



**Figure 20.**—Arctic cisco CPUE (catch/10-min tow) by sampling transect in 1990.

The average size of migrating fish was examined during three time periods approximating early, middle, and late portions of the field season. The early season included the townetting conducted prior to August 10 in Camden Bay and near the Endicott Causeway. Enough time had elapsed between the two sampling efforts (4–8 d) for juveniles to have moved from Camden Bay, into Stefansson Sound, and the causeway area. Thus, it



**Figure 21.**—Arctic cisco length frequencies, all stations.

is likely that the same cohort was sampled throughout the early season. The middle season included all the sampling that was conducted at West Dock between August 21 and 23. All fishing activities after this were pooled into a late season category. During the late period fishing was conducted between Camden and Prudhoe bays.

A comparison of the mean length of juveniles with time revealed a significant difference in average size with respect to time ( $F_{2,558} = 4.835, P = .0083$ ). The observed trend was for decreasing size with advancing summer (Fig. 22). The mean length of the juveniles during each time period was 82.1 mm, 77.7 mm, and 75.6 mm, respectively. Although the data are limited by the small sample size, it appears that larger-sized individuals lead the migration across the coast. Because the age 1 cisco captured off Endicott Causeway on August 9 could have been migrating with younger fish it was included in the analysis.

The size of juvenile Arctic ciscoes was also examined (one-way ANOVA) with respect to relative distance (in kilometers) of capture from the coast. Fish catches were pooled and partitioned into 11 distance categories based on the geographic location of the stations. A null hypothesis of equal population means was tested and rejected ( $F_{10,550} = 8.135, P < .0005$ ). This was not unexpected since we were (1) studying a migrating population, (2) the sampling was not synoptic, and (3) environmental conditions, especially winds, varied throughout the sampling period.


**Table 4.**—Length measurements of juvenile Arctic ciscoes collected by tow net in 1990.

Date	Station	Number of fish	Fork length (mm)		
			Mean	SD	Range
8/02	CC01	3	72.7	5.5	69-79
8/05	CP11	1	68.0	—	—
8/09	EC12	2	107.0	22.6	91-123
8/09	EC11	2	86.0	7.1	81-91
8/09	EC10	7	79.9	3.0	74-83
8/21	WD24	57	84.7	5.6	72-96
8/21	WD23	53	76.7	5.2	62-88
8/22	WD21	226	76.3	5.2	54-102
8/22	WD22	34	77.9	6.5	63-93
8/22	WD25	4	76.5	4.5	73-83
8/23	WD21	9	70.0	7.2	55-81
8/25	CP10	11	73.3	14.5	51-97
8/26	CP10	11	69.5	11.8	54-90
8/27	MS01	83	76.0	9.1	55-97
8/31	AI01	6	76.3	5.6	67-83
9/02	AI01	2	85.5	0.7	85-86
9/03	CP10	18	71.2	10.9	56-96
9/05	WD22	2	83.0	8.5	77-89
9/05	WD21	29	77.8	9.3	65-96

“Nearly synoptic” sampling was, however, conducted at West Dock on August 21 and 22. The length frequency data for Arctic ciscoes captured on these dates are shown in Figure 23. The average length of the 372 fish sampled was 77.83 mm. A comparison of mean lengths indicated a significant difference in the average size of the fish at stations located along the transect ( $F_{4,367} = 2.40, P < .0005$ ). The ciscoes tended to be smallest at inner and outer stations of the transect. The hydrographic properties observed at these stations were more marine than temperature and salinity conditions along the middle portion of the transect. The larger fish may have been able to avoid the marine conditions.

If the mean sizes noted above are true estimates of the population, seasonal growth can be evaluated. For the early season it was assumed that the ciscoes sampled in Camden Bay on August 2 represented the same cohort of migrating fish that was sampled 8 d later on the Endicott transect. A comparison of mean fork length size from the two dates suggests (1) a 1.64% relative increase in mean length per day, and (2) a 4.95% instantaneous rate of increase in mean body weight per day (G).

Since the distance between Collinson Pt. and West Dock is approximately 140 km, it would take a fish migrating at an assumed rate of 13 km/d about 10 d to travel this distance.




**Figure 22.**—Mean size of juvenile Arctic ciscoes for early, middle, and late periods in 1990.

This interval roughly corresponds to the time separating eastern and western sampling in the late season. Fish sampled in Simpson Cove (CC10) on August 25 had a mean length of 71.4 mm ( $n = 22$ ). Ciscoes sampled at West Dock 11 d later averaged 78.1 mm ( $n = 31$ ). Under the assumptions outlined above, this suggests late season growth rates of (1) 0.85% relative increase in mean length per day, and (2) 2.54% instantaneous increase in mean body weight per day (G).

### Time and Area

In order to assess temporal and areal aspects of the catch the study area was divided into eastern and western strata or "catch districts." The division, while arbitrary, was based on geomorphological characteristics of the Beaufort Sea coastline. The western portion of the



**Figure 23.**—Length frequencies of YOY Arctic ciscoes sampled at West Dock, August 21–22, 1990.


study area includes the open lagoon systems of eastern Stefansson Sound and Prudhoe Bay. The eastern district encompasses Camden Bay. By contrast, the western area is under greater estuarine influence during the open-water period.

The townet sampling was generally conducted in three nearshore areas (Prudhoe Bay, eastern Stefansson Sound, and Camden Bay) during four “time periods” in August and September. For analytical purposes the catch data were thus assembled in the following manner:

<i>Time period</i>	<i>Eastern catch district</i>		<i>Western catch district</i>	
	<i>Fishing area</i>	<i>Sampling dates</i>	<i>Fishing area</i>	<i>Sampling dates</i>
1	Carter Creek Collinson Pt.	8/02–8/05	Endicott	8/09
2	Simpson Cove	8/25–8/26	West Dock	8/21–8/23
3	Camden Bay	8/31–9/03	Mary Sachs	8/27
			Mary Sachs	9/03
4			West Dock	9/05

During 1990, the sampling was concentrated in fishing areas in the east during the early and late August periods and to the west during the mid-August and early September periods.





**Figure 24.**—Ninety-five percent confidence intervals for Arctic cisco CPUE by catch districts.

Mean CPUE and associated 95% CI are shown for each district in Figure 24 and by time in Table 5. It should be noted that Period 4 in these instances includes the replicate sampling that was conducted at station MS01 on September 3. The inclusion reflects the sorting of catch data into early August, mid-August, late August, and early September categories.

Regional comparisons of longitudinal abundance patterns were investigated (two-sample t tests; i.e.,  $H_0: \text{Diff} = 0$ ;  $H_a: \text{NE}$ ) between districts during similar time periods. This would help to characterize the strength and duration of the juvenile migration observed in the offshore catches. No significant differences were observed between the eastern and

**Table 5.—Mean CPUE and density of YOY Arctic cisco during four sampling periods in August and September 1990.**

Period*	Number of stations sampled	CPUE (number of fish/set)		Density (number of fish/1,000 m <sup>3</sup> )	
		Mean	95% CI	Mean	95% CI
1	21	0.71	-7.43 to 8.86	0.17	-1.65 to 2.00
2	13	29.46	19.11 to 39.81	7.32	5.00 to 9.64
3	6	17.50	2.26 to 32.74	3.55	0.13 to 6.96
4	37	1.57	-4.57 to 7.71	0.56	-0.87 to 1.93
Overall	77	7.29	3.03 to 11.54	1.83	0.87 to 2.78

\*Period 1 = August 2–9; Period 2 = August 21–23; Period 3 = August 25–27; and Period 4 = August 31–September 5.

western districts during periods 1 ( $t_{.05(2),19} = 2.093$ ;  $P = 0.1969$ ) or 2 ( $t_{.05(2),17} = 2.110$ ;  $P = 0.4985$ ). In the first instance, the comparison was between mean CPUE's from the central portion of central Camden Bay ( $\bar{x} = 0.36$ ,  $SD = 0.92$ ) and the east side of the Endicott Causeway ( $\bar{x} = 1.1$ ,  $SD = 3.43$ ). The second comparison was between mean CPUE's in Simpson Cove ( $\bar{x} = 5.5$ ,  $SD = 2.08$ ) and along the West Dock transect ( $\bar{x} = 29.46$ ,  $SD = 43.68$ ). The results suggest a relatively steady pulse of fish through the study area between August 10 and 25.

A significant change in mean abundance was detected between Camden Bay (including Simpson Cove) and Mary Sachs Entrance (MS01) during late August ( $t_{.05(2),32} = 2.037$ ;  $P = .0030$ ). No Arctic ciscoes were captured on the Brownlow Pt. transect or at station MS01 on September 3, thus the possibility of emigrational effects (i.e., sampling the same fish twice) was negated. The relatively large catch of juveniles at Mary Sachs Entrance on August 27 (83 fish in combined replicates) suggests the presence of a large pulse of migrating fish.

The large pulse of fish sampled in eastern Stefansson Sound on August 27 apparently had migrated through Prudhoe Bay by September 3. A comparison of the mean catch reported during period 4 and that of station MS01 from August 27 is indicative of a significant decline in fish abundance ( $t_{.05(2),9} = 2.262$ ;  $P = .0225$ ). While it remains possible that the juveniles were migrating closer inshore than our fishing, the small catches and apparent low abundance of juveniles suggests that we were sampling the tail end of the migration. However, juveniles were still present in coastal waters west of Barter Island in early September and thus the migration period extended beyond September 5.

Other temporal and spatial features of the migration were evaluated. Catch was shown to vary by time ( $F_{3,70} = 3.31$ ,  $P < .0005$ ) and location ( $H_0$ : equal means,  $F_{3,73} = 2.74$ ,  $P = .0869$ ). Such results would be expected for a migratory species. In the latter instance, the catch data were partitioned into 0–2, 2–5, 5–10, and >10 km categories (depending on station location); the corresponding pooled mean CPUE's were 6.4 fish/set ( $SE = 4.64$ ,  $n = 21$ ); 17.1 ( $SE = 5.31$ ,  $n = 16$ ); 4.7 ( $SE = 3.8$ ,  $n = 32$ ); and 0.5 ( $SE = 7.5$ ,  $n = 8$ ); respectively. The interaction between

treatment effects (time and location) was nonsignificant (two-way ANOVA,  $F_{3,70} = 2.74$ ,  $P = .1213$ ) and this may reflect the lack of environmental contrast in habitat (i.e., marine or nearly marine conditions) observed in 1990.

## Migration

Interruptions in the fishing schedule resulted in a sporadic time series for the offshore catch. The discontinuities render any attempts at assessing changes in fish abundance or size mode distributions with time difficult on all but very coarse scales (weekly or biweekly intervals). Over the study period longitudinal changes in juvenile density patterns were observed. For example, a one-way ANOVA of fish density versus time (periods as above except that MS01 sampling on September 3 was included in Period 4) demonstrates gross temporal changes in relative abundance ( $F_{3,73} = 2.74$ ,  $P < .0005$ ) and these may be attributable to fish migration (Table 5). These trends may parallel similar observations about the timing of migration from fyke-net catches along the coast in 1990.



The migration rate was examined using vector analysis techniques and both passive and directed fish migrations were considered. During 1990, the average-sized fish measured 77.2 mm (FL). This length was converted to a total length measurement of 84.2 mm and, under the assumption of a voluntary rate of movement of 1 body length/s, expanded to an average daily fish swimming speed of 7.3 km/d.

The time series for the mean daily rates and directions of passive and directed fish migrations are shown in Figures 25 and 26, respectively. If the juvenile migration was one of passive transport in wind-driven currents the average rate of fish movement (an average-size individual) would be 15.3 km/d (range 2.8–38.2 km/d). The mean direction of the migration off Beaufort Lagoon was 219° in the rotated coordinate system (i.e., the mean speed in the u direction was -9.6 km/d and -11.9 km/d in the v direction). This is indicative of a mean westerly migration with an onshore component. A mean speed of 18.1 km/d (range 3.9–39.3 km/d) was computed for fish undergoing a directed migration. The mean direction of such a migration would have been to the southwest (241° rotated, i.e., the mean speed in the u direction was -15.8 km/d and -8.8 km/d in the v direction). By comparison, a directed migration would not only have been faster, it would also have possessed a less significant onshore component than was predicted for the passive migration.



## Habitat

The townet effectively sampled the upper 2 m of the water column during each haul. Unless otherwise stated, "average conditions" shall refer to the mean temperature and salinity of this surficial layer.

The greatest numbers and densities of juvenile Arctic cisco were sampled in nearshore waters with temperatures ranging from 4 to 6°C and salinities from 27 to 29 ppt (Figs. 27 and 28). The average conditions were 4.1°C and 29.0 ppt, respectively. An exceptionally large catch of 154 ciscoes (32.26 fish/1,000 m<sup>3</sup>) was captured at WD21 on August 22. The mean temperature and salinity at the time of capture were 4.4°C and >28.5 ppt, respectively.



**Figure 25.**—Mean daily rates and mean daily directions of a passive migration.



**Figure 26.**—Mean daily rates and mean daily directions of a directed migration.




Figure 27.—YOY Arctic cisco catch versus mean temperatures of the upper 2 meters.





Figure 28.—YOY Arctic cisco catch versus mean salinities of the upper 2 meters.



**Figure 29.**—Cluster plot of YOY Arctic cisco CPUE with mean temperature and salinity conditions of the upper 2 meters.

The station data were assembled into brackish and marine habitats according to the hydrographic properties measured with each catch. The mean salinity conditions at 11 stations along the coast were brackish (< 28 ppt). The mean catch of juvenile Arctic ciscoes in brackish habitats was 7.45 fish/set versus 7.23 fish/set in marine (> 28 ppt) habitats. The mean densities within brackish and marine habitats were 1.77 and 1.84 fish/1,000 m<sup>3</sup>, respectively. Juvenile Arctic cisco abundance (CPUE) did not vary by habitat ( $F_{1,69} = 3.98$ ,  $P = 0.47$ ) but significant differences occurred between transects ( $F_{6,69} = 2.23$ ,  $.01 > P > .005$ ). Brackish water conditions were observed most frequently in the protected waters of Simpson Cove or in the nearshore areas of Prudhoe Bay most influenced by the Sagavanirktok River (e.g., EC10).

The Arctic cisco CPUE data cluster into four groupings defined by mean temperature and salinity habitat variables (Fig. 29). There are clear thermal differences separating the clusters with the exception of ten catches shared between Clusters III and IV. In this instance, the ability to discriminate between catches associated with marine salinities is not 100%. The distribution of townet sampling effort within each cluster was 14 sets in Cluster I, 14 in II, 17 in III, and 22 in IV. A general summary including average catch and environmental conditions for each cluster is provided in Table 6.

Cluster I represents a "High Temperature, Low Salinity" grouping from which 22% (123 fish) of the Arctic cisco catch was reported. These fish were captured in the most

**Table 6.**—Summary of cluster analysis including stations, sampling dates, and mean indices of relative abundance and environmental conditions for each cluster.

	Cluster group <sup>a</sup>			
	I	II	III	IV
Number of samples	14	14	17	22
<u>Average</u>				
CPUE (number of fish/set)	8.8	9.9	15.7	0.1
SE	4.5	4.2	9.6	0.1
Density (number of fish/1,000 m <sup>3</sup> )	2.6	2.5	3.7	0.02
SE	1.5	0.9	2.0	0.02
Temperature (°C)	5.4	4.2	3.4	2.6
Salinity (ppt)	28.1	29.6	30.9	31.8
Distance from coast (km)	5.9	4.6	4.4	6.0
range	2.0–11.5	1.5–9.0	0.3–12.0	2.0–11.5
Depth (m)	5.7	5.0	5.7	12.8
range	2.0–9.0	3.0–6.5	2.0–12.5	2.0–26.0
<u>Station information</u>				
Stations	WD23–25 EC10–11 CP10	WD22 EC12–13 MS01 CP10–11	WD21 EC14 MS01 CP10 CP12–13 CP02 WD22–23	WD21 BP01–04 CP10 CP12–15 CC01 AI01–04
Sampling dates <sup>b</sup> (in order of stations above)	821–822 809 802	822 809 827 903, 902	[822, 823, 905] 809 903 902 902, 805 825, 826 802 831 905	823 903 [825, 826] 902 802 902

<sup>a</sup>Ten samples shared between Clusters III and IV.

<sup>b</sup>Bracketed dates indicate sampling occurred at station on more than one date.

brackish water conditions we observed in 1990. The stations included in this cluster tend to be located nearest to the coast (i.e., at the Endicott transect or in Simpson Lagoon) or otherwise influenced by freshwater inflows. With respect to the latter, stations WD23–25 on the West Dock transect appear to have been located within the plume of Sagavanirktok River water that had been deflected offshore and to the north of the causeway on August 21–22. The sampling contained in Cluster I occurred during early and mid-portions of the August–September sampling period.



Cluster II can be characterized by its "Intermediate Temperature and Salinity" conditions indicating a greater degree of mixing with marine waters than was observed in Cluster I. Twenty-four percent of the total catch, or 138 fish, was reported from this cluster. Most catches were reported from inshore stations along the transects in late August. The exception was station EC13 which was sampled on August 9. The conditions at EC13 were transitional between coastal waters inshore and marine waters offshore.

Cluster III stations were sampled during the latter part of the migration period in late August and early September. Roughly 47% of the total catch (267 fish) was reported from stations included in this group. The water conditions of Cluster III indicated a greater amount of mixing with marine waters than Clusters I and II. With respect to the other clusters it could be designated the "Transitional" grouping. However, habitat conditions were really marine; perhaps the degree of modification of coastal water masses was slightly less than that of Cluster 4. The stations included in Cluster III were located at inner and middle segments of the survey transects. The largest CPUE's reported from Stump Island Lagoon (WD21), at Simpson Cove (CP10), and in eastern Stefansson Sound (MS01) are included in this association. The Cluster III habitat conditions existed coastally throughout the field season, although these may have been more common in late August. The largest catches may coincide with the period of peak migration. The high relative abundance of juveniles in marine or "near-marine" water that was observed at WD21 on August 23 indicates possible fish movement in the onshore currents which exist at the northern tip of West Dock under east wind conditions.

Cluster IV represents a "Low Temperature, High Salinity," or marine, association of stations. Two fish, composing less than 1% of the total catch, were captured in this habitat. Most of the stations occurring farthest offshore were grouped in this cluster. The sampling at WD21 (August 23), at CP10 in Simpson Cove (August 25 and 26), and at two stations off Collinson Pt. (CP12 and 13) is shared with Cluster III. All of the stations sampled on September 2 and most of those on September 3 occurred in marine habitats of Camden Bay. The majority of stations included in this cluster were sampled after August 22 although marine conditions existed in coastal waters throughout the sampling period.

The ten stations shared between Clusters III and IV represented about 6%, or 31, of the ciscoes captured in 1990. The average CPUE and density reported from these stations were 3.1 (SE = 0.95) and 0.75 (SE = 0.22), respectively. These data indicate that significant numbers of fish were present in marine conditions.

Results of a single factor analysis of variance rejected the multisample hypothesis of equal means (CPUE) between cluster groups ( $F_{4,72} = 2.50$ ,  $0.01 < P < 0.005$ ). A Tukey multiple comparison test (critical  $q_{0.05,72,5} = 3.977$ ) evaluated differences in the cluster means. The null hypothesis of equal means was rejected for Clusters II vs. IV ( $q = 5.806$ ), I vs. IV ( $q = 5.03$ ), and III vs. IV ( $q = 5.287$ ). This suggests the possibility of only two homogeneous groups (brackish and marine habitats) had the sampling effort been greater.

We conducted a series of pairwise correlation analyses seeking associations between Arctic cisco CPUE and various environmental variables that were measured or derived. These variables included salinity, temperature, depth, distance, u and v components of winds

and currents, as well as biological associations with other species in the catch. No strong relationships ( $r$  generally ranged from  $-0.2$  to  $0.2$ ) were found.

A bivariate response surface analysis was conducted to further investigate relationships between fish density and offshore thermal conditions. A matrix of  $Z$  values (normal distribution) describes the three-dimensional surface for the function describing the density-temperature relationship (Fig. 30). Parameters of the normal distribution were estimated by the variable means and their standard deviations. These were 1.83 fish/1,000  $m^3$  and 4.88 fish/1,000  $m^3$  for the density variable, and  $3.47^\circ\text{C}$  and  $1.19^\circ\text{C}$  for the temperature variable, respectively. Minimum and maximum values were 0 and  $32.26$  fish/1,000  $m^3$  and  $1.02$  and  $6.05^\circ\text{C}$ . The correlation between the two variables was  $r = 0.177$  ( $P = .1235$ ). For example, if we are interested in a random estimate of density at  $3^\circ\text{C}$ , we calculate  $X = 1.83 + (4.88)(Z_{.021})$ . The  $Z_{.021} = 0.4916$ , thus  $X = 4.23$  fish/1,000  $m^3$ .

### *Arctic Cod*


Arctic cod were present at most sampling locations in 1990. The average catch was 194 fish/set (SD = 466.9). This corresponds to a mean density of 57.9 fish/1,000  $m^3$  (SD = 137.3). Of general interest were numerous observations of cannibalism of smaller cod by larger cod while being held for field processing. Such observations have not been confirmed by stomach contents analysis but suggest a potential density dependent mechanism that may affect recruitment as has been described for other gadoids. It also suggests the apparent segregation of YOY cod that was observed in a single set near Collinson Pt. (CP02) on August 5. A catch of 397 fish was reported with fork lengths ranging from 25 to 37 mm. Smaller catches of similar-sized fish (20–25 mm) were noted in sampling at the inshore stations of the Endicott transect on August 9.

### **Length Frequencies**

Arctic cod ranged in length from 20 to 167 mm (Fig. 31). The average size was 50.4 mm (SD = 19.4). The length frequencies are clearly bimodal, with peaks between 35–45 mm and 80–120 mm. Age determinations were made from an existing age-length relationship for Arctic cod sampled in Prudhoe Bay (Craig et al. 1982). Cod measuring less than 60 mm were YOY fish. Those measuring 60–120 mm were probably age 1. Older fish, ages 2 and 3, are indicated by the remaining catch.

A trend of decreasing size of fish with advancing season was noted in the catch. The mean length of cod captured prior to August 15 was 68.1 mm ( $n = 260$ ). Between August 15 and 31 the mean size fell to 51.6 mm ( $n = 445$ ). In early September the mean length dropped even lower, to 45.8 mm ( $n = 1,115$ ).

An examination of the size of fish captured versus sampling distance from the coast indicated fish averaging 44.7 mm within 2 km of the coast ( $n = 582$ ), 64.9 mm between 2 and 5 km ( $n = 311$ ), 50.3 mm between 5 and 10 km ( $n = 822$ ), and 39.7 mm at distances greater than 10 km ( $n = 105$ ). All of the cod larger than 120 mm were captured within 2 km of the coast. Their comparatively low abundance, even at catch locations where the net was



**Figure 30.**—Bivariate normal surface plot of temperature (upper 2 m) and Arctic cisco density.


sampling virtually the entire water column, suggests gear avoidance. More likely, the larger fish were occurring at greater depths or inshore of our sampling effort.

### Habitat

The bulk of the Arctic cod catch was reported in waters having temperatures of 2–6°C and salinities of greater than 26 ppt (Figs. 32 and 33). The data indicate a definite preference for marine conditions by all sizes sampled.

### Standing Stock Estimates

Sampling in Camden Bay was of sufficient intensity and areal coverage on September 2–3, 1990 (over 31-h interval), to permit population size and weight estimations to be made for young Arctic cod. The catch was almost exclusively composed of YOY cod (Fig. 34). The mean size was 45 mm. According to Fruge et al. (1988), the average weight of a cod this size would be 11.6 g.



**Figure 31.**—Arctic cod length frequencies, all stations.

Although the offshore sampling was designed for other purposes, it was possible, given the transect and station locations, to partition the bay *a posteriori* into nearly equal subdivisions for estimation procedures. Two different gridding schemes were used to derive independent estimates of population size and biomass parameters. Assumptions regarding uniformity of fish distribution within Camden Bay were based on the widespread availability of marine habitat conditions and diminished variability in numbers observed in the replicated sampling conducted on these dates.

### ***Cell block method***

**Survey area.**—The cell block method involved the division of Camden Bay into four major subareas: Brownlow Point, Camden Bay, Arey Island, and Simpson Cove (Fig. 35). The single station (CP10) in Simpson Cove was assumed to represent the entire lagoonal system. Within the remaining three subareas, the northern and southern boundaries of the cells were delineated by the longitudinal lines connecting equidistantly located stations from the coast. The location of the transects was such that they bisected each subarea. Considering that the transects are located 41 km apart, areal dimensions for subareas and cell blocks were easily computed. Unless otherwise noted, the cell blocks have alphanumeric designations determined by transect name (BP, CP, and AI) and cell number (e.g., G1, G2) increasing from north to south.




Figure 32.—Arctic cod catch versus mean temperatures of the upper 2 meters.






Figure 33.—Arctic cod catch versus mean salinities of the upper 2 meters.



**Figure 34.**—Length frequency distribution of Arctic cod in Camden Bay, September 2–3, 1990.

The areal dimensions of the subareas were estimated as: Brownlow Pt., 215 km<sup>2</sup>; Camden Bay, 580 km<sup>2</sup>; Arey Island, 215 km<sup>2</sup>; and Simpson Cove, 60 km<sup>2</sup>. With respect to the northernmost cells in Camden Bay, several important assumptions were made for ease of area computation. First it was assumed that the eastern and western boundaries of the northern cell of the Camden subarea were of equal length. Second, that the northern cells in Brownlow Pt. and Arey Island subareas were assumed to be right triangles with hypotenuses equal to 21 km. A right triangular cell (85 km<sup>2</sup>) was also taken to encompass the bight of Camden Bay proper (CAM in Figure 35). The total area of the entire Camden Bay study area was conservatively estimated to be 1,070 km<sup>2</sup>.

Sampling stations were generally located 1.9, 3.4, 6.3, and 9.1 km offshore. A single station (CP10) in Simpson Cove was located about 1.85 km from the mainland shore, roughly halfway (north–south direction) across the cove. An additional station was located 11.75 km offshore of Collinson Pt. Because stations CP12, CP13, and CP14 were located approximately 4, 10, and 12 km offshore, respectively, their positions did not conform to specifications outlined for boundary determinations. Indices of fish abundance for stations located 3, 6, and 9 km off Collinson Pt. were therefore developed from (1) appropriate mean station data, or (2) regression analysis [ $\ln(\text{mean station density} + 1)$  vs. sampling distance (km)].



-58-

Figure 35.—Cell blocks in Camden Bay.

In the first case, the mean Arctic cod catches at stations CP12 and CP13 were not significantly different from each other ( $t_{.05(2)2} = 4.03, P > 0.50$ ). Therefore, assuming a uniform cod distribution, a mean density of 230 fish/1,000 m<sup>3</sup> was accepted for the 3.4 km station. In the second case, where regression analysis was employed, the relative abundance of cod located 6 and 9 km from the coast was estimated by:

$$\ln(\text{density}) = 4.15103 - (0.310024)(\text{km offshore})$$

$$\text{SE (a)} = 0.909965$$

$$\text{SE (b)} = 0.147277$$

$$\text{SE (est)} = 1.74906$$

$$r^2 = 26.97\%$$

$$n = 13$$

The estimated cod densities for these stations were 9.0 and 3.8 fish/1,000 m<sup>3</sup>, respectively. Stations where cod densities were estimated on the Collinson Pt. transect are shown as E12, E13, and E14 in Figure 35.

Mean density data were also used to estimate the relative abundance of cod in several other cells where little or no sampling occurred. Fish densities in the CAM cell were estimated as the mean of the innermost stations of the Collinson Pt. (CP10) and Arey Island (AI01) transects and the single station in Simpson Cove. The use of a weighted average in this instance reflects the similarity in habitat conditions at CP10 and AI01. The density estimate for CAM was 107.6 fish/1,000 m<sup>3</sup>. Mean density estimates from AI04 and BP04 were assumed to be representative of cells AIG1 and BPG1, respectively.

Because the transect lines bisected the coastal subareas, replicate samples occurred near the northern center of each cell. The townet catches at each station were assumed to be representative of cod densities throughout the upper 1.8 m of the entire cell. This depth reflects the vertical sampling dimension of the townet employed and was therefore used to compute the volume estimates for subareas and cells (i.e., area  $\times$  0.0018 km). The total volume of the study area was thus estimated to be  $2.12 \times 10^9$  m<sup>3</sup>.

Coastal waters near the Canning, Sadlerochit, and Hulahula rivers would appear to be excluded from the analysis. The inclusion of shallow and land areas in Simpson Cove and the bight of Camden Bay should compensate for such differences. Realistically, given the propensity of the species for marine conditions, these areas probably do not provide suitable habitat for Arctic cod. Other information used to develop population and biomass estimates is shown in Table 7.

**Population numbers.**—Population estimates were computed by (1) multiplying the average Arctic cod densities per cell by the cell's estimated volume, and (2) summing the population estimates for each subarea (Table 8). A total of  $83.6 \times 10^6$  YOY Arctic cod were estimated in the study area. By subarea, Arctic cod numbers were estimated as: Simpson Cove,  $2.7 \times 10^6$  fish; Camden Bay,  $76.8 \times 10^6$  fish; Arey Island,  $6.6 \times 10^6$  fish; and Brownlow Pt.,  $0.203 \times 10^6$  fish. Within Simpson Cove, east of a line running due south of Collinson Pt. the numbers were estimated to be  $0.7 \times 10^6$  Arctic cod.



Table 7.—Grid cell information for YOY Arctic cod in Camden Bay study area, September 2–3, 1990.

Subarea	Grid cell	Station	Number of samples	Station distance offshore (km)	Mean density (fish/1,000 m <sup>3</sup> )	Volume (×10 <sup>9</sup> m <sup>3</sup> )
Simpson Cove	SCG1	CP10	2	1.9	22.5	.12
Camden Bay	CPG5	CP11	2	1.9	166.9	.16
	CPG4	E12	4	3.4	230.0	.12
	CPG3	E13	13*	6.3	9.0	.24
	CPG2	E14	13*	9.1	3.8	.23
	CPG1	CP15	2	11.75	2.4	.22
	CAM	—	6	6.0	107.6	.15
Arey Island	AIG5	AI01	2	1.9	48.4	.08
	AIG4	AI02	2	3.4	13.1	.06
	AIG3	AI03	2	6.3	13.6	.12
	AIG2	AI04	2	9.1	1.9	.12
	AIG1	AI04	2	—	1.9	.06
Brownlow Point	BPG5	BP01	2	1.9	2.2	.08
	BPG4	BP02	2	3.4	0.0	.06
	BPG3	BP03	2	6.3	0.0	.12
	BPG2	BP04	2	9.1	0.2	.12
	BPG1	BP05	2	—	0.2	.06

\* Includes samples used in the regression analysis. The predicted values were assumed to have one degree of freedom associated with them.

The mean CPUE<sub>overall</sub> in Camden Bay for September 2–3 was estimated to be 39.4 cod/1,000 m<sup>3</sup> (83.6 × 10<sup>6</sup> cod/2.12 × 10<sup>9</sup> m<sup>3</sup>). The variance of this estimate was calculated as the weighted sum of the individual variances for each cell:

$$\text{VAR } \overline{\text{CPUE}}_{\text{overall}} = \Sigma [(V_i/V_T) * \text{VAR } \overline{\text{CPUE}}_{ik}] \text{ where,}$$

CPUE<sub>ik</sub> = mean density of species k (Arctic cod) in ith cell,

V<sub>i</sub> = volume of the ith cell, and

V<sub>T</sub> = total volume of all subareas combined.

The variance estimates for each cell are shown in Table 9. The estimates for stations E13 and E14 were developed from the entire catch set (all stations combined). The total variance of 209.62 is indicative of the low variability in cod catches observed in the replicate sampling.

Confidence intervals for the population estimates ( $\hat{P}_{ik}$ ) were computed as:

$$\hat{P}_{ik} \pm t_{.05, (2), v} * \text{SQRT}(\text{VAR } \overline{\text{CPUE}}_{\text{overall}}) \text{ where}$$

v = the effective degrees of freedom. The total population estimate was 83.6 × 10<sup>6</sup> with 15 effective degrees of freedom, and 95% confidence limits of 52.7 × 10<sup>6</sup> and 114.5 × 10<sup>6</sup>.

**Table 8.**—Estimated population of YOY Arctic cod in Camden Bay during September 2–3, 1990, using the cell block method.

Subarea	Grid cell	Estimated numbers ( $\times 10^6$ )
Simpson Cove	SCG1	2.7
Camden Bay	CPG1	0.5
	CPG2	0.9
	CPG3	2.2
	CPG4	27.6
	CPG5	26.7
	CAM	16.2
Arey Island	AIG1	0.1
	AIG2	0.2
	AIG3	1.6
	AIG4	0.8
	AIG5	3.9
Brownlow Point	BPG1	0.1
	BPG2	0.02
	BPG3	0.0
	BPG4	0.0
	BPG5	0.2

Subareal contributions to the total were: Simpson Cove, 2.7 million cod; Camden Bay, 74.1 million cod; Arey Island, 6.6 million cod; and Brownlow Pt., 203,000 cod.

**Population weight.**—Standard “area swept” procedures were used to compute the estimated standing stock weight (kg) of YOY Arctic cod within each subarea and throughout the study area. The population weight within a subarea was estimated by the expansion:

$$\hat{B}_{ik} = (A_i / \bar{p}_i) * \overline{CPUE}_{ik} \text{ where}$$

$\hat{B}_{ik}$  = weight (kg) of species k (Arctic cod in this instance) in subarea i, and  
 $A_i$  = area (km<sup>2</sup>) of subarea i, and  
 $\bar{p}_i$  = the effective horizontal path of the net (0.0068 km), and  
 $\overline{CPUE}_{ik}$  = CPUE (kg/km) of species k in subarea i.

To express CPUE in units of kg/km several conversions were necessary. First, standardized fish densities for each station were converted to “numbers of fish caught per volume fished per tow.” The average volume fished was 3,488.3 m<sup>3</sup> (SD = 682.1, range 2,274.2–4,513.4 m<sup>3</sup>). Next, the numbers of fish per set were divided by the actual distance (km) fished per tow ( $\bar{x}$  = 285.1 m, SD = 55.7, range 186.4–368.7 m) to compute “numbers per km.” Finally, CPUE’s (kg/km) were estimated for each station by multiplying the numbers of fish/km by 0.0116 kg. Variance estimates were derived as above except that area [i.e.,  $(A_i/A_T)^2$ ] rather than volume was used to weight the cell block estimates.

**Table 9.**—Variance estimates of  $\overline{CPUE}_{ik}$  in Simpson Cove, Camden Bay, Arey Island, and Brownlow Point subareas. Subareal estimates determined as the weighted sums of individual cell block variances.

Cell	Subarea			
	Simpson Cove	Camden Bay	Arey Island	Brownlow Point
SCG1	0.05			
Total	0.05			
CPG5		3.34		
CPG4		8.62		
CPG3		85.71		
CPG2		85.71		
CPG1		0.001		
CAM		25.31		
Total		200.08		
AIG5			0.75	
AIG4			0.06	
AIG3			0.05	
AIG2			0.01	
AIG1			0.00	
Total			0.87	
BPG5				0.01
BPG4				0.00
BPG3				0.00
BPG2				0.0003
BPG1				0.00
Total				0.0103
Grand total: 209.62 (SE = 14.48)				

Mean CPUE's (kg/km) and their respective variances were computed for each subarea. The estimate of variance for the entire study area was computed as the sum of the variances reported for each subarea.

Subarea	$\overline{CPUE}_{ik}$ (kg/km)	VAR $\overline{CPUE}_{ik}$
Simpson Cove	3.2	0.0008893
Camden Bay	17.8	3.75477
Arey Island	2.5	0.0139369
Brownlow Pt.	0.08	0.0011432
All Camden Bay	7.3	3.7707394

Population weight and variance [ $\text{VAR } \hat{B}_{ik} = (A_i/\bar{p})^2 * \text{VAR } CPUE_{ik}$ ] estimates were computed.

Estimates for the entire bay were obtained by summing the subareal components.

Subarea	$\hat{B}_{ik}$ ( $\times 10^4$ kg)	VAR $\hat{B}_{ik}$ ( $\times 10^6$ )
Simpson Cove	2.8235294	0.0695242
Camden Bay	151.823	$2.7316275 \times 10^5$
Arey Island	7.904412	$0.1393237 \times 10^2$
Brownlow Pt.	0.25294	1.14282
All Camden Bay	162.80387	$2.7331419 \times 10^5$

Confidence intervals for the population weight estimates were computed as:

$$\hat{B}_{ik} \pm t_{.05,(2),v} (\text{SQRT VAR } \hat{B}_{ik}) \text{ where}$$

$v$  = effective degrees of freedom ( $v = 15$ ). The total estimated biomass ( $\hat{B}_{ik}$ ) was 1,628,038 kg, with 95% confidence limits of 1,275,736 kg and 1,980,339 kg. Subareal contributions to the total estimated cod biomass included: Simpson Cove, 28,235 kg; Camden Bay, 1,518,230 kg; Arey Island, 79,044 kg; and Brownlow Pt., 2,529 kg.

**Biomass and density.**—The estimates of population size and weight allow computation of (1) fish density on a per meter basis, and (2) wet weight biomass ( $\text{g/m}^2$ ) for YOY Arctic cod. In the first instance, the estimated numbers of fish per square kilometer were reduced to numbers per square meter for each subarea: Simpson Cove,  $0.05/\text{m}^2$ ; Camden Bay,  $0.13/\text{m}^2$ ; Arey Island,  $0.04/\text{m}^2$ ; and Brownlow Pt.,  $0.0001/\text{m}^2$ . An overall density of  $0.08 \text{ cod}/\text{m}^2$  was estimated in Camden Bay on September 2–3, 1990.

Two independent measures of biomass can be derived from the  $\hat{P}_{ik}$  and  $\hat{B}_{ik}$  parameters. One method involves a simple conversion of  $\hat{B}_{ik}$  estimates to a per meter unit basis. The second approach involves (1) reduction of  $\hat{P}_{ik}$  to estimated numbers/ $\text{km}^2$ ; (2) multiplication of numbers/ $\text{km}^2$  by 11.6 g to express biomass in  $\text{g}/\text{km}^2$ ; and (3) conversion of  $\text{g}/\text{km}^2$  to  $\text{g}/\text{m}^2$ . A comparison of the two estimates is presented below.

Subarea	$\hat{B}_{ik}$ ( $\text{g}/\text{m}^2$ )	$\hat{P}_{ik}$ ( $\text{g}/\text{m}^2$ )
Simpson Cove	0.5	0.5
Camden Bay	2.6	1.5
Arey Island	0.4	0.5
Brownlow Pt.	0.01	0.01
All Camden Bay	0.8	0.9

### Zone method

Seven zones were described using the same station locations as identified for the cell block method (Fig. 36). Northern and southern boundaries of all zones, except for Zone 2, are equal to those of the grid cell method. The inshore limits of Zone 2 include a line paralleling the southern boundary of Zone 3 and extending between coordinates that would lie on the southernmost extensions of Collinson Pt. and Arey Island transects for which such conditions are met. Zone 1 is equal in area to cell SCG1 (Simpson Cove) above.




Figure 36.—Zones in Camden Bay.

With the exceptions of Zones 1 and 2, the eastern and western boundaries are defined by the Brownlow Pt. and Arey Island transects, respectively. Zonal areas were approximated using standard geometric relationships when length and width dimensions are known. Volume estimates were derived from area and gear depth characteristics to include only the portion of the water column fished. Most of the zones contained at least three stations where replicate sampling was conducted. The Simpson Cove density estimate is based on replicate sampling at one station. Mean densities (numbers of fish/1,000 m<sup>3</sup>) were computed for each zone using similar methods as for the cell blocks (Table 10). Therefore, Arctic cod densities at three stations on the Collinson Pt. transect were determined from mean station data (E12) or from a regression model relating mean cod densities to distance offshore (E13 and E14).

The seven zones encompass a total estimated area of 1,076 km<sup>2</sup> within Camden Bay. The total abundance of Arctic cod was computed within a volume of  $2.13 \times 10^9$  m<sup>3</sup> of water in the study area. Statistical procedures for estimating CPUE's and their variances, population size and weight and their variances, and corresponding confidence intervals have been described in the previous section.

**Table 10.**—Zone and station information used to compute population parameters for YOY Arctic cod in Camden Bay, September 2–3, 1990.

Zone	Stations in zone	Number of samples	Area of zone (km <sup>2</sup> )	Mean density (fish/1,000 m <sup>3</sup> )	Volume (×10 <sup>9</sup> m <sup>3</sup> )
1	CP10	2	60	22.5	.12
2	CP10, AI01, CP11	6	160	79.3	.29
3	CP11, BP01, AI01	6	156	72.5	.31
4	E12, BP02, AI02	8	123	118.3	.25
5	E13, BP03, AI03	5	238	46.6	.48
6	E14, BP04, AI04	5	230	45.9	.46
7	CP15, BP04, AI04	6	109	1.5	.22

A CPUE index corresponding to mean fish densities (Arctic cod/1,000 m<sup>3</sup>) in each zone was used to enumerate populations within each zone. These are the mean station densities presented in Table 8. The variances associated with these estimates are shown below.

Zone	VAR $\overline{CPUE}_{ik}$ (× 10 <sup>12</sup> )
1	0.0464341
2	91.875424
3	126.80971
4	213.94523
5	351.3299
6	325.29231
7	0.0199113
All Camden Bay	1,109.3188

The relative magnitude of the variances observed in most zones reflects the wide variation in apparent fish abundance that was observed between Barter Island and the Canning River delta.

Population estimates for each zone were calculated next:

Zone	$\hat{P}_{ik}$ (estimated numbers $\times 10^6$ )
1	2.7
2	23.0
3	22.5
4	29.6
5	22.4
6	21.1
7	0.3

A total of  $121.6 \times 10^6$  YOY Arctic cod were estimated in Camden Bay representing a mean area-wide density of 57.1 cod/1,000 m<sup>3</sup>. Upper and lower 95% confidence limits on the  $P_{ik}$  were 194.2 million and 49 million fish, respectively ( $v = 12$ ). The greatest densities were reported in zonal areas encompassing nearshore waters along the outer coast beaches of Camden Bay. Densities were highest within 9 km of the coast.

Table 11.—CPUE and population weight estimates for YOY Arctic cod in Camden Bay, September 2–3, 1990.

Zone	N	$\overline{CPUE}_{ik}$ (kg/km)	VAR $\overline{CPUE}_{ik}$	$\hat{B}_{ik}$ (kg)	VAR $\hat{B}_{ik}$ ( $\times 10^6$ )
1	2	3.2	0.0008893	$2.8235294 \times 10^4$	0.0695242
2	6	11.3	0.5029321	$2.6588234 \times 10^5$	278.43989
3	6	10.3	0.4991602	$2.3629411 \times 10^5$	262.70679
4	8	16.8	0.7744633	$3.0388234 \times 10^5$	253.39218
5	5	1.0	7.063252	$3.5000 \times 10^4$	865.524837
6	5	0.3	6.62207222	$1.0147058 \times 10^4$	7,575.8565
7	6	0.3	0.0012892	$4.8088233 \times 10^3$	0.33124
Total	38			$88.424995 \times 10^4$	17,023.274

The population weight of YOY Arctic cod in each zone was described by expansions relating mean zonal CPUE's (kg/km) and variances to the area (km<sup>2</sup>) fished within each zone. Estimates of each are shown in Table 11. The estimated population weight ( $\hat{B}_{ik}$ ) for all Camden Bay, with 9 effective degrees of freedom ( $v$ ), was 8,842,499.5 kg; 95% confidence limits were 589,119.4 kg and 1,179,380.5 kg. Individual zonal contributions to the total standing stock biomass were:

Zone	Biomass (kg)
1	28,235.3
2	265,882.3
3	236,294.1
4	303,882.3
5	35,000.0
6	10,147.1
7	4,808.8

**Density and biomass.**—Estimates of  $\hat{P}_{ik}$  were converted to mean densities (numbers/m<sup>2</sup>) per zone as follows: Zone 1, 0.05; Zone 2, 0.14; Zone 3, 0.47; Zone 4, 0.24; Zone 5, 0.20; Zone 6, 0.20; and Zone 7, 0.01. An overall mean density of 0.11 fish/m<sup>2</sup> was computed for Camden Bay.

Two indices of wet-weight biomass (g/m<sup>2</sup>) were developed from the  $\hat{P}_{ik}$  and  $\hat{B}_{ik}$  abundance estimates for each zone. The computation is the same as was described for similar estimates in the previous section.

Zone	$\hat{P}_{ik}$ Estimate (g/m <sup>2</sup> )	$\hat{B}_{ik}$ Estimate (g/m <sup>2</sup> )
1	0.5	0.5
2	1.7	1.7
3	5.4	1.5
4	2.8	2.5
5	2.3	0.2
6	2.3	0.004
7	0.2	0.04
All Camden Bay	1.3	0.8


### *Capelin*

Capelin were one of the most abundant marine species captured in 1990. Although they were found nearly 8 km off the coast, most were sampled within 4 km at depths of less than 3 m. Capelin appeared to be most abundant at inner stations along the Collinson Pt. (CP10), Endicott (EC10), and West Dock (WD21) transects. They were also common, but less abundant, in other coastal areas where the net swept most of the water column. The vertical distribution of the capelin may be such that it was not sampled effectively in deeper water habitats by our townet. The mean CPUE of capelin was 11.77 fish/tow (SD = 7.1, range 0–522). Average density was estimated to be 4.3 fish/1,000 m<sup>3</sup> (SD = 22.3, range 0–188.79).

### **Length Frequencies**

Capelin ranged in size from 41 to 95 mm (Fig. 37). The average size was 64.9 mm (SD = 10.5). Using the age–length relationships of Pahlke (1985), these were probably





**Figure 37.**—Capelin length frequencies, all stations.

yearling fish. Several capelin at the lower end of the range (around 40 mm long) could have been YOY fish.

Small differences in size of fish with distance captured from the coast were observed in the catch. Fish captured within 2 km had average lengths of 67.0 mm. Between 2 and 5 km the average length was 63.2 mm. Farther offshore at distances of between 5 and 8 km of the coast the mean capelin size was 73.8 mm.

A trend of decreasing size with time was also observed. The length of capelin reported prior to August 15 averaged 70.8 mm (n = 8). During the second half of the month the mean length fell to 66.1 mm (n = 77) and by early September it was 63.0 mm (n = 70).

### Habitat

Capelin were captured in both marine and brackish water habitats in 1990. No strong association with either habitat is apparent in the data. Yearling fish appear to be pelagically distributed in the nearshore waters. They were most frequently encountered in 2–6°C waters between 26 and 31 ppt. The capture of capelin at WD23 (August 21) and at AI01 (August 31) suggests their possible transport offshore in currents being deflected offshore by coastal landforms. At the latter station densities of yearling fish were particularly high (> 30/1,000 m<sup>3</sup>), which may suggest close proximity to an important habitat for the species in the Barter Island area.

## *Ninespine Stickleback*

The average townet catch contained 1.26 ninespine sticklebacks (SD = 3.1, range 0–22 fish). This corresponds to a mean density of 0.33 fish/1,000 m<sup>3</sup> (SD = 0.85, range 0–4.86).

Sticklebacks were most abundant at stations located within 2 km of the coast. They were collected at the inshore stations along all transects except for Brownlow Pt. and Arey Island. Marine conditions existed at all stations on the latter two transects on the dates they were sampled. The farthest offshore capture occurred in Camden Bay at station CC10 on August 2. The greatest abundance was observed at stations CP10 (August 3) and EC10 (August 9) when densities of about 4.5 fish/1,000 m<sup>3</sup> were reported in the catches.

### Length Frequencies

Ninespine sticklebacks ranged in size from 26 to 83 mm (Fig. 38). The mean length was 58.0 mm (SD = 10.4). No apparent trend in size could be noted with distance offshore. Inside of 2 km the mean length was 57.9 mm. Between 2 and 5 km from the coast the mean size was 58.3 mm.

Seasonal differences in mean size were apparent but small. Fish sampled prior to August 15 had a mean length of 58.5 mm. Between August 15 and the end of the month the mean size increased to 61.4 mm. In early September the mean size of the sticklebacks was 55.9 mm. Examination of otoliths would be required for age determinations.




Figure 38.—Ninespine stickleback length frequencies, all stations.

## Habitat

The ninespine sticklebacks were most frequently sampled in waters with temperatures ranging between 2 and 4°C and salinities > 27 ppt. The species was found in greatest abundance in areas nearby coastal sources of fresh water such as the Sagavanirktok River near the Endicott Causeway or the Katakturuk River in Simpson Cove. The data indicate that the stickleback is able to tolerate marine conditions.

## *Kelp Snailfish*

With the exception of Arctic cod, kelp snailfish were one of the most common pelagically distributed species of the offshore. They were sampled in mostly marine conditions that overlap those described above for the stickleback. The mean catch per tow was 6.8 fish (SD = 24.3, range 0–151). The mean density was 2.5 fish/1,000 m<sup>3</sup> (SD = 12.4, range 0–90.4).

## Length Frequencies

The snailfish ranged in length from 17 to 63 mm (Fig. 39) and averaged 26.4 mm (SD = 6.2). All were juvenile fish and probably young-of-the-year. No apparent differences in mean size were observed with sampling distance from the coast. The mean size ranged between 27.5 and 26.5 mm at distances of 2 km to more than 10 km offshore. Seasonal differences in mean size were less than 1 mm.




Figure 39.—Snailfish length frequencies, all stations.

## Habitat

As was noted above, the environmental conditions observed at the time of snailfish capture were very similar to those of the ninespine stickleback. The juvenile fish were found at offshore stations across the coast from Prudhoe to Camden bays. In the sampling conducted in Camden Bay during early September a very strong association between the juveniles and jellyfish medusae was apparent. Whether or not the relation offers protection to the fish from predators is not known. Their tadpole body shapes and small size do not indicate powerful swimming capabilities.

## Other Species

The remaining fishes comprised less than 1% of the total catch. Of these, the fourhorn sculpin was most abundant. Small numbers of arctic staghorn sculpin were lumped into the sculpin group. The average catch of sculpins was about 0.5 fish per tow (mean density = 0.12 fish/1,000 m<sup>3</sup>). Sculpins were always captured inshore at shallow stations where the net was able to sample demersal species. The sculpins ranged in length from 25 to 200 mm.

A single Arctic char was captured on August 27 at MS01. This fish measured 230 mm and was apparently unable to avoid the gear.

Five Arctic flounders were collected in Prudhoe Bay when the townet touched bottom at WD21. They ranged in size from 75 to 85 mm. Similarly, two species of eelblennies (12 fish total) were incidentally captured at stations in Prudhoe Bay (WD21) and Simpson Cove (CP10). They ranged from 70 to 80 mm in length.

## DISCUSSION

The strong east winds that characterized much of the open-water season effectively reduced the availability of brackish water habitats along the coast. Craig (1984) described a coastal band of brackish water with temperatures of 5–10°C and salinities of 15–25 ppt as being most important to anadromous fishes. Such conditions were rarely encountered in 1990. Nearly isohaline and isothermal conditions were observed at the base of the Endicott transect on August 9. Temperature and salinity were 6.05°C and 23.5 ppt, respectively. A pocket of relatively warm (>5°C), low salinity (24.5 ppt) water was located 7–9 km off the Prudhoe Bay shoreline (WD23) on August 21.

The majority of fish sampling occurred beyond 2 km from the coast. Closer to shore, hydrographic conditions may have been different. Offshore sampling began in early August about midway through the typical open-water season. Thus, the fish habitat conditions of July may have varied greatly from those observed a month later. Regardless, by August any brackish water influences on the nearshore appeared to have been lost and conditions were generally marine.

A major effect of the east winds was the marked lack of hydrographic contrast in the offshore stations sampled in 1990. East winds tend to depress sea level, draw brackish water

seaward, and cause upwelling near the land margin (Craig 1984). Only 11 of 77 stations sampled had salinities of less than 28 ppt. Of these, many were greater than 27 ppt. A truly coastal water mass was not evident in the sampling.

Depth-integrated temperatures averaged over the vertical dimension of the townet served as a proxy for environmental temperature of the offshore habitats. Only 18% of the stations sampled had temperatures exceeding 5°C, 18% were between 4 and 5°C, and 31% between 3 and 4°C. Thirty-three percent of the stations had temperatures of less than 3°C. Reynolds (1977) describes the role of temperature as a proximate (cue, guidepost, directing) factor affecting the movement behavior of fishes. The lack of temperature contrast coupled with the sporadic nature of the sampling precluded definitive statements about habitat preference based on disproportionate catches in various habitat types. The catches do, however, provide new information on the modal ranges and offshore distributions of fish not previously available. In the context of existing environmental issues and concerns this is especially true for YOY Arctic ciscoes.

The warmest temperatures were usually found at stations occurring nearest the coast and in close proximity to large river drainages. Thus, they were under greater local influence of freshwater mixing than their offshore counterparts. For example, the station in Simpson Cove (CP10) had consistently higher temperatures than were observed elsewhere in Camden Bay. This station is located about 2 km to the northeast of the Katakaturuk River. Similarly, the hydrographic conditions near the base of the Endicott transect were fresher (Sagavanirktok River influences) than those located farther offshore. In each instance, the water properties at each station were most affected during periods of high discharge or westerly winds.

In a review of fish use of coastal waters in the Beaufort Sea, Craig (1984) observed that the use of marine habitats by arctic fishes is poorly known. In fact, little information is available describing the occurrence of fish beyond 200 m of the shoreline at most locations along the coast. The 1990 sampling revealed pelagic fishes to be widely distributed with regular occurrence in catches taken 0.5 to 12 km off the coast seaward over depths of more than 10 m. The mean overall fish density (all species combined) was 67 fish/1,000 m<sup>3</sup> over the 1.8-m depth interval sampled. This is more than twice the density reported for Prudhoe Bay in August 1979 by Moulton and Tarbox (1987), whose estimate was for the entire water column and included pelagic and demersal species alike.

There are other indications that fish use of the offshore may be greater than has previously been described. Rose and Leggett (1989) described low densities of capelin in the northern Gulf of St. Lawrence as fewer than 100 fish/10<sup>5</sup> m<sup>3</sup> and very high densities as more than 400 fish/10<sup>5</sup> m<sup>3</sup>. In 1990, we reported a mean capelin density of nearly 12 fish/10<sup>3</sup> m<sup>3</sup>. The mean Arctic cod density was almost 5 times higher (58 fish/10<sup>3</sup> m<sup>3</sup>). Between  $83.6 \times 10^6$  (cell block method) and  $121.6 \times 10^6$  (zone method) YOY Arctic cod were estimated in Camden Bay during early September. The cod biomass was estimated at  $1.6 \times 10^6$  to  $8.8 \times 10^6$  kg with respect to the two methods shown. These estimates do not include older cod, which apparently have a different depth distribution than younger fish. The abundance of large numbers of YOY cod late in the season suggests an onshore movement of young from spawning areas removed from the bay.

The cell block method may provide more accurate estimates of relative cod abundance than the zonal method. Assuming that the true distribution of young cod is reflected in the catches, the use of less diverse (more compact spatially in a longitudinal sense) areas and volumes results in reduced variances in the estimation procedure. Similar methods have been used elsewhere to estimate population abundance and standing stocks of demersal fishes (e.g., Bering Sea). In the Bering Sea, population parameters for relative abundance of demersal species are generated from replicate bottom trawls in a network of survey areas (cells) with areas in excess of 1,300 km<sup>2</sup> per cell. By comparison, our estimates of Arctic cod abundance in Camden Bay derive from replicate sampling at 14 stations over a total area of less than 1,100 km<sup>2</sup>.

The results of the standing stock evaluations suggest a mean overall YOY cod density of 0.04 fish/m<sup>2</sup> (range 0.0001–0.13/m<sup>2</sup>). Craig and Haldorson (1981) estimated a run of  $19 \times 10^6$  Arctic cod into Simpson Lagoon during mid-August 1978. Their estimate was based on various indicators of coastal fish abundance from fyke net catch data and assumptions of equal dispersal of cod throughout the lagoon. The estimate of 19 million fish corresponds to a cod density of 0.12 fish/m<sup>2</sup>. This is very similar to the 0.13 fish/m<sup>2</sup> estimated for the central portion of Camden Bay in September 1990. From an ecological perspective, 0.13 fish/m<sup>2</sup> corresponds to a wet-weight biomass of 1.5 g/m<sup>2</sup> or 0.15 g dry weight/m<sup>2</sup> (mean weight of average fish was 11.6 g). Assuming that the daily rate of primary production is 1 gC/m<sup>2</sup> in Camden Bay, and a 10% transfer efficiency between trophic levels occurs, this implies sustainable biomass levels of about 0.1 for L2 (copepods) and 0.01 for L3 (secondary) consumers.

The habitat conditions observed at West Dock between August 21 and 23 indicate the upwelling of cold (<3.5°C), marine (salinity > 31 ppt) water near the coast (WD21). Warmer (> 5°C), less saline (<28 ppt) waters were observed farther offshore. These conditions are not unlike those described for the deflection of the Sagavanirktok River plume by the West Dock causeway under east winds (Hale et al. 1989). Wind data from Beaufort Lagoon indicate that a major storm event (east wind) had occurred just prior to our sampling at West Dock. However, since those data reflect wind conditions 220 km to the east, they cannot be extrapolated to the local conditions in Prudhoe Bay. Without an examination of the local wind record from Deadhorse, the mechanism responsible for the freshening of offshore waters remains speculative.

Existing oceanographic studies of the coastal Beaufort Sea suggest that coastal currents lag behind wind conditions by a period of 3–4 h (e.g., Hale et al. 1989). Data obtained from the eastern Beaufort Sea in 1989 (Hale 1991) show that upwelling occurs and reaches a depth of 2 m within 36 h of a west-to-east wind shift. If similar lag times can be assumed for changes in water properties at Prudhoe Bay, the vertical temperature and salinity sections constructed for the West Dock transect (August 21 and 22) argue for easterly winds at the time of sampling. Under strong east winds a strong onshore current is established along the west side of West Dock as westward-flowing coastal waters replace lagoonal water being displaced offshore. This current could explain the apparent high relative abundance of juvenile Arctic ciscoes inside Stump Island Lagoon (WD21) on August 21. They were the highest reported anywhere all season (32.26 fish/1,000 m<sup>3</sup>).

The high relative abundance of juvenile ciscoes in marine waters in the Stump Island Lagoon (WD21) may possibly reflect the passive migration of fish in strong coastal currents. The ciscoes may, on the other hand, be exhibiting a slight deviation from the shoreline dependency suspected of their migration and taking a less circuitous pathway into Simpson Lagoon. The catches at WD21 were among the largest and most diverse reported in 1990. This in part stems from the sampling of the entire water column at this site. Even so, the diversity and abundance of species observed is indicative of heightened biological activity. The catch of juvenile ciscoes in marine conditions may reflect a temporary departure from nearby brackish waters in response to nonthermal habitat factors (Reynolds 1977). Such factors might have included movements in response to prey or predators.

The mean CPUE's of young ciscoes captured at fyke-net stations on the east and west sides of West Dock on August 21 and 22 were 107 fish/d and 29 fish/d, respectively (pers. comm., W. Wilson, LGL, Anchorage, AK). Assuming that the fish were migrating through the area at a constant rate implies mean hourly catch rates of 4.5 fish (east) and 1.2 fish (west). In order to determine whether the inshore concentrations of fish differed greatly from those reported offshore, a regression analysis relating totnet CPUE's (expanded to hourly catch rates) and densities was performed. The resultant equation,  $CPUE_{1h} = -0.042658 + 6.04811 (\text{Density})$ , had an  $r^2$  of 99.63% and SE (estimate) of 1.583. This suggests that the fish densities inshore of WD21 were on the order of 0.75 fish/1,000 m<sup>3</sup> (west) and 0.21 fish/1,000 m<sup>3</sup> (east). Clearly, the indications are that fish abundance was much greater away from the immediate shoreline. The response surface relationship for temperature and fish densities indicates that the mean temperature was about 4°C at the fyke net stations. Apparently, the dispersal of juveniles in the marine waters west of West Dock was not related to thermal factors alone. This concurs with the findings of Bryan (1990) who concluded that currents and not thermoregulatory behaviors were primarily responsible for fish movements through Prudhoe Bay.

The documentation of young Arctic cisco catches at all of the West Dock stations in August provides new information about the offshore distribution of the species in the Beaufort Sea. At first glance, it suggests the possibility of a wider migration corridor through Prudhoe Bay (at least 11 km) than has been described. The lack of offshore sampling to the east and west of Prudhoe Bay negates such a broad generalization. Likewise, whether or not the offshore catches really provide evidence of a causeway effect cannot be unequivocally stated. Importantly, the ultimate fate of the young ciscoes occurring in marine, or near-marine, waters is uncertain. The occurrence of juveniles outside suspected modal ranges raises questions about the physiological tolerances of age 0 fish.

Relatively high YOY Arctic cisco abundances were observed at several of the stations sampled offshore on the West Dock transect. Replicate samples indicated juvenile densities ranging from 5 to 7 fish/1,000 m<sup>3</sup> some 5 km from the coast (WD22), 4-7 fish/1,000 m<sup>3</sup> at 7.5 km (WD23), 0-20 fish/1,000 m<sup>3</sup> at 9.5 km, and 0-1.5 fish/1,000 m<sup>3</sup> at more than 11.0 km offshore. This distribution may reflect the presence of more favorable habitat offshore and the fishes' movement responses to temperature and salinity optima (e.g., Neill et al. 1983). The fish may simply have been moving with optimal conditions found in currents being deflected offshore.

The offshore catches of juvenile Arctic ciscoes in marine, or near-marine, conditions contradicts existing paradigms of the species' tolerance of such temperature and salinity conditions (Fechhelm and Gallaway 1984). Existing experimental data for Arctic ciscoes indicates high mortalities to fish exposed to waters possessing salinities in excess of 25 ppt (Fechhelm et al., in press). The presence of juveniles in marine habitats that were sampled at many locations in 1990 suggests the possibility of acclimation.

The trend of decreasing mean size in juvenile ciscoes with advancing summer has been reported in fish recruiting to the Colville Delta (Moulton 1989). Apparently, larger individuals lead the migration across the coast. This relates not only to size-related differences in the swimming performance of larger fish but may also reflect stock-related differences in size and migration behavior. Instantaneous growth rates were observed to vary between daily increases of 4.95% (early) and 2.54% (late) in mean body weight during August 1990. These values fall within the observed growth ranges for pink salmon during their first few weeks at sea (Taylor et al. 1987). Like the juvenile pink salmon, young Arctic ciscoes move directly from their incubation sites to the sea in spring. Thus, the initial phases of their marine residence, particularly growth, may be comparable. During summer 1986, the instantaneous growth in migrant pink salmon was reported to vary between 2.92 and 9.68% of body weight per day (average 5.09%).

One would expect a diminishing expression of growth if the daily growth increment remains constant with increasing body size of the fish. The temporal difference observed in 1990 is probably related to availability of brackish habitats and greater accessibility to abundant foods early in the season. Presumably, higher growth rates are inherent to these warmer, less saline environs. Widespread marine conditions persisted throughout the study area in August.

English (in press) reported growth rates of 0.08% and 0.83% of body length per day (BL/d) for pen-reared Arctic ciscoes held at Niakuk and Endicott, respectively, in 1988. While interannual comparisons are tempting, it is difficult to compare the growth of free-ranging fish to that of fish held in captivity. Free-ranging fish actively search for patchily distributed prey and are able to respond to environmental changes via a multitude of adaptive behaviors. The occurrence of juveniles some distance away from the coast in temperatures outside what are considered modal may reflect localized dispersals in response to prey or other nonthermal factors (Reynolds 1977; Rose and Leggett 1989). Without knowledge of the abundance of young ciscoes and their prey in adjacent habitats, such inferences cannot be made. Although our estimates are based on small sample sizes, the juvenile length frequencies suggest relative rates of increase of about 1.6% mean BL/d in early August and 0.85% mean BL/d during the late August to early September period. If during 1990, the mean size of juveniles was about 35 mm at entry to the sea, and the coastal migration to Alaska was about 30 d, the implied relative rate of increase is 1.2% BL/d for the average-sized fish.

Preliminary catch data from several fyke-net sites in lagoonal waters of the eastern Beaufort Sea and in Prudhoe Bay provide additional information about the timing and age composition of the juvenile migration. They also provide a measure of how well temporal changes in abundance observed offshore parallel similar estimates made nearer the coast. Small numbers of juvenile Arctic cisco were captured at FWS field stations in Beaufort and



Kaktovik lagoons and in Camden Bay on July 9 (pers. comm., T. Underwood, FWS). More than 50%, or about 3,000, of the fish captured in the lagoons east of Barter Island during the second half of July were age 1 fish (>100 mm). During the same period, a significant but substantially smaller proportion (314 or about 25% of the catch) of the juveniles collected in Camden Bay was age 1 fish.

Season-long catch data of the FWS indicate that YOY ciscoes were consistently more abundant in Beaufort and Kaktovik lagoons than in Camden Bay. Our catches of young ciscoes at coastal stations off Arey Island, Carter Creek, and Collinson Point suggest a significant portion of the migration may bypass Simpson Cove and move directly across the bay. The migration corridor would thus extend approximately 2 km off the outer beaches of eastern Camden Bay. The sampling off Brownlow Point was too light and too late in the season to address fish movements along the western coast of the bay. It seems probable that the fish hug the coast after crossing the bay and migrate directly into Stefansson Sound through the shallow pass to the east of Flaxman Island.

Young-of-the-year Arctic cisco began appearing in Prudhoe Bay on August 4 (pers. comm., W. Wilson, LGL, Anchorage, AK). Catches remained high through August 11, at which time fishing operations were interrupted by a week-long storm. When fishing was resumed the catch rate rose on August 21 and remained high through September 1. All fishing was terminated on September 1. A spike was noted in the CPUE at or about August 30. This spike coincides with the large pulse of juveniles (densities of 5–10 fish/1,000 m<sup>3</sup>) observed in townetting conducted in eastern Stefansson Sound (MS01) on August 27.

Both passive and directed migrations in juvenile Arctic ciscoes adequately explain the apparent rates of fish movement observed in 1990. A purely passive migration of fish in wind-driven currents resulted in an estimated migration rate of about 15 km/d. Similarly, if the migrant was always assumed to be moving to the west, a daily migration rate of 18 km was computed. Assuming, that YOY ciscoes enter the Mackenzie estuary by mid-June, their arrival in Prudhoe Bay on August 2 indicates an elapsed travel time of 48 d. The apparent migration rate would thus be about 10 km/d (490 km/48 d). Since we know little about the early residence and movements of the juveniles at sea this estimate may be overly conservative. Our estimates would indicate travel times of between 27 and 30 d for directed and passive transports, respectively, of fish from Canada to Prudhoe Bay.

Neither rate accounts for the numerous adaptive behaviors the fish may employ to reduce losses (i.e., net daily westward movements) during the migration period. Inshore, such behaviors may include the movement of fish to protected areas behind islands and spits. Offshore, fishes might move to greater depths to avoid adverse current conditions. Such behaviors would minimize fish movements to the east, and thus could potentially increase the net migration rate to the west. The use of actual currents from moored instruments might improve this estimate. The probability of a longer migration period would also act to reduce the average daily growth rate in migrating ciscoes.

Both the passive and the directed migration scenarios resulted in a net southerly directional component to the 1990 juvenile migration. This would have the net overall tendency to hold the migration closer nearshore. Nearshore areas may offer greater fish

protection from prolonged encounters with the marine waters and hostile currents that may exist offshore. Given that the mortalities of migrant juveniles are probably high, a nearshore migration may increase opportunities for survival simply by proximity to refuge during storms or other periods of duress. A slight dispersal offshore offers abilities to utilize coastal refuges while at the same time avoiding predators common to the coast. Since both u and v current components act upon the fish movements, the inclusion of the cross-shore component is seen as a valid approach. The offsetting effect of a positive v on cisco dispersals offshore warrants greater attention. The degree of offsetting appears to be greatest during open coast migrations such as experienced by fish moving into Camden Bay.

The offshore catches at West Dock do not appear to reflect dispersal patterns of YOY Arctic cisco at other locations along the coast. The occurrence of juveniles off Arey Island (AI01) late in the season represents another area of potential offshore abundance. The geomorphology of the coastline in each area suggests that fish movements in deflected coastal waters (east wind conditions) may be involved. The cluster grouping of catches from inshore stations at Endicott and the offshore stations at West Dock tend to support this contention. The similarity in thermal conditions in the habitats at each area suggests freshwater influences of the Sagavanirktok River.

The offshore survey results demonstrated the outer portions of the coastal band and adjacent marine habitats to be important for arctic fishes. At certain locations, east wind conditions may promote the offshore movements of coastal fishes, including small anadromous species. Sustained westerly winds result in a deepening of the brackish water habitat along the coast. However, this would be offset by a narrowing of the horizontal width of this habitat because westerly winds displace coastal water against the shoreline. The narrowing of the coastal band could act to restrict the fish to areas closer to shore. Conversely, a widening of the coastal band could be expected during periods of sustained east winds. Offshore dispersals of anadromous fishes such as Arctic char and Arctic cisco would be expected under such conditions. Such excursions are indicated by Arctic char catches off the coast in Stefansson Sound in 1988 (Thorsteinson et al. 1990) and 1990 (this study). Factors, other than thermal conditions, should reasonably be expected to account for temporary departures from brackish to marine habitats where potential prey such as young cod and snailfish are abundant.

## CONCLUSIONS

The preponderance of east winds in 1990 had a profound effect on the active sampling objectives of the research. The east winds coupled with diminished coastal river runoff resulted in widespread marine conditions throughout the study area. This resulted in a lack of strong contrast in temperature and salinity attributes needed to clearly distinguish habitat use patterns in migratory or coastal fish assemblages. These conditions appear to be atypical of the "average" coastal conditions that occur during summer. Nevertheless, the indication is that anadromous fish continue to use the nearshore habitat for at least brief excursions that may be outside their preferences. One could speculate that the less than optimal conditions may produce higher than average mortalities in species such as Arctic char and Arctic cisco due to reduced fitness for overwintering.

The apparent affinity of the anadromous species for the shoreline may maximize opportunities to encounter warm, brackish or freshwater refugia at river and stream mouths. Such habitats would offer a temporary respite from adverse conditions. Following offshore feeding bouts the use of such refugia would promote rapid assimilation of energy reserves. The abundance of potential prey offshore (e.g., small cods, capelin, and snailfish) argues for such behavioral adaptation in species such as the Arctic char and Arctic cisco.

In years when the coastal band is more expansive the reliance of fish on such refugia may not be evident. The lack of sampling in the offshore Beaufort Sea in such circumstances leaves the question unanswered. Alternative to the refugium hypothesis, the apparent shoreline dependency of migratory fishes could reflect behavioral mechanisms such as searching for home stream odors or other cues that aid navigation. By contrast, the juvenile Arctic cisco migration involves highly motivated young fish that may be utilizing offshore waters to find current conditions that facilitate rapid westward movement. The frequent occurrence of young fish in marine conditions may indicate a greater physiological adaptability than previously considered possible. This hypothesis awaits laboratory confirmation. The near ubiquity of marine conditions in the coastal zone may have foreclosed any options for habitat selection.

## PART III. FISH TELEMETRY

### INTRODUCTION

Anadromous char and Arctic cisco are conspicuous elements of coastal ecosystems of northern Alaska. They are two of the most common and widely distributed apex consumers present in the Beaufort Sea during the open-water season (Craig 1984). The species form the basis of regionally important commercial, subsistence, and recreational fisheries (Craig 1989b; Moulton and Field 1989). Intensive studies of anadromous char and ciscoes by OCSEAP began about 1976 in response to concerns arising from oil and gas leasing and development at Prudhoe Bay and elsewhere in the North Slope region (Bendock 1979; Craig and Haldorson 1981; Truett 1983). During the ensuing years, many other fishery investigations have been conducted on the North Slope (Winters et al. 1988). The geographic scope of this work has extended from Point Barrow to the Canadian border. While these studies have provided a significant amount of information on the biology and ecology of char and Arctic cisco, almost all of the data have been obtained very near shore and in lagoons. The use of the more offshore waters of the Beaufort Sea by these and other fish species remains little studied, yet such information is a necessary constituent of informed assessments of the potential effects of offshore industrial activities on the species.

Prediction of the effects of habitat modifications requires basic information on the habitat requirements of a species (Winter and Ross 1982), as well as an understanding of the population responses to resources in the habitats (Hobbs and Hanley 1990). The Alaska Office's direct involvement in fish habitat use studies began through an association with the U.S. Fish and Wildlife Service (FWS), which conducted an OCSEAP-sponsored study of the genetic structure of char populations in North Slope drainages of Alaska and western Canada. The intent of that work was first to determine if there was sufficient variation in certain protein structures of the char to reliably identify the river of origin of fish and, if there was, then to determine the extent of intermixing of stocks in coastal waters. Such information was germane to several issues concerning the vulnerability of char to environmental perturbations. The FWS demonstrated that the North Slope char indeed consisted of unique populations, that little genetic interchange was evident, and that, with some exceptions, char could be quite reliably identified to river of origin.

The intermixing question could only be answered by collecting fish at several locations along the coast. Existing fishery investigations relied primarily on fykenetting next to beaches to capture fish, so no complementary information was available on the use of "offshore" habitats by char and other fishes. In response to this shortcoming, OCSEAP conducted an in-house fishery study in 1988. The primary objective of that work was to obtain char for the FWS's genetics work; however, an ancillary objective was to obtain information on offshore habitat use by char, Arctic cisco, and other anadromous and marine fishes. The results of the genetics studies are presented in Everett and Wilmot (1990), while our work is described in Thorsteinson et al. (1990). In 1990 we resumed the fish habitat use studies. The field work consisted of two main activities—townetting (described in Part II) and ultrasonic tracking, the results of which are described below.

It is necessary at this juncture to note some aspects of salmonid taxonomy having relevance to what follows. The taxonomy and systematics of the chars in general and some Alaskan chars in particular, are still being unraveled (Behnke 1980). The taxonomic status of the anadromous char occurring in the Alaskan sector of the Beaufort Sea and commonly called "Arctic char" is pertinent. McPhail (1961) identified it as the western Arctic Ocean-Bering Sea form of the Arctic char (*Salvelinus alpinus*) and defined its distribution as west of the Mackenzie River. An eastern form of *S. alpinus* occupies the arctic east of the Mackenzie River. However, a more recent taxonomic analysis by Morrow (1980) suggests that the former fish is rather the northern form of the Dolly Varden char (*S. malma*), the range of which extends through western Alaska to the Alaska Peninsula. According to Behnke, the naming of the northern *S. malma*, as well as that of the southern form and two lake-dwelling *S. alpinus* occurring in Alaska, will depend on their relationships to chars from type localities in Kamchatka and Sweden, respectively.

Our use of Morrow's classification produced naming inconsistencies in this report. Be aware that the bulk of the references to the anadromous North Slope char identify it as the "Arctic char." Inconsistencies also occur in species names of rainbow and cutthroat trouts. The rainbow trout, formerly *Salmo gairdneri*, was recently reclassified as *Oncorhynchus mykiss*, while its congener, the cutthroat trout, which was *Salmo clarki*, is now *Oncorhynchus clarki*.

Finally, it will be evident that we have incorporated many results of Canadian and European studies of *S. alpinus*. This occurred not only because the published literature on Arctic char is quite voluminous, but also because the northern Dolly Varden char and that species share similar arctic habitats. Considering the two species together also seems warranted because, as noted by Johnson (1980), there appears to be a great degree of overlap with respect to their respective ecologies. A comprehensive species account on the Dolly Varden char, including the northern form, is presented in Armstrong and Morrow (1980).

## OBJECTIVES AND APPROACH

The primary objectives of OCSEAP's telemetry studies in FY 90 were to determine: (1) movement patterns, (2) movement rates, (3) temperature occupancy, and (4) temperature preferences of large char in the coastal waters of the Beaufort Sea. The secondary objectives were to obtain similar information on large Arctic cisco.

Concurrent biological and physical data collection is emphasized in our habitat use studies to promote the evaluation of relationships between biotic and abiotic attributes in dynamic environments such as those prevailing in the coastal waters of the Beaufort Sea. Our approach to address the above objectives was to conduct an acoustic telemetry study in the Camden Bay area using a small vessel having a high precision navigation system.

Acoustic telemetry has been successfully employed in a large variety of studies of fishes, including investigations of habitat use and preferences (Tyus et al. 1984; Paragamian 1989), migration (Stasko et al. 1973, 1976; Ruggerone et al. 1990), homing (McCleave and LaBar 1972; Matthews 1990), home range (Clark and Green 1990; Matthews 1990), seasonal

activity patterns (Brawn 1982; Clark and Green 1990), diel patterns (Carey and Robison 1981; Clark and Green 1990), predation strategies (Sciarrotta and Nelson 1977; Carey and Robison 1981; Carey and Scharold 1990), swimming speed (Sciarrotta and Nelson 1977; Quinn 1988), temperature and oxygen preferences (Coutant and Carroll 1980; Douglas and Jahn 1987), vertical moments (Brill et al. 1984; Carey and Scharold 1990), thermoregulation (Carey and Scharold 1990), olfaction (Døving et al. 1985), and pollution effects (Kelso 1977; Martin et al. 1990).

Acoustic telemetry has several attractive features. It produces a large amount of data in a brief time (Matthews 1990), a major consideration in the arctic, where the field season is short and punctuated by stormy weather. Movement patterns of fish can be described in more detail than is possible with mark-recapture methods (Quinn and Brodeur 1991). Acoustic telemetry also avoids some of the problems inherent in mark-recapture studies of fish movements, such as biases introduced by nonrandom effort in tag application and recovery (Quinn and Leggett 1987). Habitat preference data obtained by telemetry also appear to be more accurate than data obtained from conventional fish collections because gear selectivity is not a factor (Tyus et al. 1984). Finally, because the organism is free-ranging, observations are not confounded by limitations often inherent in laboratory settings.

Until recently we had not seriously considered acoustic telemetry for fish studies in the Beaufort Sea due to the lack of a cost-effective, full-time navigation system having sufficient accuracy and precision. Good quality loran coverage is not available there. The implementation of the Global Positioning System, or GPS, and availability of relatively inexpensive GPS receivers now makes acoustic telemetry a practical technique in the region.

A small, shallow draft vessel is required for acoustic telemetry studies of anadromous fish in the Alaskan Beaufort Sea because the coastal waters occupied by the fish are shallow. Moreover, shallow draft enables a vessel to reach more of the few protected anchorages to avoid sea ice or escape rough seas. Small vessels are practical because their operating costs are quite low and thus a vessel can be dedicated to a tracking study for extended periods, as pointed out by Holland et al. (1985). The vessel should have a modicum of endurance and habitability due to the often inclement weather conditions in the Beaufort Sea and extended periods at sea that may be required while tracking a fish.

Several reasons underlie the selection of Camden Bay as the site for the acoustic telemetry study. First, the outer continental shelf off the Arctic National Wildlife Refuge is considered to have high potential for commercial quantities of petroleum and exploratory drilling has already taken place in the area. Second, Camden Bay has a protected anchorage. Third, the FWS has a multi-year fishery study under way in the area and agreed to furnish fish for our study, thereby obviating the need for us to expend time and effort to capture fish. We also have access to oceanographic data acquired during their study. Finally, the bathymetry of the central part of Camden Bay is somewhat atypical of the Beaufort Sea coast in that water depths are relatively deep near shore. This was an important consideration for our work as we would be unable to follow fish in very shallow water with the primary tracking vessel.

## STUDY AREA

Camden Bay is a 90-km-wide bight in the Alaskan Beaufort Sea coast named by Sir John Franklin in August 1826 in honor of the Marquess Camden, once lord of the British Admiralty (Orth 1971). The embayment lies in the western portion of the Arctic National Wildlife Refuge, between the Canning River and Barter Island, where the village of Kaktovik is located. Our fish tracking was conducted in the middle of the bight, near Simpson Cove (Fig. 40). Water depths in the study area range up to 16 m.

### Geomorphology and Geology

The shoreline morphology of Camden Bay varies considerably. Low bluffs composed of peat or unconsolidated glacial alluvium consisting of mixed silt, sand, and gravel are abundant; they are interspersed with areas in which tundra slopes gently to the upper part of the littoral zone. There is abundant evidence of active erosion in the form of unvegetated bluff faces and slumped materials at the feet of the bluffs. Beach materials consist largely of sands and gravels. The average rate of coastal erosion in the region is 1.5–1.6 m/yr (Hopkins and Hartz 1978).

The bathymetry of the eastern part of the study area is quite featureless (Fig. 40). The 5.5-m isobath is typically within 0.8 km of the shoreline east of Collinson Point, while to the west it extends several kilometers offshore, following a northwest trend beyond the point. A shallow (1–2 m), elongate shoal approximately 5 km long extends west-northwest from the emergent spit that terminates at Collinson Point and protects Simpson Cove. A narrow channel at Collinson Point separates the features. Simpson Cove and the area inshore of the shoal compose a shallow basin with a maximum depth of 4.6 m that extends westward to Konganevik Point. At depths less than about 18 m in central Camden Bay the bottom is composed of sands and muddy sands, while in deeper waters overconsolidated sandy muds and clays prevail (Barnes and Rearic 1986, p. 729). The seafloor appears devoid of boulders. Substantial amounts of peat are present on the bottom within Simpson Cove, as well as in bands along the shoreline (Dunton and Schonberg 1982).

### Meteorology

The proximity of the Brooks Range is reflected in the bimodal distributions of observed winds at Camden Bay, a consequence of orographic steering. During summer east-northeast winds dominate, while westerly winds occur when storms propagate through the region. The strength and persistence of the east winds vary considerably from year to year, so meteorologists and oceanographers speak of "weak" and "strong" east wind summers. Data collected in 1989 showed that meteorological events with periods greater than 36 h were responsible for 73% of alongshore and 43% of cross-shore wind variance, respectively. The highest wind speed recorded was 10.7 m/s from the west; however, 65% of the observations were 4 m/s or less.




Figure 40.—Study area for acoustic telemetry, Camden Bay, Beaufort Sea, Alaska, 1990.



## Hydrography and Oceanography

Many streams debouch into Camden Bay. Within the study area, the largest are the Sadlerochit River, whose delta is near Anderson Point, and the Katakaturuk, located just west of Simpson Cove. Marsh and Carter creeks enter Camden Bay a short distance east of Simpson Cove. The typical discharge pattern of streams on the North Slope consists of very high runoff during spring breakup, followed by diminution of flow through the summer to seasonal lows in winter. Records obtained in 1989 (Lyons 1990) from gaging stations on the Sadlerochit and Tamayariak, a stream just west of the study area, conform to the general pattern, but display occasional high daily flow rates that perhaps result from rainstorms or periods of hot weather. Records from the Sadlerochit showed a peak water temperature of 10.2°C in August 1989.

Astronomical tides along the Beaufort Sea coast are small, having amplitudes of only 10–30 cm. In contrast, storm surge runups may reach 3.4 m (Wise and Leslie 1988).

Meteorological, current meter, and CTD data obtained in 1988 and 1989 from Camden Bay reveal a very dynamic oceanographic regime (Hale 1990, 1991). The currents are event-dominated, with winds being the primary driving force. However, bathymetry, tides, and horizontal pressure gradients produced by runoff are not insignificant factors influencing local current patterns. Currents, temperature, and salinity were recorded both years at moorings placed approximately 0.5 and 2.8 km north of Carter Creek. Data obtained in 1989 from the offshore, near-surface current meter shows that 61% of the speeds were under 12 cm/s and that only very weakly expressed patterns were evident in current direction. The nearshore, shallow meter showed a more pronounced bimodal current direction pattern induced by its proximity to the shoreline. The major current direction mode was east-southeast, counter to the dominant easterly winds, and the secondary mode was west-northwest. Currents at both locations did not exceed 41 cm/s. The prevalent circulation pattern in the study area appeared to consist of an eddy-like eastward recirculation of water driven by a usually easterly nearshore current and density fluctuations caused by variable freshwater input.

The seasonal evolution of water properties and thermohaline structure in Camden Bay follows the regional pattern (cf. Colonell and Nedoroda 1990b). A relatively warm, brackish surface water mass forms at the shoreline in early summer. Runoff patterns and the relative dominance of east and west winds over the summer are influential in determining its development and persistence. West winds tend to keep the brackish water against the coast, while east winds move it offshore. In late summer the brackish water mass dissipates due to decreasing river runoff and air temperatures, as well as turbulent mixing produced by more frequent storms. CTD casts made in early August 1988 showed a strongly layered structure typical of the early part of the open-water season, while by late August vertical thermohaline structure was only weakly evident (Thorsteinson et al. 1990).

The temperature and salinity structure of the coastal waters in the study area displays a spectrum of variability that encompasses interannual to daily time scales. During the 1988 season the brackish water mass in Camden Bay was moderately developed; maximum temperatures and minimum salinities of 6°C and 10–11 ppt, respectively, were recorded. In 1989, the brackish water mass was relatively strongly developed; near-surface temperatures

as high as 14°C and salinities as low as 8 ppt were observed. Marked local changes of water properties and thermohaline structure occurring over very short periods have been observed in the study area during both years. Temperature and salinity have been observed to fluctuate by as much as 6–10°C and 17 ppt, respectively, in one to a few days. These fluctuations appear to result from sea breezes and strong wind events.

Sea ice coverage during summer may vary dramatically from year to year along the Beaufort Sea coast. In August 1988 it was very heavy in Camden Bay, but in 1990 it was absent.

## METHODS AND MATERIALS

### Fieldwork

Ultrasonic tracking was conducted from vessel *1273*. This boat is 11 m long and has a draft of about 1.3 m. It is powered by a Caterpillar Model 3208 diesel engine and is fabricated of aluminum. It has a maximum speed of 8 knots and a range of 730 km. Amenities include berthing for four people in the forecabin, an oil stove, a small dinette, and a head. The vessel's endurance is approximately 10 d.

### Tracking Apparatus

The electronic components of the tracking apparatus were manufactured by VEMCO, Halifax, Nova Scotia and the mounting hardware was fabricated locally in Anchorage.

The model V3T-1H transmitters used are 16 mm diameter by 62 mm long and weigh 22.4 g in air and 7.5 g in seawater. They have a power output of 152 dB re one microPascal at one m. The useful life of the transmitters is 5–9 d. Five different frequencies were selected—50.00, 60.00, 65.54, 69.00, and 76.80 KHz—in order to be able to unambiguously discriminate fish being tracked from others that had previously been tracked and might still be in the study area. Each transmitter was calibrated at the factory for temperature response. The stated accuracy of the calibration is to within 0.3°C of the actual temperature. Transmitters were activated just prior to attachment to fish by soldering together two wires extending from one end of the transmitter. The connection was covered with quick-drying epoxy to waterproof it.

The deck unit consisted of a VR-60 receiver with an 02 power supply, 01 decoder/display, and 03A data storage/logging module, all of which are enclosed in a weathertight metal case. The receiver amplifies and converts signals from the transmitters into an audible signal as well as a visual display of signal strength. It has a frequency range of 10–99.9 kHz and 900 channels are available for use. The internal power supply consists of a rechargeable 12 volt battery; it was necessary for tracking from the skiff. Individual transmitter calibration data stored in the decoder/display were used to convert transmitter pulse rates into temperature data, which were shown in a liquid crystal display as well as being available for storage in the data storage/logging module. Coupled date/time data were

provided by an internal clock. The data logger memory has a capacity of 256K bytes; it can be downloaded in the field into a laptop computer.

Two V-10 hydrophones were employed for tracking. This model has vertical and horizontal directivities between -3 dB points of 160 and 80 degrees, respectively, and a sensitivity of -148 dB re one microPascal. The primary transducer was mounted on a faired aluminum boom abeam of the cabin off the port gunwale. The shielded cable from the transducer was led through a window into the cabin. The boom extended approximately 0.3 m below the vessel's keel and was designed with a quick detachable mount in order to facilitate docking and to avoid damage to the gear if heavy ice was encountered. The selection of boom location and use of a fairing was based on the experiences of Stasko and Polar (1973) and Holland et al. (1985); these expedients were intended to minimize turbulence-caused noise at the hydrophone as well as received noise from the vessel's electronics and engine. The other transducer was mounted on a lightweight detachable boom that was installed on the 5.5-m auxiliary skiff. Tracking from the skiff occurred only when contact with the fish was lost in shoal waters and weather and sea conditions allowed safe skiff operations.

A VH-65 omnidirectional test hydrophone was used for functional tests prior to release of tagged fish. In brief, the hydrophone was placed in the fish holding tank and monitored briefly to ascertain that the transmitter was activated and that it and the receiving unit were operating properly.

## Navigation

The primary navigation instrument was a Magnavox MX 5400 GPS Satellite Navigation System, which receives transmissions from visible satellites. The GPS unit determines present position and speed over ground, as well as performing a variety of other functions. Latitude and longitude (to 0.001 minute) are shown in a visual display. The manufacturer states that the receiver is capable of position accuracies of approximately 15 m RMS. Achieving this level of accuracy requires that 3-4 satellites are visible to the receiving unit and that high quality transmissions are available from the satellites. The degraded data transmitted for public use are reported to produce position accuracies of 100 m.

Because the GPS was not yet fully operational, satellite coverage was intermittent during the 1990 field season. However, during about 61 h of tracking there were only 10 periods longer than 30 min when we were without GPS navigation or had poor fixes. Some 93 percent of the fixes were at frequencies of 10 min or less. The initial screening of GPS position data occurred during fish tracking, when obviously erroneous positions were discarded. Another onboard means of flagging possibly bad positions was the vessel ground speed output of the GPS, which occasionally indicated excessive speeds. As a check of the precision of the GPS in the working area, we casually recorded 28 fixes over a 4-h period while at anchor on August 6, 1990. Analysis of the data set gave standard deviations of latitude and longitude of 64 m and 49 m, respectively. This precision estimate is conservative as it includes some movement of 1273 about the anchor line.

A Furuno Model FR805D 48 nm radar was employed for navigation when satellites were not available. Radar fixes were taken at roughly half-hour intervals when the radar was the sole navigation instrument. The radar was also employed when the skiff was being used to track fish. Radar ranges and bearings to the skiff, as well as GPS positions and 1273's heading, were recorded at 5-min intervals for subsequent plotting of the skiff's position. Finally, while tracking fish near shore, the radar was occasionally used to determine the distance to shore.

Water depths were determined with a Ray Jefferson Color Telescan 2000 sounder.

### **Oceanographic Data Collection**

Thermohaline structure was determined at each fish release location and periodically while tracking. An Applied Microsystems CTD-12 instrument was used. The CTD is self-contained and requires no outside power or deck recorder; it stores data in an internal memory and records eight samples of temperature, conductivity, and pressure per second. The instrument's depth, temperature, and salinity sensors are accurate to within 0.10 m, 0.03°C, and 0.2 ppt, respectively. During CTD casts, the instrument was lowered over the vessel's side until the sensors were submerged, then allowed to equilibrate before commencing lowering. The lowering speed was slow (mean 5 m/min, range 3–8 m/min) to minimize turbulence around the sensors and to reduce any errors introduced by sensor response lags while passing through strong gradients of water properties. Cast depths extended to within 1 m of the sea bottom at all stations. Upon completion of a cast the CTD data were downloaded to a Compac laptop computer, reviewed for accuracy, and stored on minidiskettes for processing.

### **Tagging and Tracking Techniques**

Tests of the tracking system were conducted in Simpson Cove prior to tracking fish. We placed a transmitter on a buoy anchored in shallow water (about 2 m depth) in the east end of the cove, then proceeded toward it in 1273 to determine the aural and visual (signal strength meter) detection threshold ranges and the range at which a "loud and clear" acoustic signal was emitted by the receiver's loudspeaker. This test indicated a visual detection threshold of 0.57 km and an acoustic detection threshold of 0.45 km. A loud and clear signal was observed at distances of 0.21 km and less. In very shallow water (about 1 m deep) near shore the transmitted signal was strongly degraded and the detection threshold from the skiff was estimated to be only 0.1 km. Because temperature outliers were infrequent when a strong signal was received, we attempted to maintain close contact with fish being tracked.

Intercomparisons of ultrasonic transmitters, the bucket thermometer, and the CTD were conducted by simultaneously immersing all instruments into a container of seawater and reading each. One comparison was conducted prior to the release of Fish 3, the other prior to the release of Fish 6. They gave the following results (°C):

<i>Instrument</i>	<i>Transmitter #7191</i>	<i>Transmitter #7194</i>
Bucket Thermometer	4.6	5.6
Temperature Transmitter	4.3-4.4	5.8
CTD	4.1	5.2

The fish used for tracking were captured by FWS personnel at a fyke net located in the eastern end of Simpson Cove. Large individuals were selected in order to minimize tag effects on swimming speed and behavior (cf. Mellas and Haynes 1985). We followed (with the exception of the cisco) the guideline presented in Winter (1983) that transmitters should weigh no more than 2 percent in air of the fish's weight out of water. For the transmitters used in this study, this equates to a fish weighing at least 1,125 g. A char of that weight would be about 475 mm long.

Fish to be used for tracking were transferred to a floating pen located approximately 100 m from the fyke net and held there until needed. Normally only one or two fish were held at a time and holding times were no more than 1-2 h. One fish would be dipped from the net into a plastic container and transferred by skiff to 1273. Immediately after transfer, the fish was placed into a tagging cradle lined with a wet towel, its length measured to the nearest millimeter (FL), and then the transmitter was attached. A 6.5 cm long Petersen pin was passed through the fish's musculature immediately behind the dorsal fin. The sharpened end was then twisted with needlenose pliers to create a small pigtail loop. Earlier, the transmitter had been loosely connected to the other end of the pin by a similar pigtail around the nylon attachment loop at one end of the transmitter, thereby creating a flexible linkage. Petersen discs (2.5 cm diameter) were placed on both the entry and exit sides of the wire to reduce scale abrasion by the transmitter and the pigtailed end of the wire. The measurement and tag attachment procedure took less than 2 minutes. The fish were then placed in a large plastic fish tote (dimensions approx. 2 x 1 x 1 m) filled with seawater and allowed to recover during the approximately 1-hour trip to the release area. The fishes' condition was observed frequently during the journey to the release area. Seawater was periodically added to the container for oxygenation and to aid acclimation to conditions at the release area. Water temperatures and salinities within the cove did not differ greatly from those outside during the tracking period. The release procedure consisted of putting the fish into a smaller, covered container, transferring it to the skiff, and running the skiff about 50 m ahead of 1273 for the release to ensure that the fish entered the water gently and that it was immediately available for detection by the forward-scanning hydrophone.

Prior to releasing a fish the vessel was stopped, a GPS position determined, and a CTD cast made. Initially, the tracking data collection protocol consisted of using the data storage/logger for time/temperature data, logging at 30-s intervals. Alaska Daylight Savings Time (nearest minute), temperature (to 0.1°C), position (to the nearest 0.01 min), and corrected depth (ft) were hand logged at approximately 10-min intervals. This procedure was abandoned shortly after tracking commenced due to an apparent failure of the data logger. Manual logging of all data at nominal 5-minute intervals was performed thereafter. During tracking, the receiver's automatic gain control circuit was employed rather than manually altering the amplification of the received signal.

The most efficient method for maintaining contact with a tagged fish was a stop-and-go procedure, which consisted of slowly turning the vessel through 360° when audio signal strength weakened perceptibly, noting the bearing at which the received signal was strongest, and then proceeding slowly on that bearing until a loud, clear signal was received, at which point the vessel was stopped again. This procedure decreased the possibility of losing contact due to a marked change of direction by the fish. It was rarely necessary while tracking to exceed 1 knot to maintain contact with the fish. Of the seven fish tracked from 1273, contact with two was lost offshore, while the others were lost when they entered shoal and/or surf zone areas. On three of the latter occasions, the skiff was employed in attempts to regain contact. We successfully tracked one fish from the skiff into waters too shallow (less than 1 m) for outboard operations. When tracking fish from the skiff, an ICOM handheld VHF radio was used to communicate with 1273 and to transmit temperature data.

## **Data Analysis**

### **Telemetry Data**

Time, position, and temperature data arrays for each fish were entered into the computer for processing, conversion, and analysis. Programs written in-house were used to calculate ground speeds and to plot fish tracks, ground speeds, and temperature records. The coastline contours used for the track plots had been previously digitized from 1:250,000 scale U.S. Geological Survey maps. Many of the other analyses and graphics plots were done on an IBM PC using Statgraphics statistical analysis and plotting programs. Due to the heterogeneity of the data distributions and small (hourly-averaged) sample sizes, non-parametric methods were used for statistical testing. An alpha level of 0.05 was used for all tests of significance. The vessel's and fishes' positions were assumed to coincide.

A directedness-of-movement index was determined for each fish. The index is simply the great circle distance between the beginning and end of a fish's track divided by the total length of the track. Therefore, a value of 1 indicates completely directed movement, while a value of 0 indicates no directed movement. The direction of net movement of each fish was determined from the starting and ending positions of each track. Both unweighted and weighted Rayleigh tests (Zar 1984) were used to test for non-randomness of direction of movement of the char as a group. Moore's modification was employed in the latter test, which used net displacement as a ranking factor.

Our fish tracking generated ground speed data. Ground speed is the vector sum of the fish's swimming velocity, the ambient current velocity, and an error term representing the variable location of the fish relative to the tracking vessel as well as navigational error. Ground speeds between successive positions were calculated assuming straight-line movements. Time and distance were iteratively summed to generate gross ground speeds for entire tracks. Net ground speed, or net rate of displacement, also was calculated. It is the shortest water distance between the starting and ending positions of a fish track divided by the duration of the track.

Several methods were used to evaluate ground speed data quality. Ground speed data from individual fish were first standardized (as described below) and then screened for

anomalously high values. Our criterion for deleting data was based on the results of several experiments on the swimming performance of salmonids. Using 34-cm Arctic char, Beamish (1980) showed that approximately 1 min was required to reach 50% fatigue at 3 L/s and 5 min at 2.5 L/s. Brett (1973) determined that adult sockeye salmon had a mean critical swimming speed (the approximate maximum speed that a fish can maintain without fatiguing) of 3.1 L/s. Williams and Brett (1987) examined differences in swimming performance of migrating adult pink salmon due to sex, stock unit, and stage of maturation. They found the fish could attain critical swimming speeds as great as  $3.39 \pm 0.48$  L/s. Based on the above, we selected 3.3 L/s as a cutoff point. This cutoff includes an arbitrary 0.3 L/s (about 15 cm/s) allowance for currents. In cases where calculated L/s exceeded this value, the position data bounding that speed estimate were examined, one was deleted, and ground speeds were recalculated over the longer time interval. Zero speed values bounded by acceptable speeds were retained in the absence of any compelling reasons to question their validity.

Following data screening, gross ground speed was subtracted from each fix-to-fix speed estimate to produce demeaned speed data for each fish. These data were plotted against time to identify possible tag effects or escape responses.

Screened ground speeds were binned into 1-hour segments and averaged for statistical analyses. Linear interpolation was used when observed data points did not coincide with the bins. The averaged data were used to generate standard descriptive statistics (mean, standard deviation, range) for each fish. A Mann-Whitney U test (Zar 1984) was used to test for significant differences in offshore and alongshore ground speeds of Fishes 2 and 4. Runs tests (Zar 1984) were applied to the hourly-averaged data from Fishes 2, 4, and 6 to detect trends in swimming speeds that might reflect escape responses or tag effects. Possible diurnal variations in ground speed were only cursorily evaluated due to the absence of darkness during the study period.

Standardized ground speed data were used to compare movement rates of fish of differing sizes, to pool data from different fish for testing, and to make comparisons with results of tracking studies conducted on other species. Fork lengths were converted to total lengths using regressions derived from meristic data for char and Arctic cisco (presented in Appendix B of Thorsteinson et al. 1990). Ground speed data were then divided by the derived values to generate L/s data. A L/s vs. track duration plot was created to identify tag effects on swimming speeds. The Kruskal-Wallis test (Zar 1984) was used to test for significant differences in standardized ground speeds among fish.

The initial screening of transmitted temperature data consisted of the examination of individual time-at-temperature plots to detect obviously errant values and possible escape responses (cf. Quinn and terHart 1987). Temperature records from all char were combined to generate a temperature occupancy histogram. A similar histogram was produced for the lone tagged Arctic cisco. Temperature data from fish and CTD casts were used to evaluate temperature preferences and to estimate swimming depths of tagged fish. Spot comparisons were made for the start of each track; i.e., initial transmissions were compared to the temperature data obtained from the CTD cast taken just prior to release of a fish. Underway comparisons consisted of the comparison of coincident-in-time data from CTD casts and

transmitters, as well the transmissions during the 1-hour period bracketing a CTD cast. Hourly average temperatures and ranges were determined from the transmitter data, allowing an evaluation of the representativeness of each preference observation.

An exploratory examination of the effect of water temperature on char ground speeds was conducted using selected track segments. Segments sought were those in which temperatures remained nearly constant (less than 1°C change) over periods of at least 30 min and high quality navigation was continuously available. This was intended to minimize the potential effects of temperature gradients on behavior (see, e.g., Laurs et al. 1977) and position errors on speed estimates. Data from Fishes 2, 4, and 6, which had no significant differences in standardized ground speeds, were pooled for analysis. Seven segments were selected. An initial plot of the speed and temperature data suggested a relationship, so statistical testing followed. A Spearman rank correlation test was run on the temperature and ground speed data averaged over each of the segments. Next, the segments were broken into 50–70-min bins and average temperatures and speeds calculated as before for each bin. One 35 min segment was included because of the sparse suitable data. The average speeds from each bin, grouped by temperature, formed the data set for a Kruskal–Wallis test of intergroup differences of ground speed.

## **Oceanographic Data**

The CTD data were quality controlled and processed by conventional methods. Sensor readings were converted to engineering units and sorted by pressure into 0.1-m bins. Data in each bin were then averaged to derive the temperature and conductivity for that pressure. Empty bins were filled by a value derived from interpolating between closest non-empty bins. Salinity and density were calculated and stored along with temperature as a function of depth. Data recorded during the lowering of the instrument were used for analyses to minimize turbulence-induced effects on the sensors due to passage of the instrument through the water. Plots of temperature, salinity, and density ( $\sigma_t$ ) versus depth and plots of temperature vs. salinity were examined to detect erroneous data and to categorize data in terms of thermohaline structure and properties.

## **RESULTS**

### **Fish Tracks**

Table 12 summarizes the results of the tracking study. Six northern Dolly Varden char and one Arctic cisco were tracked for a total of 3,599 min over a distance of 119.5 km. Individual fish were tracked for periods of 140–1,260 min and distances of 6.2–36.9 km. Gross ground speeds of individual char were 48.8–74.2 cm/s, while the group's average ground speed (summed gross track length divided by summed time tracked) was 55.8 cm/s. Net ground speeds, or net movement rates, were 9.5–63.2 cm/s. The chars' individual standardized ground speeds were 0.96–1.37 L/s, while their mean standardized ground speed was 1.04 L/s. The cisco's gross ground speed was 48.9 cm/s, or 1.14 L/s, while its net ground speed was 44.1 cm/s.



**Table 12.**—Summary of results from ultrasonic tracking of northern Dolly Varden char and Arctic cisco in Beaufort Sea coastal waters, 1990.

Fish number	Total length (cm)	Start date	Duration of track (min)	Movement		Gross ground speed		Net ground speed (cm/s)
				Gross (km)	Net (km)	(cm/s)	(L/s)	
Northern Dolly Varden char								
1	53.1	7/30	166	6.9	6.3	69.7	1.31	63.2
2	54.0	7/31	935	31.0	12.0	55.3	1.02	21.4
3	48.5	8/01	292	11.5	3.1	65.5	1.35	17.7
4	56.9	8/03	1,260	36.9	7.2	48.8	0.96	9.5
5	54.2	8/10	140	6.2	5.4	74.2	1.37	64.3
6	53.5	8/11	545	19.3	5.2	59.0	1.10	15.9
1-6 combined			3,338	111.8		55.8	1.04	
Arctic cisco								
7	42.9	8/12	261	7.7	7.0	48.9	1.14	44.1

Computer drawn fish tracks illustrate results described below. Note that portions of Tracks 2 and 4 extend inside the shoreline. As these are vessel positions, the actual fish positions would be even farther ashore. The tracks were not adjusted seaward because we are confident in the quality of the GPS positions. The discrepancies could derive from four sources. The first is errors introduced during the digitization of the coastline from 1:250,000 scale USGS maps. The second is inaccuracies in the maps themselves. The National Map Accuracy Standard for well-located points on a 1:250,000 scale map is 127 m. The Mt. Michelson and Flaxman Island maps, which encompass the study area, were prepared from aerial surveys conducted in 1955 and have not been field checked. The third is coastal erosion subsequent to the mapping surveys of the area, which reportedly took place 35 years ago. Assuming that the average coastal erosion rate of 1.5–1.6 m/yr in the region noted earlier in this report is correct, the shoreline could have retreated over 50 m since the surveys. Finally, the datums used for the USGS maps and GPS system differ. The former is based on the 1927 North American Datum (NAD), while the latter uses the 1983 datum. Local differences between the two datums are about 110 m and 25 m for longitude and latitude, respectively.

**Fish 1.**—This char tracking session took place on July 30. At the beginning of the period there was a 0.3-m swell and no seas. Winds were from the northeast; they increased from 1 m/s to 5.4–5.6 m/s during the day. CTD cast 1 was made at 09:19, 53 min prior to fish release. It showed a 3-m-thick surface mixed layer with a temperature of 0.5°C and salinity of about 27.5 ppt. Temperature decreased and salinity increased in the pycnocline immediately below to values of about 0°C and 29.5 ppt, respectively, at 9 m depth.

Fish 1 was released at 10:12 at 70°00.5'N, 144°43.1'W, about 4.6 km north of Carter Creek. The fish displayed directed movement toward the northwest until contact was lost




Figure 41.—Track of Fish 1.




Figure 42.—Temperature vs. elapsed time, Fish 1.

at 13:31 (Fig. 41). Temperatures received from the transmitter during tracking ranged from 1.0 to 2.1°C; they oscillated through about one degree for about 1 h after tracking began and one-half degree or less thereafter (Fig. 42). Water depths increased monotonically from 8.5 m at the release point to 12.8 m at the time contact was lost.

**Fish 2.**—Tracking began on July 31. There was a 0.6-m swell and no seas during the duration of tracking. Winds were light (1–2.2 m/s) during the day, increased to 3.6–4.5 m/s in the evening, then weakened to 0–2.2 m/s after midnight. CTD casts 2, 3, and 4 were made at 09:07, 12:30, and 20:00, respectively. The thermohaline structures at the first and third CTD stations were similar; both were almost isohaline (about 31 ppt) from surface to bottom and had only about a 1°C temperature decrease between the 1°C surface waters and near-bottom waters. In contrast, CTD cast 3 displayed pronounced temperature and salinity gradients. Surface water was dilute (about 14 ppt) and cool (2.5°C). A pycnocline was evident between 6 and 8 m, below which temperatures were below 0°C and salinities above 23 ppt. The GPS was unavailable for two periods exceeding 30 min in the middle of the tracking session.

Fish 2 was released at 70°01.0'N, 144°48.2'W, about 4.2 km north of Marsh Creek. Tracking began at 10:50 (Fig. 43). The fish initially moved northward. At about 12:30 it began to veer in a clockwise direction; at about 20:00, it was moving almost due south. The transmitter signal was lost at 21:55, when 1273 was about 140 m off the beach in the eastern part of the study area. Shortly thereafter we began searching along the beach from the skiff and at 23:46 the fish was relocated. The fish moved northeastward within 50 m of the beach until tracking was terminated at 02:25 when it entered very shallow waters (less than 1 m depth) behind the barrier spits at Anderson Point.





Figure 43.—Track of Fish 2.



**Figure 44.**—Temperature vs. elapsed time, Fish 2. Note the very cold water after about 800 min, which presumably is due to upwelling, and the ensuing rapid temperature increase near the end of the record, which is presumed to reflect entry of the fish into the plume of the Sadlerochit River.

Fish 2 traversed waters having temperatures between  $-0.3$  and  $3.4^{\circ}\text{C}$  (Fig. 44). Beginning in waters slightly over  $0^{\circ}\text{C}$ , the fish moved into warmer waters and reached the warmest waters encountered offshore at about 13:30, when temperatures exceeded  $3^{\circ}\text{C}$ . From then until 01:00 the overall trend was for it to occupy colder temperatures, albeit with some oscillation about the trend line. The coldest waters traversed by the fish occurred along shore near Anderson Point, where temperatures below  $0^{\circ}\text{C}$  were transmitted. At about 02:00 a sharp increase in water temperature ensued; the warming trend continued until tracking terminated, when a temperature of  $3.3^{\circ}\text{C}$  was recorded. The fish was exposed to a temperature increase averaging  $0.14^{\circ}\text{C}/\text{min}$  while passing through this relatively strong gradient. Water depths during the tracking session began at 11 m, increased to 13 m, and then decreased to 3.5 m while tracking from 1273. They were less than 2 m while tracking from the skiff.

**Fish 3.**—Tracking of this fish commenced on August 1. At the time of release, seas were 0.6–0.9 m and winds were 3.6–4.5 m/s. Fog was present after 22:45. CTD cast 5 was taken at 17:46. The data showed little variation of salinity with depth and values near 31 ppt. The temperature profile indicated a 4-m-thick mixed layer of water slightly over  $2^{\circ}\text{C}$  capping a thermocline that extended to about 7 m depth, below which temperatures were isothermal and roughly  $-0.5^{\circ}\text{C}$ . The GPS was unavailable for three periods exceeding 30 min while following Fish 3.




Figure 45.—Track of Fish 3.




Figure 46.—Temperature vs. elapsed time, Fish 3.

This char was released at 17:53 at 70°00.9'N, 144°52.3'W, roughly 2.5 km north of the opening in the spit that forms the northern shore of Simpson Cove. It moved generally northwestward for about 2 h, then turned sharply back southeastward, maintaining that direction until about 21:55, when it veered southwestward (Fig. 45). Contact with the fish was lost at 22:45 in 1.4 m of water and foggy conditions west of Collinson Point. Temperatures transmitted from the fish during tracking ranged from 1.7 to 2.8°C (Fig. 46). The general trend observed was a slow temperature decrease with oscillations of less than one degree about the trend line. Water depths during tracking ranged from 10.7 to 1.4 m.

**Fish 4.**—We commenced this track on August 3. At 11:00 the wind was westerly at 3.6–4.5 m/s and seas were 0.6–0.9 m high. By 20:00 the wind had weakened to 1–1.8 m/s. Later, by 02:00, it was calm and fog was developing. A shift in wind direction occurred in early morning and by 07:00 winds were from the northeast at 1 m/s. Winds gradually increased to 1.8–2.2 m/s by 09:00, then freshened to 8.9–10.7 m/s by 13:10. Seven CTD casts were completed during this tracking period. Casts 16–20 were made while in contact with the fish. (As CTD casts 21 and 22 occurred while searching, they will not be considered here.) The times at which casts 16–20 were made are as follows: 11:01, 14:55, 20:55, 23:10, and 04:20. Surface water temperatures and salinities varied little at CTD stations 16–18, being about 4°C and 31 ppt, respectively. However, the mixed layer at station 18 was about 6 m thick, in contrast to the 4-m-thick layers at the other two stations. CTD station 16 had the most well-developed thermocline, reaching temperatures near 0°C below 9 m depth. CTD cast 19 was taken near shore in the eastern part of the study area. It was distinguished by essentially invariant salinity and weakly negative temperature profiles; salinity was about 30 ppt and temperature about 4–4.5°C. CTD cast 20 also was taken near shore, west of CTD cast 19. A two-layer structure was present. Surface temperature and salinity were about





Figure 47.—Track of Fish 4.



6°C and 29 ppt, respectively, in the 1-m-thick mixed layer. A 1-m-thick pycnocline separated slightly more saline and cool waters from those above. The GPS was unavailable for two periods exceeding 30 min.

The release of Fish 4 was delayed as an apparent weak signal from a previously tracked fish was detected upon arrival at the release location. After searching unsuccessfully for the source of the signal for about 1 hour, we released Fish 4 at 11:53 at 70°01.4'N, 144°45.7'W, some 4.9 km north of Marsh Creek. This was the largest and slowest char we tracked. The fish seemed to mill about for 6 h; its track during that time includes a loop about 0.5 km across (Fig. 47). At 18:00 the fish began directed easterly movement that brought it within 240 m of the shore just off a radio repeater station located in the eastern part of the study area at 23:35. The char then moved northeast close inshore until 02:03, whereupon it reversed direction and proceeded southwest until contact was lost at 09:05 in the vicinity of Carter Creek. 1273 was usually within 250 m of shore during the period from 23:35 to 09:05. The area between Marsh Creek and the repeater station was searched unsuccessfully until 15:55.

Water temperatures traversed by the fish ranged from 2.6 to 6.2°C. The temperature record from Fish 4 displayed a step-like structure (Fig. 48). From the time of release until about 22:00, temperatures remained between roughly 3 and 4°C. Between 22:00 and 23:30 a rapid increase of temperature to values exceeding 6°C occurred, after which temperatures displayed relatively little short term change but a gradual decreasing trend such that at the



**Figure 48.**—Temperature vs. elapsed time, Fish 4. Note the rapid increase of temperature occurring about 10 h after tracking commenced. Along the shore, relatively warm waters had replaced waters with temperatures below 0°C that were present 3 days earlier when Fish 2 traversed the area.





Figure 49.—Track of Fish 5.

time contact with the fish was lost they were 4.5–5°C. Between 22:00 and 23:20 water temperatures traversed by the fish increased by 3°C, or at an average rate of 0.04°C/min. It is notable that the highest temperatures occurred in the same inshore area of eastern Camden Bay that had water colder than 0°C when traversed by Fish 2 about 3 d earlier.

**Fish 5.**—Wind speeds were about 1 m/s at the time this tracking session began on August 10. GPS coverage was excellent throughout; no interval with GPS unavailability exceeding 20 min occurred. CTD cast 44 was taken at the release point at 12:14 and cast 45 at 14:30. The salinity profiles at the stations showed almost no vertical variation, both being slightly over 31 ppt throughout the water column. The temperature profiles at both locations were weakly negative; however, surface temperatures were about 4°C at CTD station 45, which was quite close to shore, in contrast to 2.5°C at station 44.

This char was released at 12:26 at 70°01.6'N, 144°47.6'W, about 5.3 km north of Marsh Creek. The fish displayed directed movement to the south and at 14:45, 1273 was 90 m off the shore at Marsh Creek in 3.3 m of water (Fig. 49). The transmitter signal was lost at 15:00. At 16:10 searching commenced from the skiff. The search was broken off at 18:00.

Fish 5 traversed a considerable temperature range in a short time (Fig. 50). For the first 90 min the fish was in waters having temperatures of 1.5–2.5°C. However, after about 14:00 the temperature record displays a sharp increase, reaching values over 6°C by 14:40. At about 14:28 an apparent 2°C discrepancy between the warmest temperature recorded by the CTD (about 3.9°C) and that at the fish's location (6°C) occurred. At the time, 1273 was about 900 m from the shoreline. We believe that the difference was real and that the fish probably was in a narrow, shallow inshore band of relatively warm, dilute water off Marsh



**Figure 50.**—Temperature vs. elapsed time, Fish 5.





Figure 51.—Track of Fish 6.

Creek. The average rate of temperature increase that Fish 5 was exposed to while moving into warmer water was  $0.08^{\circ}\text{C}/\text{min}$ . Water depths ranged from 10.7 m at the release point to 3 m when the fish was lost.

**Fish 6.**—This tracking session began on August 11. Winds were calm at time of release, but strengthened to about 2–3.5 m/s from the northeast after 16:25. Seas remained less than 0.5 m high. Radar fixes were used for position determination from 14:15 to 18:05 due to the unavailability of GPS positions. CTD casts 46 and 47 were made while in contact with the fish and cast 48 shortly after contact was lost; the cast times were 10:20, 12:25, and 19:50, respectively. The salinity structures at the three stations typified those seen earlier, being almost invariant from surface to bottom and roughly 31 ppt. The temperature profiles showed evidence of wind mixing in the form of nearly isothermal surface layers overlying thermoclines. The mixed layers were 2–4 m thick. Temperatures at the bottom were  $1\text{--}2^{\circ}\text{C}$ .

This char was released at 10:20 about 5.8 km north of Marsh Creek at  $70^{\circ}01.9'\text{N}$ ,  $144^{\circ}47.5'\text{W}$ . The fish moved southward—similar to Fish 5—and reached the vicinity of the shoreline at about 13:30, after which time it moved eastward. Fish 6 reached its easternmost position at about 17:00, then turned northward. At 18:05 the fish turned east-northeast, the direction it followed until contact was lost at 19:30 (Fig. 51). An unsuccessful search for Fish 6 was terminated at 23:30.

Fish 6 traversed waters having temperatures between  $2.7$  and  $5.5^{\circ}\text{C}$ . The temperature-at-time record was characterized by frequent, roughly  $1^{\circ}\text{C}$  oscillations about the trendline throughout most of the record (Fig. 52). Fish 6 proceeded from roughly  $3^{\circ}\text{C}$  water



**Figure 52.**—Temperature vs. elapsed time, Fish 6. Note the oscillatory structure of the temperature record about the general trend—most notably, between 140 and 300 min.




Figure 53.—Track of Fish 7.

at release to 5°C water by 12:00. Between 12:00 and about 17:00 the mean temperature was near 5°C. However, oscillations were notably abundant in this portion of the temperature record; their maxima and minima were 5.3–5.4°C and 4.6–4.7°C, respectively. The periods of the oscillations were 15–25 min. If a ground speed of 59 cm/s and a stationary temperature field are assumed, a 20-min period would represent a 710-m wavelength for an oscillation. A decreasing temperature trend occurred thereafter to the end of the record. Water depths during the tracking session ranging from 6.4 to 11 m.

**Fish 7.**—This tracking session began on August 12. Winds were about 1.3 m/s at 11:00, strengthening to 1.5 m/s by 13:00 and 3 m/s by 15:20. The GPS was unavailable for one period exceeding 30 min near the end of the session. CTD cast 49 was made at 10:54. A pronounced two-layer structure, manifested by a sharp thermocline lying between 5 and 7 m depth, was evident. The mixed layer was 5 m thick and had waters with temperatures of 3°C. Again, salinity was quite invariant, being about 31 ppt in the surface layer and 31.75 ppt below the pycnocline. Below the thermocline, water temperature was invariant and only slightly over 1°C.

Fish 7 was the only Arctic cisco tracked due to the unavailability of fish deemed sufficiently large to be unaffected by the transmitter. The release location was 70°01.8'N, 144°47.7'W, which is approximately 5.7 km off Marsh Creek. From the time tracking began at 10:59 until contact was lost in the channel off Collinson Point at 15:20, the cisco displayed very directed southwest movement (Fig. 53). An extensive search of the channel area employing both 1273 and the skiff was unsuccessful and was terminated at 18:40.

Temperatures transmitted from the cisco did not change much during tracking; they ranged from 2.0 to 3.4°C (Fig. 54). Some decrease of temperature was observed after about





Figure 54.—Temperature vs. elapsed time, Fish 7.

14:00. Water depths decreased monotonically from 11 m at the fish release point to 2.1 m when the signal was lost.

### Movement Patterns

Directedness indices varied widely among the char, ranging from 0.20 to 0.92. The compass bearings of Fishes 2–6 were between  $88^{\circ}\text{T}$  and  $236^{\circ}\text{T}$ , or generally shoreward (Fig. 55). Fish 1 moved offshore on a bearing of  $329^{\circ}\text{T}$ . Given the short duration of contact with this fish and similar initial offshore movements shown by Fishes 2 and 3, it is possible that Fish 1 also eventually turned shoreward. The mean direction of movement of all char was  $116^{\circ}\text{T}$ . While the direction of movement of char was dominantly shoreward, both the unweighted and weighted Rayleigh tests produced a conclusion of random net movement.



**Figure 55.**—Directedness indices and compass bearings of fish. Length of arrow indicates directedness of movement as a ratio of net to gross ground speed, direction of arrow indicates direction of movement based on starting and ending positions of track. Number at end of arrow indicates fish.

### Movement Rates

Ground speed statistics based on data averaged over 1-h bins are shown in Table 13. The average ground speeds of the char ranged from 48.8 cm/s to 74.7 cm/s. These values differ slightly from the gross ground speeds due to the omission of some data by truncation. Fluctuations in the ground speeds of individual fish reached 50 cm/s. The standard




**Table 13.**—Ground speed statistics of northern Dolly Varden char and Arctic cisco based on hourly averaged data.

Fish number	Number of samples	Speed			
		Average		SD	Range
		(cm/s)	(L/s)	(cm/s)	(cm/s)
Northern Dolly Varden char					
1	2	74.7	1.49	11.7	66.4–83.0
2	13	58.2	1.14	13.3	37.1–82.8
3	4	65.4	1.44	8.9	52.5–72.5
4	21	48.8	0.91	12.4	26.8–85.4
5	2	75.0	1.47	10.5	67.6–82.5
6	9	59.4	1.18	18.3	41.1–92.7
Arctic cisco					
7	4	48.6	1.25	5.8	41.3–55.2

deviations of the ground speeds also displayed considerable variability; the greater deviations occurred among fish tracked for relatively long periods. The Kruskal–Wallis test showed that the standardized ground speeds of Fishes 2, 4, and 6, which had the three longest records, were not significantly different. The Arctic cisco's average ground speed and standardized speed were 48.9 cm/s and 1.14 L/s, respectively. The standardized gross ground speeds of all fish vs. total time tracked are shown in Figure 56. They suggest a negative relationship of speed and time tracked—possibly a tag effect. This relationship was examined in more detail by reviewing demeaned ground speed records for each fish. No consistent pattern of decreasing speed with time was evident in the individual records. Furthermore, all three of the runs tests on the longest records (Tracks 2, 4, and 6) detected no trends. Examination of the hourly-averaged speed records also did not indicate any diurnal patterns. Differences in alongshore and offshore ground speeds of char were evident. Fish 2 averaged 56.8 cm/s offshore between 10:50 and 20:50 and 72.5 cm/s alongshore between 00:00 and 02:00. Fish 4 averaged 50.5 cm/s offshore between 12:05 and 22:05 and 48.4 cm/s alongshore between 00:00 and 09:00. However, neither difference was statistically significant according to the Mann–Whitney U test.

Fish 4 provided us a serendipitous opportunity to make a swimming speed estimate when it reversed direction of movement along shore. By assuming the fish's swimming speed and the alongshore current speed were invariant, swimming and current speeds were calculated from simultaneous equations employing ground speeds attained while moving in opposite directions; i.e.,  $V_f + V_c = V_{g1}$  and  $V_f - V_c = V_{g2}$ , where  $V_f$ ,  $V_c$ , and  $V_g$  represent swimming, current, and ground speeds, respectively, and  $V_{g1}$  is greater than  $V_{g2}$ . The 23:50–02:03 and 02:03–05:00 track segments selected were nearly coincident in space. The calculated values of Fish 4's northeasterly and southwesterly ground speeds were 54 and 42 cm/s, respectively, while its swimming speed was 48 cm/s, and the current speed was 6 cm/s to the northeast. The swimming speed estimate agrees closely with the fish's gross



**Figure 56.**—Standardized ground speed vs. elapsed time. Note the apparent decrease of speeds occurring as elapsed time increases. Numbers identify fish.

ground speed of 48.8 cm/s over the entire track. The inferred current is consistent with an observation logged at 23:55 that the tracking vessel was being set northeast parallel to the beach, as well as with the flow patterns deduced from oceanographic observations off Carter Creek (Hale 1991).

A relationship between ground speed and temperature was not evident from an examination of scatter plots for individual fish. As illustrated in Figure 57, the full range of speeds appeared possible at any given temperature. However, speed and temperature data from seven selected track segments (Fishes 2, 4, and 6) showed an apparent negative relationship (Fig. 58). Speeds decreased fairly steadily from about 1.4 L/s at approximately  $-0.2^{\circ}\text{C}$  to 0.7–0.8 L/s at  $4.4\text{--}5.8^{\circ}\text{C}$ . A Spearman rank correlation of temperature and ground speed data averaged over each segment indicated a significant negative correlation. The same data averaged over 35–70-min bins and used in a Kruskal–Wallis ANOVA, produced a conclusion of no significant difference in ground speeds at the various temperatures.

### Temperature Occupancy

A total of 493 observations of temperatures was obtained from the six char tracked. The mean water temperature occupied was  $3.6^{\circ}\text{C}$ , while the range was  $-0.5$  to  $6.5^{\circ}\text{C}$ . The frequency distribution of the temperatures was multimodal (Fig. 59). A total of 51 observations of temperature was obtained from the single Arctic cisco. The mean water temperature occupied was  $2.85^{\circ}\text{C}$  and the range was  $1.5\text{--}3.5^{\circ}\text{C}$ . The temperature frequency distribution for the Arctic cisco was unimodal (Fig. 60).




Figure 57.—Raw ground speed vs. temperature, Fish 2.






Figure 58.—Standardized ground speed vs. temperature. Selected temperature and associated speed data from Fishes 2, 4, and 6. Bars indicate standard errors about the means. Data segments used range from 35 to 255 min.




**Figure 59.**—Temperature occupancy, all char. The range of temperature occupied by the fish was similar to the observed range of temperature in the study area.



**Figure 60.**—Temperature occupancy, Arctic cisco.

## Temperature Preference

Temperatures available to fish and those used are shown in Figure 61. Six of the seven comparisons made at the beginning of tracking sessions showed the fish preferred water at the upper end of the available temperature range, while one indicated preference for the middle of the available range. However, these data are not considered to be representative because the fish may have been exhibiting escape responses and the time interval between the CTD cast and initial transmission from the fish often was considerable. Thus the following discussion focuses on comparisons made after tracking was under way.



**Figure 61.**—Temperature selection observations. Horizontal bars indicate range of temperature available as indicated by CTD. Time of CTD cast is shown to left of bar. Filled circle indicates transmitted temperature. Time of transmitted temperature is shown above circle if differing from CTD. Shaded bars indicate range of transmitted temperatures during 1-h period bracketing CTD cast, while x's indicate average temperature during the 1-h period. The fish that the observations come from is indicated at the far left of the figure.

Two underway temperature preference observations were made for Fish 2. In the first instance, a fairly wide range of temperatures was available ( $-0.3$  to  $2.5^{\circ}\text{C}$ ) and the fish used the high end of the range, which was in the uppermost 3 m of the water column. In the second, the fish occupied the middle of the  $0.5$ – $1.2^{\circ}\text{C}$  range and the uppermost 4 m.

Four underway temperature preference observations were made for Fish 4. During the first observation, the fish occupied intermediate temperature water from the available  $2.2$ – $4.0^{\circ}\text{C}$  range; it appears that it was in the thermocline at about 4 m depth. Similar selection was apparent during the second observation, when  $2.8$ – $3.6^{\circ}\text{C}$  waters were available; the fish appeared to be just below the mixed layer in  $3.2^{\circ}\text{C}$  water at 6 m depth. In the third and fourth observations, the fish selected the warmest waters available ( $4.7$  and  $5.9^{\circ}\text{C}$ , respectively). In the former instance it appeared to be in the uppermost 5 m, while in the latter it apparently used the 1-m-thick surface mixed layer. The rapid increase of water temperatures that occurred during the third observation confounds an unambiguous interpretation of selection, although the coincident-in-time temperature values agree quite well.

The underway observation for Fish 5 produced the greatest disparity between concurrent CTD and transmitter temperature readings and also the greatest transmitted temperature range during the hour-long period around the CTD cast—some  $4^{\circ}\text{C}$ . Transmitted temperature increased very rapidly prior to the CTD cast, then became relatively stable at  $5.9$ – $6.4^{\circ}\text{C}$ . Although nearby, this fish evidently was in waters having properties that differed dramatically from those at the tracking vessel, so a temperature preference inference is unwarranted.

Fish 6 had a wide range of temperatures available ( $1.7$ – $4.6^{\circ}\text{C}$ ) and selected the upper end of the range, which was represented by a 2-m-thick surface mixed layer.

## DISCUSSION

### Habitat Attributes

The unusually strong and persistent easterly winds during the summer of 1990 had a major influence on oceanographic conditions in the study area. Pack ice was driven far offshore, creating a broad band of open water along the coast. The resultant large wind fetch promoted active wind-driven currents and associated mixing processes in the area. As a consequence, temperature and salinity contrasts between surface and deeper waters were weakly developed as compared to those observed in Camden Bay in early August 1988 and much weaker than those in 1989. The difference was most evident with respect to salinity, where all but a few surface values were 30–32 ppt in 1990, as compared to about 10–12 ppt during the same period in 1988 and as little as 5 ppt in 1989. However, near-surface water temperatures in 1990 were also considerably lower than those during the preceding year. If the temperature and salinity criteria used to define water masses in 1988 (Thorsteinson et al. 1990) are applied to CTD stations occupied in 1990, only one of the latter had coastal water in the surface layer, one had transitional waters in the surface layer, and the remainder were purely marine throughout the water column.

## Methodology

The changeable spatial relationship between the tracking vessel and fish, navigation errors, escape responses and tag effects may have biased our ground speed estimates.

The screening procedure should have reduced the effects of positional errors on instantaneous speed estimates because positional errors would have remained relatively constant while distances used for the estimates increased. On the negative side, the screening may have produced some straightening of tracks and concomitantly decreased gross speed estimates. Upward biased speeds may have been produced because the tracking vessel meandered about the fish tracks and thereby covered more ground in a given time than the fish did. One avenue to address those biases involves comparisons of alongshore and offshore data. The former are relatively well constrained because of the close proximity of the vessel to the fish in shallow water, dictated by the decreased signal detection range, and the consistent spatial relationship of the vessel and fish, which is dictated by the shoreline. The best data available to us are from the track of Fish 4, from which we have an estimate of alongshore swimming speed that may be unbiased by currents. The fish's swimming speed was only 0.8 cm/s less than its offshore ground speed. The result is encouraging; however, it is based on very little data and should be considered tentative.

Screening of the temperature and ground speed data for tag effects or escape responses showed no evidence of such trends, so we do not believe that the speed estimates are biased due to those factors. The apparent negative relationship between gross ground speeds of individual char and elapsed time tracked seems to have arisen by chance.

The question arises as to how closely the gross ground speeds of the char and cisco approximate their actual swimming speeds. The good agreement of the swimming speed and gross ground speed estimates for Fish 4 suggests only small differences. Additional evidence is available from other telemetry studies in which swimming speeds of fish were calculated or actually measured. Quinn (1988) estimated sockeye salmon swimming speeds by vector subtraction of observed currents from fish tracks over ground. While there were considerable differences between average swimming and ground speeds for individual fish, agreement was good for the entire data set (66.76 and 64.7 cm/s, respectively;  $n = 25$  fish). Quinn surmised that the salmon did not respond strongly to currents and, thus, over extended temporal and spatial scales, currents became in effect random perturbations that were averaged out. Additional evidence supporting a close relation between swimming and ground speeds comes from a telemetry study conducted in deep ocean waters with no major currents. Carey and Scharold (1990) observed that the average swimming speed based on speedometer readings from a shark tracked for 19 h was comparable to its gross ground speed estimated from the tracking vessel's positions. The speeds were  $40 \pm 3$  cm/s and  $43 \pm 8$  cm/s, respectively.

When a large data set is available and the study area has weak and/or periodic currents, it is probably reasonable to equate gross ground speeds with swimming speeds. However, the Eulerian current data from the vicinity of Carter Creek have demonstrated that the current regime in Camden Bay is event-dominated, highly variable, and relatively little influenced by tides. Lagrangian observations likely would reveal comparable richness in the horizontal structure of the current field. Thus we conclude that until a more substantial data

base on swimming speeds of char and cisco is available, it is unwarranted to conclude that the gross ground speeds closely approximate the average swimming speeds of the fish.

## Northern Dolly Varden Char

### Movement Patterns

No consistent pattern of movement of char was shown by the Rayleigh tests or the directedness index. However, these results may be confounded by currents prevailing in 1990, about which no information is available at present. The movements of fishes can be strongly influenced by currents. Quinn and Leggett (1987) graphically illustrate this in figures depicting the track over ground and the actual swimming course of a telemetered Atlantic salmon (*Salmo salar*). The former shows a looping track indicative of milling, while the latter is quite straight and reflective of directed movement.

A recurring movement pattern of the char we tracked was the tendency to seek inshore waters. After being released offshore, four of six fish were observed to return to the vicinity of the shoreline. Relatively heavy use of shoreline habitats by char in the Beaufort Sea has been documented by Craig and Haldorson (1981) and others. In Cresswell Bay, Northwest Territories, no Arctic char were taken in water depths greater than 3.1 m and 60% were caught within 45 m of the shore, despite the fact that only 33% of the netting effort was within that area (Johnson 1980). Gillnet catches of char off Carter Creek in Camden Bay during the period July 30–September 12, 1988, do not demonstrate quite as strong a pattern. While the highest catch rate (0.58 fish/h) was in a net set 90 m from shore, the rates at the nets set 200 m and 1.6 km off shore were only slightly lower (0.44 fish/h) (Fruge et al. 1989).

The northern Dolly Varden char's association with shorelines is comparable to other anadromous salmonids that use coastal, rather than open ocean, habitats. In a tracking study of the southern form of Dolly Varden char conducted in the fjord habitats of southeastern Alaska, Armstrong and Reed (1971) found that char, once encountering a shoreline after being released in open water, did not return to open water, but remained within 6 m of the shoreline most of the time. Washington (1977) noted that in the fjord habitats of Puget Sound, cutthroat trout (*Oncorhynchus clarki*) feed in shallow estuaries, over gravel beaches and in or near "saltchucks" during high tide periods. Similarly, Jones (1976) observed that anadromous cutthroat trout in southeastern Alaska were reluctant to cross bodies of water 3–8 km wide and preferred to follow shorelines. However, off the open Oregon–Washington coast, cutthroat trout have been found to be relatively abundant from 9 to as much as 46 km offshore and much less so within 9 km of the coast. Significantly, these fish appear to be associated with the plume of the Columbia River, which persists far from the river's mouth (Trotter 1989; Percy et al. 1990).

Actively migrating adult pink salmon (*O. gorbuscha*) and sockeye salmon (*O. nerka*) returning to the Fraser River, British Columbia, did not follow shorelines as they passed through the San Juan Islands (Stasko et al. 1973, 1976). The anadromous form of the rainbow trout (*O. mykiss*), the steelhead, tracked in the Dean and Fisher Channels, British Columbia, displayed inconsistent use of shorelines during their return migration (Ruggerone et al. 1990).



## Movement Rates

The average and range of gross ground speed of the northern Dolly Varden char are very similar to those reported for steelhead trout and pink salmon and considerably higher than those for southern Dolly Varden, cutthroat trout, and Atlantic salmon (Table 14). They are somewhat lower than the speeds reported for sockeye salmon. However, given the considerable size differences between the species, standardized speeds provide a more meaningful comparison. On that basis, the northern Dolly Varden swam more rapidly than the Atlantic salmon, steelhead, or cutthroat trout and about as fast as sockeye salmon. It is also notable that laboratory studies (Beamish 1980) showed that, among Arctic char, lake trout (*Salvelinus namaycush*), and brook trout (*S. fontinalis*), the critical swimming speed of Arctic char is markedly higher than those of its congeners for fish of equal size.

Table 14.—Gross ground speeds of salmonids.

Species (sample size)	Total length (cm)	Group average speed		Individual average speed		Reference
		(cm/s)	(L/s)	(cm/s)	(L/s)	
Northern Dolly Varden (n = 6)	48.5–56.9	55.8 <sup>a</sup>	1.04	48.8–74.2	0.96–1.37	This study
Southern Dolly Varden (n = 1)	—	—	—	30.8	—	Armstrong and Reed (1971)
Cutthroat trout <sup>b</sup> (n = 14)	31–40	33.6 <sup>a</sup>	0.86	6.8–44.9	0.20–1.31	McCleave and LaBar (1972)
Steelhead, rainbow trout (n = 6)	72–89	57.4 <sup>a</sup>	0.74	38.9–69.4	0.54–0.96	Ruggerone et al. (1990)
Sockeye salmon <sup>b</sup> (n = 14)	51–71	61	—	35–101	—	Stasko et al. (1976)
Sockeye salmon (n = 13)	—	61	0.99	—	—	Blair and Quinn (1991)
Sockeye salmon (n = 25)	60.6–73.1	64.7 66.75 <sup>d</sup>	— 1.0 <sup>d</sup>	— 31.9–108.0 <sup>d</sup>	— 0.48–1.66 <sup>d</sup>	Quinn (1988)
Pink salmon <sup>c</sup> (n = 5)	58.5–68.5	62	—	40–78	—	Stasko et al. (1973)
Pink salmon <sup>e</sup> (n = 37)	41–60	48.3 <sup>f</sup>	—	—	—	Martin et al. (1990)
Atlantic salmon <sup>g</sup> (n = 1)	83	—	—	27.1	0.33	Døving et al. (1985)
Arctic cisco (n = 1)	42.9	—	—	48.9	1.14	This study

<sup>a</sup>Weighted average; summed track lengths/summed times.

<sup>b</sup>Entire track. Tracks 1–4, 6–9, 14, 17, 19, 21, 23, 25.

<sup>c</sup>Active fish only.

<sup>d</sup>Swimming speed; currents removed from ground speeds.

<sup>e</sup>Three control groups; sustained swimming toward stream.

<sup>f</sup>Grand mean of means weighted by numbers of fish in each of three control groups.

<sup>g</sup>Control fish "C."

The hourly-averaged ground speed statistics revealed considerable variation among individual char. The standard deviations of the fish's ground speeds always were considerably less than their average speeds, but increased as sample sizes became larger. Similarly, speed ranges were greater for fish tracked for comparatively long periods. While we have no basis to partition the variation between swimming speeds and currents, it is probable that the fish tracked for longer periods were exposed to more variability in current speed and direction and that this factor was primarily responsible for the larger standard deviations and ground speed ranges as tracking durations increased. It is notable that the speed ranges of the three fish tracked for 9- to 21-h periods all were about 50 cm/s, so that value may approximate the maximum variability of char ground speeds in the study area.

The only comparative data on char movement rates in arctic waters that we have are from tag recaptures. Dempson (1984) estimated a net ground speed of 16.4 km/d over 5 d by an Arctic char in the coastal waters of northern Labrador. Craig and Haldorson (1981) calculated an average net movement rate of 2.8 km/d from 30 char recaptures in the Beaufort Sea. However, they noted that recaptures of migrating fish in the fall indicate net movement rates of up to 78 km/d (90 cm/s), which exceeds the gross ground speed of any char we tracked. An Arctic char tagged in northern Norway was captured 97 d later near Murmansk in the USSR (Johnson 1980). This probably represents the greatest documented migratory movement by that species. The average rate of movement of the fish over 940 km was 9.7 km/d. The rates of net movement of char in Camden Bay ranged from 8.1 to 51.9 km/d, thus they fell between the extremes noted above.

The high gross ground speeds (average 48 km/d) of our char may reflect their motivational state. All of the char were large and probably 7–10 years old based on age-length data presented by Palmer and Dugan (1990). Assuming adult northern Dolly Varden char remain at sea for some 5–7 wk, as is the case with Arctic char in the Northwest Territories and northern Norway (Dempson and Kristofferson 1987), they would be expected to begin migration toward overwintering areas in August and, therefore, to be swimming in a relatively directed, rapid fashion. Palmer and Dugan (1990) observed that char larger than 350 mm fork length were absent from coastal waters of the Arctic National Wildlife Range after the end of August, although smaller fish remained until mid-September. A similar pattern of return to fresh water by Arctic char—mature fish returning before immature fish—was noted by Dempson and Kristofferson (1987) in northern Labrador. At-sea movement rates based on tag returns and presumably reflecting sustained migratory speeds are available for several Pacific salmonids. Ruggerone et al. (1990) noted that travel rates of 49 km/d were observed for steelhead on the high seas. Tag recaptures indicate that pink salmon achieve net ground speeds of 50–69 cm/s (43–60 km/d) over distances up to 2,000 km (Hartt 1966). A notable tag recovery from a 42-cm sockeye salmon demonstrated that the fish traversed a straight line distance of 2,200 km in 57 d (French et al. 1976); this is equivalent to a net ground speed of 44 cm/s, or better than 1 L/s, over the entire distance.

McCleave and Horral (1970) observed that cutthroat trout in Yellowstone Lake swam at significantly higher speeds when along shorelines than they did offshore, while Blair and Quinn (1991) found the opposite to be true of sockeye salmon in Lake Iliamna. This led us to investigate such potential differences in char. The available ground speed data are inconsistent; Fish 2 moved faster along shore than offshore, while Fish 4 did the opposite.

Runs tests disclosed no significant temporal trends in the ground speeds of the fish, and Mann-Whitney tests indicated no significant differences in onshore and offshore speeds. It is possible that our tests were confounded by biases in the data due to currents.

Immigrant salmon and steelhead tracked in coastal British Columbia and Washington showed significantly decreased net movement during darkness (cf. Ruggerone et al. 1990; Quinn 1988; Stasko et al. 1973, 1976). No such diurnal pattern was evident in the char we tracked or in sockeye salmon tracked in Lake Iliamna, Alaska (Blair and Quinn 1991). The lack of diurnal variation could be due to the absence of darkness during our study. Alternatively, it may reflect strong motivation by char to reach sites used for spawning (as Blair and Quinn surmised was the case for the Iliamna sockeye) or overwintering.


A negative relationship between water temperature and char ground speeds was apparent. Speed increases at cold temperatures could represent physiological or avoidance responses by the fish. A comparison of free-living fish with ones that are forced to perform may not be warranted, but it is notable that laboratory experiments on the effects of temperature on critical swimming speeds of Arctic char, lake trout, and brook trout (Beamish 1980) produced contrary results. All showed a dome-shaped relation between 5 and 25°C. Critical swimming performance of Arctic char peaked at 15°C. Given our equivocal statistical results and the lack of a larger suite of data spanning a broader temperature range, at this juncture the negative relationship should be considered more suggestive than substantive.

### **Temperature Occupation**

Northern Dolly Varden char used the entire range of water temperatures present in Camden Bay during our study. Perhaps this occurred because the fishes' options were limited by the paucity of brackish water habitat in the area at the time. Strong horizontal temperature gradients produced no discernible responses from the char we tracked. Fishes 2, 4, and 5 passed through relatively sharp, positive temperature gradients at speeds that differed little from their gross ground speeds. We had no comparable examples of the char encountering sharp negative gradients, thus their behavior when encountering those conditions is uncertain. It should be noted, however, that Fish 2 appeared to move readily through the cold, upwelled water in eastern Camden Bay (Fig. 62).

### **Temperature Preference**

Temperature preference is indicated by the disproportionate occupancy of certain temperatures within the total range available to a fish. The available range is ultimately determined by the fish's physiological limits. When the temperatures occupied by the char we tracked are compared to those available at the time in the water column, it appears that they selected the upper halves of the available ranges. However, we observed no avoidance of cold waters by Fish 2, which moved through water having temperatures below 0°C. At Cresswell Bay, Somerset Island in the Canadian high arctic, Johnson (1980) found Arctic char in -1.42 to -1.44°C water in late June. He concluded that the char tolerated the cold temperatures during foraging, but then returned to the 4-6°C waters at the mouth of the Union River to digest their food. Fyke net catch patterns in the vicinity of Prudhoe Bay



**Figure 62.**—Track of Fish 2 relative to horizontal temperature field as indicated by temperatures transmitted from the fish. Approximate shoreline contour sketched in. Note the non-avoidance of relatively cold water adjacent to the shoreline.

indicate that North Slope char prefer temperatures of 4–12°C during the early July to mid-August period (Houghton et al. 1990).

It also appears there are size-related differences in temperature selection or tolerance by char and other salmonids. Moulton et al. (1985) observed that in the Beaufort Sea larger char occurred more frequently in relatively cold, saline water than smaller ones. Large rainbow trout were observed to use cooler portions of a thermal discharge into Lake Michigan more frequently than smaller fish (Spigarelli and Thommes 1979).

### Swimming Depth

The use of the warmer waters by char that we observed may be less a reflection of temperature selection than one of depth selection. While we have no direct measurements of swimming depths, transmitted temperature data suggest that the fish used only the uppermost 8 m and most of the time were in the uppermost 4 m. The apparent preference for surface waters is consistent with other observations on char, as well as those for several other salmonids. The deepest occurrence of Arctic char we noted in the literature was of fish captured at 15 m depth in gillnets at Cresswell Bay, Somerset Island (Johnson 1980). The vertical distribution of char in gillnets in the Camden Bay area during 1988 and 1989 (Früge et al. 1989; Palmer and Dugan 1990) suggests heavy use of the uppermost 2.4 m, limited use of the 2.4–4.9-m stratum, and little or no use of the 4.9–7.3-m stratum.

Preferences for near-surface waters have also been observed in the southern form of the Dolly Varden char (Armstrong and Reed 1971), steelhead trout (Ruggerone et al. 1990), pink salmon (Martin et al. 1990), and sockeye salmon (Stasko et al. 1976; Quinn and terHart 1987). Concurrent depth telemetry and hydrographic observations showed that sockeye salmon preferred the immediate surface when in unstratified water, but selected greater depths (about 15–20 m) when vertical temperature and salinity gradients were present (Quinn and Leggett 1987).

The temperature records from several char contained oscillations suggestive of periodic vertical movements. Such behavior has previously been observed in Atlantic salmon (Døving et al. 1985), steelhead (Ruggerone et al. 1990), sockeye salmon (Quinn and terHart 1987), and pink salmon (Martin et al. 1990), but apparently not in char. The studies noted above were focused on immigrant adult fish. Tracked in deep fjords, Atlantic and sockeye salmon dived to as much as 80 m depth and steelhead to 30 m. The large amplitude dives of the steelhead averaged 28 min and those of the Atlantic salmon were of similar duration. In contrast, the pink salmon were studied in a shallow embayment, where they attained a depth of 4 m. The vertical amplitudes of the pink salmon's movements ranged from 0.5 to 2 m. While we have no direct observations of vertical excursions from the char we tracked, comparisons of temperature–depth profiles from CTD casts in relation to the observed temperature changes registered by the ultrasonic transmitters suggest that they would have been 2 m or less.

Many interpretations have been advanced for the repetitive vertical movements of fishes, including predatory strategies (Sciarrotta and Nelson 1977; Carey and Scharold 1990), energy conservation (Holland et al. 1990), behavioral thermoregulation (Holland et al. 1990; Carey and Scharold 1990), and orientation or homing (Døving et al. 1985; Martin et al. 1990). The last-mentioned one is compelling with respect to the char we observed. Field experiments by Døving et al. (1985) and Martin et al. (1990) suggest that migrating salmonids move vertically in the water column in order to detect odors and that much of the movement is concentrated in the pycnocline.

It is also possible to attribute the oscillatory features in the chars' temperature records to a purely physical phenomenon. While swimming at a constant depth, they may have traversed internal wave packets. Such features are commonly observed in coastal waters.

## Arctic Cisco

### Movement Patterns

The sole Arctic cisco we tracked displayed very directed movement toward the vicinity of its capture site, Simpson Cove. In common with other anadromous fish species in the Beaufort Sea, Arctic cisco appear most concentrated along shoreline habitats. However, they (and Arctic char) are more tolerant of marine conditions than the other species. Craig and Haldorson (1981) observed that, while cisco occurred offshore in Simpson Lagoon in relatively low densities, due to the greater expanse of available habitat, the numbers of fish offshore may have been equivalent to those close to shore. Finally, our townet sampling in 1990 (described elsewhere in this report) demonstrated the presence of significant numbers of migrating young-of-the-year Arctic cisco at distances up to several kilometers offshore.

## **Movement Rates**

The average ground speed of the Arctic cisco we tracked was not exceptionally great. However, its speed in lengths per second was only slightly less than that of the fastest char we tracked. In an evaluation of the swimming performance of nine Arctic cisco (mean FL of 30.4 cm), Jones et al. (1974) found a mean critical velocity of  $42.5 \pm 6.5$  cm/s, which equates to nearly 1.4 L/s. Craig and Haldorson (1981) noted several observations of mixed schools of Arctic char and Arctic cisco in the Beaufort Sea, which implies the latter species is capable of sustained swimming speeds comparable to those of char. A net movement rate of 24 km/d (28 cm/s) over 170 km was calculated from a tag recapture of a large Arctic cisco (Fechhelm et al. 1989). Another Arctic cisco, tagged at Prudhoe Bay and recaptured at Kaktovik, had a net movement rate of 12.5 km/d (14.5 cm/s) (Fruge et al. 1989).

## **Temperature Occupation and Preference**

The cisco we tracked traversed a relatively small range of temperatures, which was not representative of the entire temperature range available to the species in the Beaufort Sea. In Simpson Lagoon, Craig and Haldorson (1981) captured Arctic cisco in waters ranging from 0 to 13.5°C. Results of analyses of the large, multi-year catch and temperature/salinity databases from Prudhoe Bay and vicinity by Houghton et al. (1989) suggest that Arctic cisco prefer waters having temperatures of 4–12°C and 0–20 ppt salinity.

As we observed in the case of char, the occurrence of Arctic cisco in cold, saline water in Camden Bay during August 1990 may reflect tolerance of less-than-optimal environmental conditions by actively migrating fish more than it does habitat selection by feeding fish. Temperature preference investigations by Fechhelm and Gallaway (1984) showed that juvenile Arctic cisco initially acclimated at 5°C and 30 ppt—conditions similar to those prevailing in Camden Bay during our study—selected 12.8°C temperatures. In addition, they noted that the juvenile cisco acclimated to 5°C appeared lethargic, as well as being less agile and feeding less frequently than fish held at 10 and 15°C. Larger fish may have lower temperature preferences than the juveniles studied in the laboratory. Moulton et al. (1985) found that larger Arctic cisco occurred more frequently in colder, more saline water than smaller fish.

## **CONCLUSIONS**

The acoustic telemetry data from the northern Dolly Varden char and Arctic cisco tracked in Camden Bay provide some new information on the species' movements and habitat use in arctic marine waters, as well as an opportunity to compare the findings with those observed in other salmonid species. However, it is an incomplete picture. The timing of the observations makes it likely that the fish were returning to overwintering streams. Observations made earlier in the open-water season, when foraging was the dominant activity, might produce differing results. It is also possible that size-related differences in movement patterns and habitat use by the char occur. Refinements of tracking procedures have the potential to produce more precise ground speed, swimming speed, and course estimates. Finally, because brackish water habitat fluctuates so markedly in the Beaufort

Sea, habitat use studies will likely require more time and effort to reach fruition than would be the case in a more static setting.

Camden Bay is perhaps the most marine-like section of the Beaufort Sea coast in Alaska. The major regional sources of fresh water, the Mackenzie River to the east and the Colville to the west, are distant. Runoff from the relatively small local streams drops off sharply after spring breakup and would be low in August. The lack of barrier islands and comparatively deep waters near shore facilitate vertical and lateral mixing by strong winds, thereby quickly eroding what brackish water habitat is present inshore. This was especially evident in August 1990, when there was little warm, brackish water in Camden Bay compared to the preceding two years. One is led to the conclusion that the quality of the available habitat for anadromous fish—at least with respect to the thermohaline regime—was quite poor in 1990.

Fechhelm and Gallaway (1984) hypothesized that the thermal structure of the coastal environment in the Beaufort Sea causes an orientation bias favoring shoreward movement by juvenile Arctic cisco. The fishes' temperature preferences lie near or above the highest water temperatures, which are attained inshore. The hypothesis can be extended to all anadromous fish species in the Beaufort Sea, with the proviso that the strength of expression of the bias will vary in response to species-specific, age- and size-related physiological requirements and motivations.

Our temperature preference observations showed a preponderance of use by char and Arctic cisco of the warmer halves of the temperature ranges occurring in the water column. This may have been more reflective of depth selection than temperature selection because we observed no avoidance of cold water in the horizontal dimension. However, most of the fish tracked moved shoreward. Perhaps this behavior has survival value when available habitat is absent, discontinuous, or of marginal water quality. Association with shorelines may increase the fishes' opportunities to escape adverse conditions by maximizing their encounters with stream mouths. Conversely, when available habitat for char is expansive, the shoreline affinity may be relaxed. An example of this was noted earlier with respect to the cutthroat trout observed far off the Washington-Oregon coast in the Columbia River plume.

There is abundant evidence that with respect to use of arctic marine habitats, adult northern Dolly Varden char and Arctic cisco are the most resilient components of the anadromous fish fauna. Like the Arctic char, they are well adapted to disperse rapidly throughout the coastal waters of the Beaufort Sea and exploit a variety of habitats during the brief summer season (cf. Craig 1984). Our observations reinforce those conclusions.

## **RECOMMENDATIONS FOR FURTHER WORK**

### **Instrumentation**

Following the complete implementation of the Global Positioning System, extremely accurate, full-time position determination will be possible if the proposed signal degradation for civilian users is not implemented. The navigational error term in a fish track will be an order of magnitude smaller than the error due to the uncertainty of the position of the fish

vis-à-vis the tracking vessel (i.e., about 20 m vs. roughly 200 m). To capitalize on that possibility, positions should be logged to the nearest 0.01 minute if a test of navigation precision shows high quality fixes are being transmitted.

More accurate data on swimming speeds and courses of fish can be obtained if information is available on currents in the vicinity of fish being tracked. Currents appear to exert a substantial influence on fishes' ground speeds in the study area. Vector subtraction can be employed to derive swimming speeds from ground speeds and swimming courses from tracks over ground. In addition, we will be able to estimate bias resulting from movements of the tracking vessel about fish courses while tracking offshore when concurrent data on alongshore ground speeds and currents are available. An inexpensive, easily implemented technique to do this would be to launch current drogues in the path of a fish being tracked and to periodically determine the drogue's range and bearing by radar. The vessel heading and radar range and bearing data could be used later in conjunction with GPS positions to calculate drogue positions and current speeds. Drogues could be deployed, recovered and redeployed as needed by using a skiff in conjunction with the tracking vessel.

If a second ultrasonic receiver were available, it could be placed in the skiff when it became evident that a fish was entering waters too shallow for *I273* to enter. A second point of contact with the fish could be rapidly established. This should considerably increase our ability to maintain contact with fish in shallow water.

An intercomparison of each temperature/depth tag with the CTD should be conducted in order to be able later to adjust the former data to agree with the CTD "standard." This should be done through a range of temperatures and depths in case differences in readings between instruments are not constant. Intercomparisons could be quickly conducted by selecting a site having waters with a substantial temperature range and sufficient depth, temporarily activating each transmitter, attaching it near the sensor of the CTD, and then lowering the units through the water column. Transmitter readings would be logged at discrete depths for later comparison with the CTD data from those depths. This procedure should result in better quality information for habitat preference evaluations.

## **Data**

An additional data field should be added to the fish tracking log sheet—a signal quality code. A 0 could represent no audio or VU meter signal received; 1 could represent a VU meter deflection, but no audio signal; 2 could represent a VU meter deflection and audio signal, but the latter not clear; and 3 could represent a strong VU meter deflection and "loud and clear" audio signal. Signal strength information would improve the quality of the ground speed estimates by reducing the uncertainty of the fish position relative to the vessel. By selecting only Code 3 positions for ground speed calculations the analyst would be reasonably confident that the potential fish-boat position uncertainty was no more than about 200 m.

The telemetry data from northern Dolly Varden char that were obtained during 1990 provide information useful for assessment applications. However, additional fish should be tracked. Data obtained earlier in the open-water season would provide insights about movement patterns and habitat use during the chars' feeding period. The ranges of water



temperatures and salinities in the study area in 1990 were not representative of the total ranges occurring there or in the coastal Beaufort Sea. It would be desirable to obtain data when a strongly developed warm, brackish water band is present. Additional temperature data also would increase the database available for evaluation of temperature and swimming speed relationships. We have no data on the vertical movements of char other than those inferred from the temperature transmitter and CTD records. It would be desirable to equip at least two fish with pressure transmitters to obtain such information. Due to the shallow water in the study area the transmitters would need a depth resolution of 1 m or less to provide useful data. Further tracking of char would augment our meager data on movement patterns and habitat use.

Several Arctic ciscoes should be instrumented with temperature and depth transmitters and tracked. Camden Bay may not be the best site for an Arctic cisco tracking study. Large fish may be more readily available at some other location.

#### PART IV. REFERENCES CITED

- Armstrong, R. H., and J. E. Morrow. 1980. The dolly varden charr, *Salvelinus malma*. Pages 99–140 in E. K. Balon (ed.), Charrs: Salmonid fishes of the genus *Salvelinus*. Dr. W. Junk, The Hague, The Netherlands.
- Armstrong, R. H., and R. D. Reed. 1971. A study of dolly varden in Alaska. Alaska Dep. Fish and Game, Fed. Aid in Fish Restoration, Annual Prog. Rep., 1970–71, Project F-9-3-12. 11 pp.
- Bakkala, R. G., and G. B. Smith. 1978. Demersal fish resources of the eastern Bering Sea: Spring 1976. U.S. Dep. Commer., Northwest and Alaska Fisheries Center, Seattle, Washington. Processed Report, 233 pp.
- Barnes, P. W., and D. M. Rearic. 1986. Rates of sediment disruption by sea ice as determined from characteristics of dated ice gouges created since 1975 on the inner shelf of the Beaufort Sea, Alaska. U.S. Dep. Commer. and U.S. Dep. Inter., OCSEAP Final Rep. 52: 707–820.
- Beamish, F. W. H. 1980. Swimming performance and oxygen consumption of the charrs. Pages 739–748 in E. K. Balon (ed.), Charrs: Salmonid fishes of the genus *Salvelinus*. Dr. W. Junk, The Hague, The Netherlands.
- Behnke, R. J. 1980. A systematic review of the genus *Salvelinus*. Pages 441–480 in E. K. Balon (ed.), Charrs: Salmonid fishes of the genus *Salvelinus*. Dr. W. Junk, The Hague, The Netherlands.
- Bendock, T. 1979. Beaufort Sea estuarine fishery study. U.S. Dep. Commer. and U.S. Dep. Inter. (OCSEAP), Environ. Assess. Alaskan Continental Shelf, Final Rep. 14: 670–729.
- Berg, O. K., and M. Berg. 1989. Sea growth and time of migration of anadromous arctic char (*Salvelinus alpinus*) from the Vardnes River, in northern Norway. Can. J. Aquat. Sci. 46: 955–960.
- Blair, G. R., and T. P. Quinn. 1991. Homing and spawning site selection by sockeye salmon (*Oncorhynchus nerka*) in Iliamna Lake, Alaska. Can. J. Zool. 69: 176–181.
- Brawn, V. M. 1982. Behavior of Atlantic salmon (*Salmo salar*) during suspended migration in an estuary, Sheet Harbour, Nova Scotia, observed visually and by ultrasonic tracking. Can. J. Fish. Aquat. Sci. 39: 248–256.
- Brett, J. R. 1973. Energy expenditure of sockeye salmon, *Oncorhynchus nerka*, during sustained performance. J. Fish. Res. Board Can. 30: 1799–1809.
- Brill, R. W., K. N. Holland, and J. S. Ferguson. 1984. Use of ultrasonic telemetry to determine the short-term movements and residence times of tunas around fish

aggregating devices. Pages MRM1/1–MRM1/7 in Proceedings, Pacific Congress on Marine Technology, PACON 84, Honolulu, Hawaii, April 24–27, 1984.

- Bryan, J. D. 1990. Fine-scale biodiffusion model of the arctic cisco movement and distribution. Chapter 9 in *The 1988 Endicott Development Fish Monitoring Program: a monograph of project papers, 1988*. Draft final report by LGL Alaska Research Associates, Inc. for BP Exploration (Alaska), Inc. and North Slope Borough. Unpaginated.
- Carey, F. G., and B. H. Robison. 1981. Daily patterns in the activities of swordfish, *Xiphias gladius*, observed by acoustic telemetry. *Fish. Bull.* 79: 277–292.
- Carey, F. G., and J. V. Scharold. 1990. Movements of blue sharks (*Prionace glauca*) in depth and course. *Mar. Biol.* 106: 329–342.
- Clark, D. S., and J. M. Green. 1990. Activity and movement patterns of juvenile Atlantic cod, *Gadus morhua*, in Conception Bay, Newfoundland, as determined by sonic telemetry. *Can. J. Zool.* 68: 1434–1442.
- Cochran, W. G. 1963. *Sampling techniques*. Wiley & Sons, New York. 413 pp.
- Colonell, J. M., and B. J. Gallaway (eds.). 1990. An assessment of the marine environmental impacts of West Dock Causeway. Report for the Prudhoe Bay Unit Owners represented by Arco Alaska, Inc., prepared by LGL Alaska Research Associates, Inc., and Environmental Science and Engineering, Inc., Anchorage, AK. 132 pp. + appendixes.
- Colonell, J. M., and A. W. Niedoroda. 1990a. Nearshore oceanographic processes of potential importance to anadromous fishes. Pages 35–42 in R. M. Meyers and T. M. Johnson (eds.), *Fisheries oceanography—a comprehensive formulation of technical objectives for offshore application in the arctic*. Proceedings of a workshop held in Anchorage, AK, on April 5–6, 1988. U.S. Dep. Inter., Minerals Management Service, Anchorage, AK, and MBC Applied Environmental Sciences, Costa Mesa, CA.
- Colonell, J. M., and A. W. Niedoroda. 1990b. Appendix B. Coastal oceanography of the Alaska Beaufort Sea. Pages B-1–B-74 in J. M. Colonell and B. J. Gallaway (eds.), *An assessment of the marine environmental impacts of West Dock Causeway*. Report for the Prudhoe Bay Unit Owners represented by Arco Alaska, Inc., prepared by LGL Alaska Research Associates, Inc., and Environmental Science and Engineering, Inc., Anchorage, AK. 132 pp. + appendixes.
- Coutant, C. C., and D. S. Carroll. 1980. Temperatures occupied by ten ultrasonic-tagged striped bass in freshwater lakes. *Trans. Am. Fish. Soc.* 109: 195–202.
- Craig, P. C. 1984. Fish use of coastal waters of the Alaskan Beaufort Sea: a review. *Trans. Am. Fish. Soc.* 113: 265–282.

- Craig, P. C. 1989a. An introduction to anadromous fishes in the Alaskan Arctic. Biol. Pap. Univ. Alaska 24: 27-54.
- Craig, P. C. 1989b. Subsistence fisheries at coastal villages in the Alaskan arctic, 1970-1986. Biol. Pap. Univ. Alaska 24: 131-152.
- Craig, P. C., W. B. Griffiths, L. Haldorson, and H. McElderry. 1982. Ecological studies of arctic cod (*Boreogadus saida*) in Beaufort Sea coastal waters, Alaska. Can. J. Fish. Aquat. Sci. 39: 395-406.
- Craig, P. C., and L. S. Haldorson. 1981. Fish. Pages 384-678 in S. R. Johnson and W. J. Richardson (eds.), Beaufort Sea barrier island-lagoon ecological process studies. Final report, Simpson Lagoon. U.S. Dep. Commer. and U.S. Dep. Inter. (OCSEAP), Environ. Assess. Alaskan Continental Shelf, Final Rep. 7, Biol. Stud.
- Dempson, J. B. 1984. Identification of anadromous arctic char stocks in coastal areas of northern Labrador. Pages 143-162 in L. Johnson and B. Burns (eds.), Biology of the arctic char: Proceedings of the international symposium on arctic char, 1981, Winnipeg, Manitoba. University of Manitoba Press.
- Dempson, J. B., and A. H. Kristofferson. 1987. Spatial and temporal aspects of the ocean migration of anadromous arctic char. Am. Fish. Soc. Symposium 1: 340-357.
- Douglas, D. R., and L. A. Jahn. 1987. Radiotracking hybrid striped bass in Spring Lake, Illinois, to determine temperature and oxygen preferences. N. Am. J. Fish. Manage. 7: 531-534.
- Doving, K. B., H. Westerberg, and P. B. Johnsen. 1985. Role of olfaction in the behavioral and neuronal responses of Atlantic salmon, *Salmo salar*, to hydrographic stratification. Can. J. Fish. Aquat. Sci. 42: 1658-1667.
- Dunton, K. H., and S. V. Schonberg. 1982. The Canning River to Demarcation Bay: a preliminary survey of macrophyte communities. Cruise and summary report. Prepared for OCSEAP by Western Washington University, Bellingham, WA. 47 pp.
- English, K. K. In press. Effects of temperature, salinity and prey abundance on the growth of Arctic cisco and broad whitefish feeding on epibenthic prey *in-situ* enclosures. Am. Fish. Soc. Symposium 12.
- Everett, R. J., and R. L. Wilmot. 1990. Genetic stock structure of arctic char (*Salvelinus alpinus*) from drainages to the Beaufort Sea in Alaska and Canada. Report prepared for Minerals Management Service and National Oceanic and Atmospheric Administration by U.S. Fish & Wildlife Service, Anchorage, AK. 59 pp.
- Fechhelm, R. G., J. S. Baker, W. B. Griffiths, and D. R. Schmidt. 1989. Localized movement patterns of least cisco (*Coregonus sardinella*) and arctic cisco (*C. autumnalis*) in the vicinity of a solid-fill causeway. Biol. Pap. Univ. Alaska 24: 75-106.

- Fechhelm, R. G., J. D. Fechhelm, C. J. Herlugson, and D. K. Beaubien. In press. Bioassay tests of acute temperature and salinity shock to arctic cisco. Am. Fish. Soc. Symposium 12.
- Fechhelm, R. G., and D. B. Fissel. 1988. Wind-aided recruitment of arctic cisco (*Coregonus autumnalis*) into Alaskan waters. Can. J. Fish. Aquat. Sci. 45: 906-910.
- Fechhelm, R. G., and B. J. Gallaway. 1982. Temperature preference of juvenile arctic cisco (*Coregonus autumnalis*) from the Alaskan Beaufort Sea, in relation to salinity and temperature acclimation. U.S. Dep. Commer. and U.S. Dep. Inter., OCSEAP Final Rep. 23: 207-231.
- Fechhelm, R. G., and W. B. Griffiths. 1990. Effect of wind on the recruitment of Canadian arctic cisco (*Coregonus autumnalis*) into the central Alaskan Beaufort Sea. Can. J. Fish. Aquat. Sci. 47: 2164-2171.
- French, R., W. Bilton, M. Osako, and A. Hartt. 1976. Distribution and origin of sockeye salmon (*Oncorhynchus nerka*) in offshore waters of the North Pacific Ocean. Int. North Pac. Fish. Comm. Bull. 34. 113 pp.
- Fruge, D. J., D. W. Wiswar, L. J. Dugan, and D. E. Palmer. 1989. Fish population and hydrographic characteristics of Arctic National Wildlife Refuge coastal waters, summer 1988. Report by U.S. Fish and Wildl. Serv., Fairbanks, AK. 65 pp. + 6 append.
- Gallaway, B. J. 1990. Factors limiting the growth of arctic anadromous fish populations. Pages 57-62 in R. M. Meyer and T. M. Johnson (eds.), Fisheries oceanography—a comprehensive formulation of technical objectives for offshore application in the Arctic. U.S. Dep. Inter., Minerals Management Service, OCS Study MMS 88-0042. 112 pp. + appendixes.
- Gallaway, B. J., W. B. Griffiths, P. C. Craig, W. J. Gazey, and J. W. Helmericks. 1983. An assessment of the Colville River stock of arctic cisco—migrants from Canada? Biol. Pap. Univ. Alaska 21: 4-22.
- Grant, W. E. 1986. Systems analysis and simulation in wildlife and fisheries sciences. John Wiley & Sons, New York.
- Green, R. H. 1979. Sampling design and statistical methods for environmental biologists. John Wiley & Sons, New York.
- Grossman, G. D., J. F. Dowd, and M. Crawford. 1990. Assemblage stability in stream fishes: a review. Environmental Management 14(5): 661-671.
- Hair, J. F. Jr., R. E. Anderson, and R. L. Tatham. 1987. Multivariate data analysis (second edition). Macmillan Publishing Company, New York.

- Hale, D. A. 1990. A description of the physical characteristics of nearshore and lagoonal waters in the eastern Beaufort Sea. Report prepared for U.S. Fish and Wildl. Serv. by Ocean Assessments Division, NOAA, Anchorage, AK. 72 pp. + appendixes.
- Hale, D. A. 1991. A description of the physical characteristics of nearshore and lagoonal waters in the eastern Beaufort Sea, 1989. Final report prepared for U.S. Fish and Wildl. Serv. by Ocean Assessments Division, NOAA, Anchorage, AK.
- Hale, D. A., M. J. Hameedi, L. E. Hachmeister, and W. J. Stringer. 1989. Effects of the West Dock Causeway on nearshore oceanographic processes in the vicinity of Prudhoe Bay, Alaska. Special Report prepared for the U.S. Environmental Protection Agency, Region 10, by the U.S. Department of Commerce, NOAA, Ocean Assessments Division, Alaska Office, Anchorage, AK. 50 pp.
- Hartt, A. C. 1966. Migrations of salmon in the North Pacific Ocean and Bering Sea as determined by seining and tagging, 1959–1960. Int. N. Pac. Fish. Comm. Bull. 19: 141 pp.
- Hobbs, N. T., and T. A. Hanley. 1990. Habitat evaluation: Do use/availability data reflect carrying capacity? J. Wildl. Manage. 54(4): 515–522.
- Holland, K. N., R. W. Brill, and R. K. C. Chang. 1990. Horizontal and vertical movements of yellowfin and bigeye tuna associated with fish aggregating devices. U.S. Natl. Mar. Fish. Serv. Fish. Bull. 88(3): 493–507.
- Holland, K. N., R. W. Brill, S. Ferguson, R. Chang, and R. Yost. 1985. A small vessel technique for tracking pelagic fish. U.S. Natl. Mar. Fish. Serv. Mar. Fish. Rev. 47(4): 26–32.
- Hopkins, D. M., and R. W. Hartz. 1978. Coastal morphology, coastal erosion, and barrier islands of the Beaufort Sea, Alaska. USGS Open-File Rep. 78–1063. 54 pp.
- Houghton, J. P., and C. J. Whitmus. 1988. Shallow neritic fish of the central Beaufort Sea. Draft final report to Standard Alaska Production Company from Dames and Moore. Contract No. 172–0002–20. 26 pp. + appendixes.
- Houghton, J. P., C. J. Whitmus, and A. W. Maki. 1990. Habitat relationships of Beaufort Sea anadromous fish: integration of oceanographic information and fish-catch data. Pages 73–88 in R. M. Meyer and T. M. Johnson (eds.), Fisheries oceanography—a comprehensive formulation of technical objectives for offshore application in the Arctic. U.S. Dep. Inter., Minerals Management Service, OCS Study MMS 88–0042. 112 pp. + appendixes.
- Johnson, L. 1980. The arctic charr, *Salvelinus alpinus*. Pages 15–98 in E. G. Balon (ed.), Charrs: salmonid fishes of the genus *Salvelinus*. Dr. W. Junk, The Hague, The Netherlands.

- Jones, D. E. 1976. Life history of sea-run cutthroat trout in southeastern Alaska: Anadromous fish studies, annual performance report, 1975-76. Alaska Dep. Fish and Game Proj. AFS-42, 117 (AFS-42-4B): Section M, 29-53.
- Jones, D. K., J. W. Kiseniuk, and O. S. Bamford. 1974. Evaluation of the swimming performance of several fish species from the Mackenzie River. J. Fish. Res. Board Can. 31: 1641-1647.
- Kelso, J. R. M. 1977. Density, distribution, and movement of Nipigon Bay fishes in relation to a pulp and paper mill effluent. J. Fish. Res. Board Can. 34: 879-885.
- Laurs, R. M., H. S. H. Yuen, and J. H. Johnson. 1977. Small-scale movements of albacore, *Thunnus alalunga*, in relation to ocean features as indicated by ultrasonic tracking and oceanographic sampling. U.S. Natl. Mar. Fish. Serv. Fish. Bull. 75(2): 347-355.
- Lyons, S. 1990. Water resource inventory and assessment, Arctic National Wildlife Refuge: 1989 stream discharge gaging data. Fisheries Management Services, Water Resources Branch, U.S. Fish & Wildlife Service, Anchorage, AK. 9 pp. + appendixes.
- Martin, D. J., C. J. Whitmus, and L. A. Brocklehurst. 1990. Effects of petroleum contaminated waterways on migratory behavior of adult pink salmon. U.S. Dep. Commer. and U.S. Dep. Inter., OCSEAP Final Rep. 66: 281-529.
- Matthews, K. R. 1990. A telemetric study of the home ranges and homing routes of copper and quillback rockfishes on shallow rocky reefs. Can. J. Zool. 68: 2243-2250.
- Maughan, O. E. 1990. Habitat. Pages 85-96 in MBC Applied Environmental Sciences (eds.), A synthesis of environmental information on causeways in the nearshore Beaufort Sea. U.S. Dep. Inter., Minerals Management Service, OCS Study MMS 89-0038. 117 pp. + appendixes.
- MBC Applied Environmental Sciences. 1990. A synthesis of environmental information on causeways in the nearshore Beaufort Sea, workshop proceedings. U.S. Dep. Inter., Minerals Management Service, OCS Study MMS 89-0038. 117 pp. + appendixes.
- McCleave, J. D., and R. M. Horrall. 1970. Ultrasonic tracking of homing cutthroat trout *Salmo clarki* in Yellowstone Lake. J. Fish. Res. Board Can. 27: 715-730.
- McCleave, J. D., and G. W. LaBar. 1972. Further ultrasonic tracking and tagging studies of homing cutthroat trout (*Salmo clarki*) in Yellowstone Lake. Trans. Am. Fish. Soc. 101: 44-54.
- McDowall, R. M. 1987. The occurrence and distribution of diadromy among fishes. Am. Fish. Soc. Symposium 1: 1-13.
- McPhail, J. D. 1961. A systematic study of the *Salvelinus alpinus* complex in North America. J. Fish. Res. Board Can. 5: 131-137.


- Mellas, E. J., and J. M. Haynes. 1985. Swimming performance and behavior of rainbow trout (*Salmo gairdneri*) and white perch (*Morone americana*): effects of attaching telemetry transmitters. *Can. J. Fish. Aquat. Sci.* 42: 488-493.
- Menke, W. W. 1973. Response surface methodology applications to biological data from field measurements. *Ecology* 54(4): 920-923.
- Morrow, J. E. 1980. Analysis of the dolly varden char, *Salvelinus malma*, of northwestern North America and northeastern Siberia. Pages 323-338 in E. K. Balon (ed.), *Charrs: salmonid fishes of the genus Salvelinus*. Dr. W. Junk, The Hague, The Netherlands.
- Moulton, L. L. 1989. Recruitment of arctic cisco (*Coregonus autumnalis*) to the Colville Delta, Alaska, in 1985. *Biol. Pap. Univ. Alaska* 24: 107-111.
- Moulton, L. L., M. H. Fawcett, and T. A. Carpenter. 1985. Lisburne Development environmental studies: 1984. Fish, Final Report, Chap. 5, Vol. 4. Prepared for ARCO Alaska, Inc., by Woodward-Clyde Consultants, Anchorage, AK.
- Moulton, L. L., and L. J. Field. 1989. Assessment of the Colville River fall fishery, 1985-87. Pages 47-53 in R. M. Meyer and T. M. Johnson (eds.), *Fisheries oceanography—a comprehensive formulation of technical objectives for offshore application in the arctic*. Proceedings of a workshop held in Anchorage, AK, April 5-6, 1988. U.S. Dep. Inter., Minerals Management Service, Anchorage, AK, and MBC Environmental Sciences, Costa Mesa, CA.
- Moulton, L. L., and K. E. Tarbox. 1987. Analysis of arctic cod movements in the Beaufort Sea nearshore region, 1978-79. *Arctic* 40(1): 43-49.
- Neill, W. H., R. J. Fechhelm, B. J. Gallaway, J. D. Bryan, and S. W. Anderson. 1983. Modeling movement and distribution of arctic cisco (*Coregonus autumnalis*) relative to temperature-salinity regimes of the Beaufort Sea near the Waterflood Causeway, Prudhoe Bay, Alaska. *Biol. Pap. Univ. Alaska* 21: 39-61.
- Orth, D. J. 1971. Dictionary of Alaska place names. USGS Prof. Paper 567. U.S. Government Printing Office, Washington, DC. 1084 pp. + 12 maps.
- Pahlke, K. A. 1985. Preliminary studies of capelin (*Mallotus villosus*) in Alaskan waters. Alaska Dep. Fish and Game Informational Leaflet 250. 65 pp.
- Palmer, D. E., and L. J. Dugan. 1990. Fish population characteristics of Arctic National Wildlife Refuge coastal waters, summer 1989. Progress Report. Fishery Assistance Office, U.S. Fish & Wildlife Service, Fairbanks, AK. 83 pp.
- Paragamian, V. L. 1989. Seasonal habitat use by walleye in a warmwater river system, as determined by radiotelemetry. *N. Am. J. Fish. Manage.* 9: 392-401.











- Pearcy, W. G., R. D. Brodeur, and J. P. Fisher. 1990. Distribution and biology of juvenile cutthroat trout *Oncorhynchus clarki clarki* and steelhead *O. mykiss* in coastal waters off Oregon and Washington. U.S. Natl. Mar. Fish. Serv. Fish. Bull. 88(4): 697-711.
- Quinn, T. P. 1988. Estimated swimming speeds of migrating adult sockeye salmon. Can. J. Zool. 66: 2160-2163.
- Quinn, T. P., and R. D. Brodeur. 1991. Intra-specific variations in the movement patterns of marine animals. Am. Zool. 31: 231-241.
- Quinn, T. P., and W. C. Leggett. 1987. Perspectives on marine migrations of diadromous fishes. Am. Fish. Soc. Symposium 1: 377-388.
- Quinn, T. P., and B. A. terHart. 1987. Movements of adult sockeye salmon (*Oncorhynchus nerka*) in British Columbia coastal waters in relation to temperature and salinity stratification: ultrasonic telemetry results. Pages 61-77 in H. D. Smith, L. Margolis, and C. C. Wood (ed.), Sockeye salmon (*Oncorhynchus nerka*) population and future management. Can. Spec. Publ. Fish. Aquat. Sci. 96.
- Reynolds, W. W. 1977. Temperature as a proximate factor in orientation behavior. J. Fish. Res. Board Can. 34: 734-739.
- Ricker, W. E. 1975. Computation and interpretation of biological statistics of fish populations. Bull. Fish. Res. Board Can. 191. 382 pp.
- Rose, G. A., and W. C. Leggett. 1989. Interactive effects of the geophysically-forced sea temperatures and prey abundance on mesoscale coastal distributions of a marine predator, Atlantic cod (*Gadus morhua*). Can. J. Fish. Aquat. Sci. 46: 1904-1913.
- Ross, B. D. 1988. Causeways in the Alaskan Beaufort Sea. U.S. Environmental Protection Agency, Region 10, Anchorage, AK. EPA 910/0-88-218. 25 pp.
- Ruggerone, G. T., T. P. Quinn, I. A. McGregor, and T. D. Wilkinson. 1990. Horizontal and vertical movements of adult steelhead trout, *Oncorhynchus mykiss*, in the Dean and Fisher Channels, British Columbia. Can. J. Fish. Aquat. Sci. 47: 1963-1969.
- Schimdt, D. R., W. B. Griffiths, D. K. Beaubien, and C. J. Herlugson. In press. Movement of young-of-the-year arctic cisco across the Beaufort Sea coast. Am. Fish. Soc. Symposium 12.
- Sciarrotta, T. C., and D. R. Nelson. 1977. Diel behavior of the blue shark, *Prionace glauca*, near Santa Catalina Island, California. U.S. Natl. Mar. Fish. Serv. Fish. Bull. 75: 519-528.
- Spigarelli, S. A., and M. M. Thommes. 1979. Temperature selection and estimated thermal acclimation by rainbow trout (*Salmo gairdneri*) in a thermal plume. J. Fish. Res. Board Can. 36: 366-376.

- Standard Alaska Production Company. 1989. Assessing the impact of causeways on Beaufort Sea fish. A white paper prepared by Standard Alaska Production Company, Environmental and Regulatory Affairs, P.O. Box 196612, Anchorage, AK. 6 pp. + 6 Issue Papers + Appendixes.
- Stasko, A. B., R. M. Horral, and A. D. Hasler. 1976. Coastal movements of adult Fraser River sockeye salmon (*Oncorhynchus nerka*) observed by ultrasonic tracking. *Trans. Am. Fish. Soc.* 105: 64-71.
- Stasko, A. B., R. M. Horral, A. D. Hasler and D. Stasko. 1973. Coastal movements of mature Fraser River pink salmon (*Oncorhynchus gorbuscha*) as revealed by ultrasonic tracking. *J. Fish. Res. Board Can.* 30: 1309-1316.
- Stasko, A. B., and D. G. Pincock. 1977. Review of underwater biotelemetry, with emphasis on ultrasonic techniques. *J. Fish. Res. Board Can.* 34: 1261-1285.
- Stasko, A. B., and S. M. Polar. 1973. Hydrophone and bow mount for tracking fish by ultrasonic telemetry. *J. Fish. Res. Board Can.* 30: 119-121.
- Taylor, S. G., J. C. Landingham, D. G. Mortensen, and A. C. Wertheimer. 1987. Pink salmon early life history in Auke Bay: growth, diet, and survival. Component 8 in Annual report summary of 1986 APPRISE research program. Volume I. Technical Report. UAJ SFS-8702. 422 pp.
- Thorsteinson, L. K., L. E. Jarvela, and D. A. Hale. 1990. Arctic fish habitat investigations: nearshore studies in the Alaskan Beaufort Sea, summer 1988. U.S. Dep. Commer. and U.S. Dep. Inter., OCSEAP Final Rep. 71: 349-485.
- Trotter, P. C. 1989. Coastal cutthroat trout: A life history compendium. *Trans. Am. Fish. Soc.* 118: 463-473.
- Truett, J. C. (ed.). 1983. Environmental characterization and biological use of lagoons in the eastern Beaufort Sea. U.S. Dep. Commer. and U.S. Dep. Inter., OCSEAP Final Rep. 24: 129-579.
- Tyus, H. M., B. D. Burdick, and C. W. McAda. 1984. Use of radiotelemetry for obtaining habitat preference data on Colorado squawfish. *N. Am. J. Fish. Manage.* 4: 177-180.
- Washington, P. 1977. The sea-run cutthroat trout resource and sport fishery. U.S. Natl. Mar. Fish. Serv. *Mar. Fish. Rev.* 39(12): 20-22.
- Williams, I. V., and J. R. Brett. 1987. Critical swimming speed of Fraser and Thompson River pink salmon (*Oncorhynchus gorbuscha*). *Can. J. Fish. Aquat. Sci.* 44: 348-356.
- Winter, J. D. 1983. Underwater biotelemetry. Pages 371-395 in L. A. Nielson and D. L. Johnson (eds.), *Fisheries techniques*. American Fisheries Society, Bethesda, MD.


- Winter, J. D., and M. J. Ross. 1982. Methods in analyzing fish habitat utilization from telemetry data. Pages 273-279 in N. B. Armantrout (ed.), Proceedings of the symposium on the acquisition and utilization of aquatic habitat inventory information. Western Division, American Fisheries Society, Bethesda, MD.
- Winters, J. F., P. K. Weber, A. L. DeCicco, and N. Shishido. 1988. An annotated bibliography of selected references of fishes of the North Slope of Alaska, with emphasis on research conducted in National Petroleum Reserve-Alaska. Prepared for North Slope Borough by Alaska Dep. Fish and Game, Fairbanks, AK. 159 pp.
- Wise, J. L., and L. D. Leslie. 1988. Section I: Selected topics in marine and coastal meteorology. Climatic atlas, Vol. III: Chukchi-Beaufort Sea. National Climatic Data Center, Asheville, NC, and Arctic Environmental Information and Data Center, Anchorage, AK.
- Zar, J. H. 1984. Biostatistical analysis. Prentice-Hall, Englewood Cliffs, NJ.











Appendix A—CTD Station Data, 1990











Appendix A—CTD Station Data, 1990 (continued)



Appendix A—CTD Station Data, 1990 (continued)











Appendix A—CTD Station Data, 1990 (continued)












Appendix A—CTD Station Data, 1990 (continued)





Appendix A—CTD Station Data, 1990 (continued)





**Appendix B—Townet Catch Per Unit Effort (CPUE), 1990<sup>a</sup>**

Date	Station/cast	Arctic cisco	Arctic char	Arctic cod	Kelp snailfish	Capelin	Ninespine stickleback	Sculpins	Arctic flounder	Eelblennies
8/02	CC01/007						1	5		
8/02	CC01/008	3								
8/02	CC02/011	water haul								
8/02	CP10/013	water haul								
8/02	CP10/014	water haul								
8/02	CP10/015	water haul								
8/05	CP01/024	1		8		4				
8/05	CP01/025			9		3				
8/05	CP02/027			397						
8/05	CP02/028	water haul								
8/09	EC14/030				4					
8/09	EC14/031				1					
8/09	EC13/033				1					
8/09	EC13/034	water haul								
8/09	EC12/036	2		411						
8/09	EC12/037			64						
8/09	EC11/039	1		1						
8/09	EC11/040	1								
8/09	EC10/042	1				1	1	1		
8/09	EC10/043	6			1		18	1		
8/21	WD24/051	1		22	27					
8/21	WD24/052	56		174	9					
8/21	WD24/053			1	6			2		
8/21	WD23/055	33		3,418						
8/21	WD23/056	20		533		1				
8/22	WD21/058	154		1	4		2			
8/22	WD21/059	72			5					
8/22	WD22/061	14		1		2				

Appendix B—Townet CPUE, 1990<sup>a</sup> (continued)

Date	Station/cast	Arctic cisco	Arctic char	Arctic cod	Kelp snailfish	Capelin	Ninespine stickleback	Sculpins	Arctic flounder	Eelblennies
8/22	WD22/062	20		1	2	10				
8/22	WD25/064	4			50					
8/22	WD25/065				45					
8/23	WD21/067	6		74	2	151	2	9		
8/23	WD21/068	3		481		522	4	9	2	3
8/25	CP10/070	3		8		2	4			
8/25	CP10/071	8		2			22			
8/26	CP10/073	6		1	3			2		
8/26	CP10/074	5		5	1		5	5		
8/27	MS01/081	54		560		5				
8/27	MS01/082	28	1	210			1			
8/31	AI01/084	7		1,190	140	60				
8/31	AI01/085			950	151	75		1		
9/02	AI01/087			251						
9/02	AI01/088	2		125						
9/02	AI02/090			75						
9/02	AI02/091			27						
9/02	AI03/093			64						
9/02	AI03/094			42						
9/02	AI04/096			13						
9/02	AI04/097			2						
9/02	CP15/099			8	5					
9/02	CP15/100			5	6					
9/02	CP14/102			132	6					
9/02	CP14/103			46	3					
9/02	CP13/105			700	7					
9/02	CP13/106			991						
9/02	CP12/108			907						

**Appendix B—Townet CPUE, 1990<sup>a</sup> (continued)**

Date	Station/cast	Arctic cisco	Arctic char	Arctic cod	Kelp snailfish	Capelin	Ninespine stickleback	Sculpins	Arctic flounder	Eelblennies
9/02	CP12/109			904						
9/02	CP11/111			677						
9/02	CP11/112			572				1		
9/03	CP10/114	6		98			17			4
9/03	CP10/115	12		77	6		9			
9/03	BP04/117				1					
9/03	BP04/118			1						
9/03	BP03/120	water haul								
9/03	BP03/121	water haul								
9/03	BP02/123				14					
9/03	BP02/124				7					
9/03	BP01/126			1	3					
9/03	BP01/127			11	2					
9/03	MS01/129			40	5					
9/03	MS01/130			82	1		3			
9/04	EC10/132	terminated								
9/05	WD22/134	1		2		1				
9/05	WD22/135	1					1			
9/05	WD23/137			13						
9/05	WD21/139	15		417	3	46			2	1
9/05	WD21/140	14		140		23	7		1	

<sup>a</sup> Absence of data denotes zero catch.

Appendix C—Fish Densities (numbers/1,000 m<sup>3</sup>), 1990

Date	Station/cast	Arctic cisco	Arctic char	Arctic cod	Kelp snailfish	Capelin	Ninespine stickleback	Sculpins	Arctic flounder	Eelblennies	Unknown	Total
8/02	CC01/007							0.17	0.85			1.02
8/02	CC01/008	0.55										0.55
8/02	CC02/011											0.00
8/02	CP10/013											0.00
8/02	CP10/014											0.00
8/02	CP10/015											0.00
8/05	CP01/024	0.31		2.45		1.23						3.99
8/05	CP01/025			2.72		0.91					1.21	4.84
8/05	CP02/027			170.22								170.22
8/05	CP02/028											0.00
8/09	EC14/030				0.88							0.88
8/09	EC14/031				0.22							0.22
8/09	EC13/033				0.19							0.19
8/09	EC13/034											0.00
8/09	EC12/036	0.44		91.17								91.61
8/09	EC12/037			13.68								13.68
8/09	EC11/039	0.26		0.26								0.52
8/09	EC11/040	0.26										0.26
8/09	EC10/042	0.26				0.26	0.26	0.26				1.04
8/09	EC10/043	1.54			0.26		4.63	0.26				6.69
8/21	WD24/051	0.20		4.30	5.27							9.77
8/21	WD24/052	20.32		63.12	3.27							86.71
8/21	WD24/053			0.44	2.64			0.88				3.96
8/21	WD23/055	7.14		740.03								747.17
8/21	WD23/056	4.30		114.59		0.21						119.11
8/22	WD21/058	32.26		0.21	0.84		0.42					33.73
8/22	WD21/059	14.97			1.04							16.00
8/22	WD22/061	4.97		0.36		0.71						6.04

**Appendix C—Fish Densities, 1990 (continued)**

Date	Station/cast	Arctic cisco	Arctic char	Arctic cod	Kelp snailfish	Capelin	Ninespine stickleback	Sculpins	Arctic flounder	Eelblennies	Unknown	Total
8/22	WD22/062	7.23		0.36	0.72	3.62						11.93
8/22	WD25/064	1.45			18.12							19.56
8/22	WD25/065				18.67							18.67
8/23	WD21/067	1.17		14.47	0.39	29.52	0.39	0.39				47.70
8/23	WD21/068	1.09		173.96		188.79	1.45	3.26	0.72	1.09		370.36
8/25	CP10/070	0.66		1.77		0.44	0.88					3.75
8/25	CP10/071	1.77		0.44			4.86					7.07
8/26	CP10/073	1.55		0.26	0.77			0.52				3.09
8/26	CP10/074	1.29		1.29	0.26		1.29	1.29				5.41
8/27	MS01/081	10.55		109.40		0.98						120.93
8/27	MS01/082	5.45	0.20	40.85			0.20					46.69
8/31	AI01/084	4.50		765.35	90.04	38.59						898.48
8/31	AI01/085			396.64	63.05	31.31		0.42				491.42
9/02	AI01/087			64.61								64.61
9/02	AI01/088	0.52		32.18								32.69
9/02	AI02/090			19.23								19.23
9/02	AI02/091			6.92								6.92
9/02	AI03/093			16.48								16.48
9/02	AI03/094			10.81								10.81
9/02	AI04/096			3.35								3.35
9/02	AI04/097			0.52								0.52
9/02	CP15/099			2.57	1.61							4.18
9/02	CP15/100			2.14	2.57							4.72
9/02	CP14/102			29.25	1.33							30.58
9/02	CP14/103			11.85	0.77							12.62
9/02	CP13/105			181.51	1.82							183.32
9/02	CP13/106			303.26								303.26
9/02	CP12/108			234.81								234.81



**Appendix C—Fish Densities, 1990 (continued)**

Date	Station/cast	Arctic cisco	Arctic char	Arctic cod	Kelp snailfish	Capelin	Ninespine stickleback	Sculpins	Arctic flounder	Eelblennies	Unknown	Total
9/02	CP12/109			200.29								200.29
9/02	CP11/111			150.17								150.17
9/02	CP11/112			183.52				0.32				183.34
9/03	CP10/114	1.55		25.23			4.38			1.03		32.18
9/03	CP10/115	3.86		19.82	1.55		2.32			1.03		27.80
9/03	BP04/117				0.43							0.43
9/03	BP04/118			0.43								0.43
9/03	BP03/120											0.00
9/03	BP03/121											0.00
9/03	BP02/123				4.30							4.30
9/03	BP02/124				2.12							2.12
9/03	BP01/126			0.44	1.32							1.76
9/03	BP01/127				3.98	0.72						4.70
9/03	MS01/129			6.76	0.85							7.61
9/03	MS01/130			35.94	0.44		1.31					37.69
9/04	EC10/132											—
9/05	WD22/134	0.19		0.39		0.19						0.77
9/05	WD22/135	0.19					0.19					0.39
9/05	WD23/137			2.70								2.70
9/05	WD21/139	3.86		107.34	0.77	11.84			0.52	0.26		124.59
9/05	WD21/140	6.01		60.07		9.87	3.00		0.42			79.37

## Appendix D—Catch Summary, 1990

Date	Station	Location	Species	Catch Summary			
				$\Sigma$	$\bar{X}$	SD	CV
8/02	CC01	Carter Creek	Arctic cisco	3	1.50	2.12	1.14
			Ninespine stickleback	1	0.50	0.71	1.42
			Sculpins	5	2.50	3.54	1.42
			Total	9	4.50	2.12	0.47
8/02	CP10	Simpson Cove	Three (3) water hauls				
8/05	CP01	Collinson Point	Arctic cisco	1	0.50	0.71	1.42
			Arctic cod	17	8.50	0.71	0.08
			Capelin	7	3.50	0.71	0.20
			Unknown	4	—	—	—
			Total	29	14.50	2.12	0.15
8/05	CP02	Collinson Point	Arctic cod	397	198.50	280.72	1.41
8/09	EC14	Endicott Causeway	Kelp snailfish	5	2.50	2.12	0.85
8/09	EC13	Endicott Causeway	Kelp snailfish	1	0.50	0.71	1.42
8/09	EC12	Endicott Causeway	Arctic cisco	2	1.00	1.41	1.41
			Arctic cod	475	237.50	245.37	1.03
8/09	EC11	Endicott Causeway	Arctic cisco	2	1.00	0.00	0.00
8/09	EC10	Endicott Causeway	Arctic cisco	7	4.00	3.54	1.01
			Kelp snailfish	1	0.50	0.71	1.42
			Capelin	1	5.00	0.71	1.42
			Ninespine stickleback	19	9.50	12.02	0.13
			Sculpins	2	1.00	0.00	0.00
			Total	30	15.00	15.56	1.04
8/21	WD24	West Dock Causeway	Arctic cisco	56	28.00	39.60	1.40
			Arctic cod	175	87.50	122.33	1.40
			Kelp snailfish	15	7.50	2.12	0.28
			Sculpins	2	1.00	1.41	1.41
			Total	248	124.00	162.63	1.31
8/21	WD23	West Dock Causeway	Arctic cisco	53	26.50	9.19	0.35
			Arctic cod	3,951	1,975.50	2,040.00	1.03
			Capelin	1	0.50	0.71	1.42
			Total	4,005	2,002.50	2,048.48	1.02
8/21	WD21	West Dock Causeway Comment: Northward tows	Arctic cisco	226	113.00	57.98	0.51
			Arctic cod	1	0.50	0.71	1.42
			Kelp snailfish	9	4.50	0.71	0.16
			Ninespine stickleback	2	1.00	1.41	1.41
			Total	238	119.00	59.40	0.50
8/22	WD22	West Dock Causeway	Arctic cisco	34	17.00	4.24	0.25
			Arctic cod	2	1.00	0.00	0.00
			Kelp snailfish	2	1.00	1.41	1.41
			Capelin	12	60.00	5.66	0.94
			Total	50	25.00	11.31	0.45

**Appendix D—Catch Summary (continued)**

Date	Station	Location	Species	Catch Summary			
				$\Sigma$	$\bar{X}$	SD	CV
8/22	WD25	West Dock Causeway	Arctic cisco	4	2.00	2.83	1.41
			Kelp snailfish	95	47.50	3.53	0.07
			Total	99	49.50	6.36	0.13
8/23	WD21	West Dock Causeway	Arctic cisco	9	4.50	2.12	0.05
			Arctic cod	555	277.50	287.79	1.04
			Kelp snailfish	2	1.00	1.41	1.41
			Capelin	673	336.50	262.34	0.78
			Ninespine stickleback	6	3.00	1.41	0.47
			Sculpins	18	9.00	—	—
			Arctic flounder	2	1.00	1.41	1.41
			Eelblennies	3	1.50	2.12	1.41
Total	1,268	634.00	548.71	0.87			
8/25	CP10	Collinson Point	Arctic cisco	11	5.50	3.54	0.64
			Arctic cod	10	5.00	4.24	0.85
			Capelin	2	1.00	1.41	1.41
			Ninespine stickleback	26	13.00	12.73	0.98
			Total	49	24.50	10.61	0.43
8/26	CP10	Simpson Cove	Arctic cisco	11	5.50	0.71	0.13
			Arctic cod	6	3.00	2.83	0.94
			Kelp snailfish	4	2.00	1.41	0.71
			Capelin	5	2.50	3.54	1.41
			Sculpins	7	3.50	2.12	0.61
			Total	33	16.50	6.36	0.39
8/27	MS01	Mary Sachs Entrance	Arctic cisco	82	41.00	18.38	0.45
			Arctic char	1	0.50	0.71	1.41
			Arctic cod	770	385.00	247.49	0.64
			Capelin	5	2.50	3.54	1.41
			Ninespine stickleback	1	0.50	0.71	1.41
			Total	859	429.50	267.00	0.62
8/31	AI01	Arey Island	Arctic cisco	7	3.50	4.95	1.41
			Arctic cod	2,140	1,070.00	169.71	0.16
			Kelp snailfish	291	145.50	7.78	0.05
			Capelin	135	67.50	10.61	0.16
			Sculpin	1	0.50	0.71	1.41
			Total	2,574	1,287.00	155.56	0.12
9/02	AI01	Arey Island	Arctic cisco	2	1.00	1.41	0.71
			Arctic cod	376	188.00	89.10	0.47
			Total	378	189.00	87.68	0.46
9/02	AI02	Arey Island	Arctic cod	102	51.00	33.94	0.67
9/02	AI03	Arey Island	Arctic cod	106	53.00	15.56	0.29
9/02	AI04	Arey Island	Arctic cod	15	7.50	7.78	1.04
9/02	CP15	Collinson Point	Arctic cod	13	6.50	2.12	0.33
			Kelp snailfish	11	5.50	0.71	0.13
			Total	24	12.00	1.41	0.12

**Appendix D—Catch Summary (continued)**

Date	Station	Location	Species	Catch Summary			
				$\Sigma$	$\bar{X}$	SD	CV
9/02	CP14	Collinson Point	Arctic cod	178	89.00	60.81	0.68
			Kelp snailfish	9	4.50	2.12	0.47
			Total	187	93.50	62.93	0.67
9/02	CP13	Collinson Point	Arctic cod	1,691	845.50	205.77	0.24
			Kelp snailfish	7	3.50	4.95	1.41
			Total	1,698	849.00	200.82	0.23
9/02	CP12	Collinson Point	Arctic cod	1,811	905.50	2.12	0.02
9/02	CP11	Collinson Point	Arctic cod	1,249	624.50	74.25	0.12
9/03	CP10	Collinson Point	Arctic cisco	18	9.00	4.24	0.47
			Arctic cod	175	87.50	14.85	0.17
			Kelp snailfish	6	3.00	4.24	1.41
			Ninespine stickleback	26	13.00	5.66	0.44
			Eelblennies	8	4.00	0.00	0.00
			Total	233	116.50	12.02	0.10
9/03	BP04	Brownlow Point	Arctic cod	1	0.50	0.71	1.42
			Kelp snailfish	1	0.05	0.71	1.42
9/03	BP03	Brownlow Point	Water hauls				
9/03	BP02	Brownlow Point	Arctic cod	21	10.50	4.95	0.47
9/03	BP01	Brownlow Point	Arctic cod	12	6.00	7.07	1.18
			Kelp snailfish	5	2.50	0.71	0.28
			Total	17	8.50	6.36	0.75
9/03	MS01	Mary Sachs Entrance	Arctic cod	122	61.00	29.70	0.49
			Kelp snailfish	6	3.00	2.83	0.94
			Ninespine stickleback	3	1.50	2.12	1.41
			Total	131	65.50	28.99	0.44
9/04	EC10	Endicott Causeway		0			
		No replicate					
9/05	WD22	West Dock Causeway	Arctic cisco	2	1.00	0.00	0.00
			Arctic cod	2	1.00	1.41	1.41
			Capelin	1	0.50	0.71	1.41
			Ninespine stickleback	1	0.50	0.71	1.41
			Total	6	3.00	1.41	0.47
9/05	WD23	West Dock Causeway	Arctic cod	13	—	—	—
		No replicate					
9/05	WD21	West Dock Causeway	Arctic cisco	29	14.50	0.71	0.05
			Arctic cod	547	273.50	188.80	0.69
			Kelp snailfish	3	1.50	2.12	1.41
			Capelin	23	11.50	16.26	1.41
			Ninespine stickleback	7	3.50	4.95	1.41
			Arctic flounder	3	1.50	0.71	0.47
			Eelblenny	1	0.50	0.71	1.41
			Total	613	306.50	171.83	0.56

Appendix E—Townet and CTD Station Locations, 1990

<u>Date</u>	<u>Station</u>	<u>FISH CAST</u>	<u>OCEAN CAST</u>	<u>LAT</u>	<u>LONG</u>
8/02	CC01	007 008	006	69 58.1	144 41.9
8/02	CC02	010 011	009	70 00.1	144 41.8
8/02	CP10	013 014 015	012	69 59.0	144 53.8
8/05	CP11	024 025	023	70 00.3	144 55.2
8/05	CP02	027 028	026	70 02.3	155 55.3
8/09	EC14	030 031	029	70 18.9	147 35.2
8/09	EC13	033 034	032	70 17.4	147 35.4
8/09	EC12	036 037	035	70 15.8	147 35.6
8/09	EC11	039 040	038	70 14.2	147 35.4
8/10	EC10	042 043	041	70 13.5	147 35.5
8/21	WD24	051 052 053	050	70 27.8	148 33.0
8/21	WD23	055 056	054	70 26.7	148 32.9
8/22	WD21	058 059	057	70 24.4	148 32.9
8/22	WD22	061 062	060	70 25.7	148 32.8
8/22	WD25	064 065	063	70 28.8	148 33.0
8/23	WD21	067 068	066	70 24.4	148 32.9
8/25	CP10	070 071	069	69 58.8	144 55.2
8/26	CP10	073 074	072	69 59.0	144 55.1
8/27	MS01	081 082	080	70 11.5	146 15.9
8/31	AI01	084 085	083	70 08.4	143 54.3

**Appendix E—Townet and CTD Station Locations, 1990 (continued)**

<u>Date</u>	<u>Station</u>	<u>FISH CAST</u>	<u>OCEAN CAST</u>	<u>LAT</u>	<u>LONG</u>
9/02	AI01	087	086	70 08.4	143 54.1
		088			
9/02	AI02	090	089	70 09.1	143 53.4
		091			
9/02	AI03	093	092	70 10.9	143 53.9
		094			
9/02	AI04	096	095	70 12.3	143 54.0
		097			
9/02	CP15	099	098	70 06.0	144 55.2
		100			
9/02	CP14	102	101	70 04.3	144 55.1
		103			
9/02	CP13	105	104	70 02.8	144 55.2
		106			
9/02	CP12	108	107	70 01.2	144 55.2
		109			
9/03	CP11	111	110	70 00.5	144 55.1
		112			
9/03	CP10	114	113	69 58.8	144 55.2
		115			
9/03	BP04	117	116	70 14.6	145 50.0
		118			
9/03	BP03	120	119	70 13.7	145 51.0
		121			
9/03	BP02	123	122	70 11.6	145 51.2
		124			
9/03	BP01	126	125	70 10.7	145 51.2
		127			
9/03	MS01	129	128	70 11.3	146 16.9
		130			
9/04	EC10		131	70 13.6	147 35.8
9/05	WD22	134	133	70 25.6	148 33.0
		135			
9/05	WD23	137	136	70 26.6	148 33.7
9/05	WD21	139	138	70 28.8	148 32.9
		140			

## Appendix F—Ground Speed Data, 1990

Fish 1      Start Date 7/30/90

	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELASPED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
1	1039	.0	.0	.0	.0	.0
2	1046	676.4	420.0	161.0	676.4	161.0
3	1054	315.1	480.0	65.6	991.4	110.2
4	1105	445.8	660.0	67.5	1437.2	92.1
5	1115	489.5	600.0	81.6	1926.7	89.2
6	1125	532.9	600.0	88.8	2459.6	89.1
7	1135	368.3	600.0	61.4	2827.9	84.2
8	1206	1254.3	1860.0	67.4	4082.3	78.2
9	1215	418.2	540.0	77.4	4500.4	78.1
10	1225	377.3	600.0	62.9	4877.8	76.7
11	1236	377.3	660.0	57.2	5255.1	74.9
12	1245	377.4	540.0	69.9	5632.6	74.5
13	1253	185.6	480.0	38.7	5818.2	72.4
14	1305	377.3	720.0	52.4	6195.5	70.7
15	1315	372.0	600.0	62.0	6567.5	70.2
16	1325	372.0	600.0	62.0	6939.4	69.7

	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELAPSED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
1	1050	.0	.0	.0	.0	.0
2	1100	417.4	600.0	69.6	417.4	69.6
3	1110	186.4	600.0	31.1	603.8	50.3
4	1120	489.5	600.0	81.6	1093.3	60.7
5	1130	393.2	600.0	65.5	1486.6	61.9
6	1140	571.9	600.0	95.3	2058.5	68.6
7	1200	641.9	1200.0	53.5	2700.4	64.3
8	1210	377.3	600.0	62.9	3077.8	64.1
9	1220	377.4	600.0	62.9	3455.2	64.0
10	1230	186.4	600.0	31.1	3641.6	60.7
11	1240	377.3	600.0	62.9	4019.0	60.9
12	1300	737.1	1200.0	61.4	4756.0	61.0
13	1310	224.9	600.0	37.5	4980.9	59.3
14	1320	127.6	600.0	21.3	5108.5	56.8
15	1330	424.4	600.0	70.7	5532.9	57.6
16	1340	127.6	600.0	21.3	5660.5	55.5
17	1345	.0	300.0	.0	5660.5	53.9
18	1350	63.5	300.0	21.2	5724.0	53.0
19	1355	314.6	300.0	104.9	6038.6	54.4
20	1400	.0	300.0	.0	6038.6	53.0
21	1405	64.1	300.0	21.4	6102.7	52.2
22	1411	185.6	360.0	51.5	6288.3	52.1
23	1415	190.5	240.0	79.4	6478.8	52.7
24	1420	64.1	300.0	21.4	6542.9	51.9
25	1425	196.9	300.0	65.6	6739.9	52.2
26	1430	190.5	300.0	63.5	6930.4	52.5
27	1435	191.1	300.0	63.7	7121.5	52.8
28	1440	196.1	300.0	65.4	7317.7	53.0
29	1445	127.6	300.0	42.5	7445.3	52.8
30	1450	63.5	300.0	21.2	7508.8	52.1
31	1455	63.5	300.0	21.2	7572.3	51.5
32	1500	127.0	300.0	42.3	7699.3	51.3
33	1515	724.8	900.0	80.5	8424.1	53.0
34	1520	63.5	300.0	21.2	8487.6	52.4
35	1525	63.5	300.0	21.2	8551.2	51.8
36	1530	63.5	300.0	21.2	8614.7	51.3
37	1535	127.6	300.0	42.5	8742.3	51.1
38	1540	224.9	300.0	75.0	8967.2	51.5
39	1545	318.2	300.0	106.1	9285.3	52.5
40	1550	.0	300.0	.0	9285.3	51.6
41	1556	318.7	360.0	88.5	9604.0	52.3
42	1605	196.9	540.0	36.5	9801.0	51.9
43	1610	.0	300.0	.0	9801.0	51.0
44	1615	372.0	300.0	124.0	10172.9	52.2
45	1620	267.0	300.0	89.0	10439.9	52.7
46	1625	196.9	300.0	65.6	10636.9	52.9
47	1630	266.0	300.0	88.7	10902.8	53.4
48	1635	186.4	300.0	62.1	11089.2	53.6
49	1642	191.1	420.0	45.5	11280.4	53.4
50	1645	225.9	180.0	125.5	11506.3	54.0
51	1650	196.9	300.0	65.6	11703.2	54.2



	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELAPSED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
52	1655	63.5	300.0	21.2	11766.7	53.7
53	1700	196.1	300.0	65.4	11962.8	53.9
54	1705	185.6	300.0	61.9	12148.4	54.0
55	1710	225.2	300.0	75.1	12373.6	54.3
56	1715	196.1	300.0	65.4	12569.7	54.4
57	1720	.0	300.0	.0	12569.7	53.7
58	1725	196.9	300.0	65.6	12766.7	53.9
59	1750	683.0	1500.0	45.5	13449.6	53.4
60	1755	185.6	300.0	61.9	13635.2	53.5
61	1800	127.6	300.0	42.5	13762.8	53.3
62	1805	445.2	300.0	148.4	14208.0	54.4
63	1820	809.1	900.0	89.9	15017.1	55.6
64	1825	186.4	300.0	62.1	15203.5	55.7
65	1830	185.6	300.0	61.9	15389.1	55.8
66	1907	1126.4	2220.0	50.7	16515.5	55.4
67	1910	.0	180.0	.0	16515.5	55.1
68	1940	393.1	1800.0	21.8	16908.6	53.2
69	1945	418.2	300.0	139.4	17326.7	54.0
70	2000	1487.9	900.0	165.3	18814.6	57.0
71	2010	224.9	600.0	37.5	19039.5	56.7
72	2015	196.9	300.0	65.6	19236.4	56.7
73	2020	63.5	300.0	21.2	19299.9	56.4
74	2030	372.0	600.0	62.0	19671.9	56.5
75	2035	224.9	300.0	75.0	19896.8	56.7
76	2040	372.0	300.0	124.0	20268.7	57.3
77	2050	196.9	600.0	32.8	20465.7	56.8
78	2055	.0	300.0	.0	20465.7	56.4
79	2100	196.1	300.0	65.4	20661.8	56.5
80	2105	196.1	300.0	65.4	20857.9	56.5
81	2110	63.5	300.0	21.2	20921.4	56.2
82	2115	186.4	300.0	62.1	21107.8	56.3
83	2120	.0	300.0	.0	21107.8	55.8
84	2130	377.3	600.0	62.9	21485.2	56.0
85	2135	196.1	300.0	65.4	21681.3	56.0
86	2155	482.8	1200.0	40.2	22164.2	55.5
87	0000	2758.0	7500.0	36.8	24922.2	52.6
88	0005	225.6	300.0	75.2	25147.7	52.7
89	0010	372.0	300.0	124.0	25519.7	53.2
90	0015	418.2	300.0	139.4	25937.9	53.7
91	0020	185.6	300.0	61.9	26123.4	53.8
92	0025	450.8	300.0	150.3	26574.2	54.3
93	0035	196.1	600.0	32.7	26770.4	54.1
94	0040	190.5	300.0	63.5	26960.9	54.1
95	0050	450.8	600.0	75.1	27411.7	54.4
96	0055	266.4	300.0	88.8	27678.1	54.6
97	0100	225.6	300.0	75.2	27903.6	54.7
98	0110	318.2	600.0	53.0	28221.8	54.7
99	0115	127.0	300.0	42.3	28348.8	54.6
100	0120	185.6	300.0	61.9	28534.4	54.7
101	0125	196.9	300.0	65.6	28731.3	54.7
102	0130	191.1	300.0	63.7	28922.4	54.8

	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELASPED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
103	0135	196.1	300.0	65.4	29118.6	54.8
104	0140	.0	300.0	.0	29118.6	54.5
105	0145	196.9	300.0	65.6	29315.5	54.6
106	0150	185.6	300.0	61.9	29501.0	54.6
107	0155	266.4	300.0	88.8	29767.4	54.8
108	0200	377.3	300.0	125.8	30144.8	55.2
109	0205	63.5	300.0	21.2	30208.3	55.0
110	0210	376.5	300.0	125.5	30584.8	55.4
111	0215	186.4	300.0	62.1	30771.2	55.4
112	0220	225.9	300.0	75.3	30997.1	55.6
113	0225	.0	300.0	.0	30997.1	55.3

	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELAPSED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
1	1753	.0	.0	.0	.0	.0
2	1755	186.4	120.0	155.3	186.4	155.3
3	1805	63.5	600.0	10.6	249.9	34.7
4	1810	417.2	300.0	139.1	667.1	65.4
5	1815	.0	300.0	.0	667.1	50.5
6	1820	225.9	300.0	75.3	893.0	55.1
7	1825	368.3	300.0	122.8	1261.3	65.7
8	1830	315.6	300.0	105.2	1576.9	71.0
9	1905	1448.3	2100.0	69.0	3025.2	70.0
10	1940	1375.5	2100.0	65.5	4400.7	68.5
11	1945	418.2	300.0	139.4	4818.9	71.7
12	1950	63.5	300.0	21.2	4882.4	69.6
13	1955	225.2	300.0	75.1	5107.6	69.8
14	2005	948.9	600.0	158.2	6056.6	76.5
15	2010	254.6	300.0	84.9	6311.2	76.8
16	2015	225.6	300.0	75.2	6536.8	76.7
17	2020	127.6	300.0	42.5	6664.4	75.6
18	2025	254.6	300.0	84.9	6919.0	75.9
19	2030	127.0	300.0	42.3	7046.1	74.8
20	2035	127.6	300.0	42.5	7173.7	73.8
21	2040	127.0	300.0	42.3	7300.7	72.9
22	2045	127.6	300.0	42.5	7428.3	72.0
23	2050	224.9	300.0	75.0	7653.2	72.1
24	2055	63.5	300.0	21.2	7716.7	70.7
25	2100	127.0	300.0	42.3	7843.7	69.9
26	2105	191.1	300.0	63.7	8034.9	69.7
27	2110	63.5	300.0	21.2	8098.4	68.5
28	2115	254.6	300.0	84.9	8353.0	68.9
29	2155	1254.3	2400.0	52.3	9607.3	66.2
30	2200	315.1	300.0	105.0	9922.4	67.0
31	2205	196.9	300.0	65.6	10119.3	66.9
32	2210	254.6	300.0	84.9	10374.0	67.3
33	2220	318.2	600.0	53.0	10692.1	66.7
34	2225	196.1	300.0	65.4	10888.3	66.7
35	2230	267.0	300.0	89.0	11155.2	67.1
36	2235	63.5	300.0	21.2	11218.8	66.3
37	2240	185.6	300.0	61.9	11404.3	66.2
38	2245	63.5	300.0	21.2	11467.8	65.5

	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELAPSED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
1	1205	.0	.0	.0	.0	.0
2	1210	224.9	300.0	75.0	224.9	75.0
3	1215	63.5	300.0	21.2	288.4	48.1
4	1220	196.9	300.0	65.6	485.3	53.9
5	1225	63.5	300.0	21.2	548.8	45.7
6	1230	.0	300.0	.0	548.8	36.6
7	1235	127.6	300.0	42.5	676.4	37.6
8	1240	225.2	300.0	75.1	901.6	42.9
9	1245	63.5	300.0	21.2	965.2	40.2
10	1250	64.1	300.0	21.4	1029.3	38.1
11	1255	.0	300.0	.0	1029.3	34.3
12	1300	196.1	300.0	65.4	1225.4	37.1
13	1305	63.5	300.0	21.2	1288.9	35.8
14	1310	267.0	300.0	89.0	1555.9	39.9
15	1315	127.6	300.0	42.5	1683.5	40.1
16	1320	196.1	300.0	65.4	1879.6	41.8
17	1325	197.1	300.0	65.7	2076.7	43.3
18	1335	754.7	600.0	125.8	2831.4	52.4
19	1350	714.5	900.0	79.4	3545.9	56.3
20	1355	63.5	300.0	21.2	3609.4	54.7
21	1400	196.9	300.0	65.6	3806.4	55.2
22	1405	196.9	300.0	65.6	4003.3	55.6
23	1410	191.1	300.0	63.7	4194.4	55.9
24	1415	196.9	300.0	65.6	4391.4	56.3
25	1420	196.9	300.0	65.6	4588.3	56.6
26	1425	.0	300.0	.0	4588.3	54.6
27	1430	63.5	300.0	21.2	4651.8	53.5
28	1435	.0	300.0	.0	4651.8	51.7
29	1440	.0	300.0	.0	4651.8	50.0
30	1445	191.1	300.0	63.7	4842.9	50.4
31	1450	.0	300.0	.0	4842.9	48.9
32	1455	.0	300.0	.0	4842.9	47.5
33	1500	196.9	300.0	65.6	5039.9	48.0
34	1505	127.6	300.0	42.5	5167.5	47.8
35	1510	127.6	300.0	42.5	5295.1	47.7
36	1515	63.5	300.0	21.2	5358.6	47.0
37	1520	186.4	300.0	62.1	5545.0	47.4
38	1525	186.4	300.0	62.1	5731.4	47.8
39	1530	225.6	300.0	75.2	5957.0	48.4
40	1535	190.5	300.0	63.5	6147.5	48.8
41	1540	63.5	300.0	21.2	6211.1	48.1
42	1545	127.6	300.0	42.5	6338.7	48.0
43	1550	63.5	300.0	21.2	6402.2	47.4
44	1555	196.1	300.0	65.4	6598.3	47.8
45	1600	.0	300.0	.0	6598.3	46.8
46	1605	63.5	300.0	21.2	6661.8	46.3
47	1610	225.9	300.0	75.3	6887.7	46.9
48	1615	186.4	300.0	62.1	7074.1	47.2
49	1620	186.4	300.0	62.1	7260.5	47.5
50	1625	196.9	300.0	65.6	7457.5	47.8
51	1630	186.4	300.0	62.1	7643.9	48.1

	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELAPSED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
52	1705	792.5	2100.0	37.7	8436.4	46.9
53	1710	64.1	300.0	21.4	8500.5	46.5
54	1735	450.8	1500.0	30.1	8951.3	45.2
55	1740	225.6	300.0	75.2	9176.9	45.7
56	1745	63.5	300.0	21.2	9240.4	45.3
57	1750	127.0	300.0	42.3	9367.4	45.3
58	1755	196.1	300.0	65.4	9563.6	45.5
59	1800	186.4	300.0	62.1	9750.0	45.8
60	1805	.0	300.0	.0	9750.0	45.1
61	1810	196.1	300.0	65.4	9946.1	45.4
62	1815	64.1	300.0	21.4	10010.2	45.1
63	1820	185.6	300.0	61.9	10195.7	45.3
64	1825	368.3	300.0	122.8	10564.1	46.3
65	1830	127.0	300.0	42.3	10691.1	46.3
66	1835	63.5	300.0	21.2	10754.6	46.0
67	1840	64.1	300.0	21.4	10818.7	45.6
68	1845	63.5	300.0	21.2	10882.2	45.3
69	1850	.0	300.0	.0	10882.2	44.8
70	1855	.0	300.0	.0	10882.2	44.2
71	1900	191.1	300.0	63.7	11073.3	44.5
72	1910	867.0	600.0	144.5	11940.3	46.8
73	1915	225.9	300.0	75.3	12166.2	47.2
74	1920	127.0	300.0	42.3	12293.2	47.1
75	1925	196.1	300.0	65.4	12489.4	47.3
76	1930	197.1	300.0	65.7	12686.5	47.5
77	1935	127.0	300.0	42.3	12813.5	47.5
78	1940	368.3	300.0	122.8	13181.8	48.3
79	1945	196.1	300.0	65.4	13378.0	48.5
80	1950	266.4	300.0	88.8	13644.3	48.9
81	1955	127.0	300.0	42.3	13771.4	48.8
82	2000	254.6	300.0	84.9	14026.0	49.2
83	2005	.0	300.0	.0	14026.0	48.7
84	2010	63.5	300.0	21.2	14089.5	48.4
85	2015	191.1	300.0	63.7	14280.6	48.6
86	2020	127.0	300.0	42.3	14407.7	48.5
87	2025	393.2	300.0	131.1	14800.9	49.3
88	2030	417.9	300.0	139.3	15218.9	50.2
89	2035	254.6	300.0	84.9	15473.5	50.6
90	2040	196.1	300.0	65.4	15669.6	50.7
91	2045	127.6	300.0	42.5	15797.2	50.6
92	2050	127.0	300.0	42.3	15924.3	50.6
93	2055	63.5	300.0	21.2	15987.8	50.3
94	2100	63.5	300.0	21.2	16051.3	50.0
95	2105	191.1	300.0	63.7	16242.4	50.1
96	2110	63.5	300.0	21.2	16305.9	49.9
97	2115	266.4	300.0	88.8	16572.3	50.2
98	2120	127.6	300.0	42.5	16699.9	50.1
99	2125	196.1	300.0	65.4	16896.1	50.3
100	2130	224.9	300.0	75.0	17120.9	50.5
101	2135	63.5	300.0	21.2	17184.4	50.2
102	2140	254.6	300.0	84.9	17439.1	50.5

	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELASPED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
103	2145	.0	300.0	.0	17439.1	50.1
104	2150	225.2	300.0	75.1	17664.3	50.3
105	2155	224.9	300.0	75.0	17889.2	50.5
106	2200	225.2	300.0	75.1	18114.4	50.7
107	2205	63.5	300.0	21.2	18177.9	50.5
108	2210	267.0	300.0	89.0	18444.9	50.8
109	2215	127.0	300.0	42.3	18571.9	50.7
110	2220	63.5	300.0	21.2	18635.4	50.5
111	2225	254.6	300.0	84.9	18890.0	50.8
112	2350	2050.3	5100.0	40.2	20940.3	49.5
113	2355	185.6	300.0	61.9	21125.9	49.6
114	0000	.0	300.0	.0	21125.9	49.2
115	0005	.0	300.0	.0	21125.9	48.9
116	0010	185.6	300.0	61.9	21311.4	49.0
117	0015	185.6	300.0	61.9	21497.0	49.1
118	0020	127.6	300.0	42.5	21624.6	49.0
119	0025	186.4	300.0	62.1	21811.0	49.1
120	0030	417.9	300.0	139.3	22228.9	49.7
121	0035	127.6	300.0	42.5	22356.6	49.7
122	0040	266.0	300.0	88.7	22622.5	49.9
123	0045	.0	300.0	.0	22622.5	49.6
124	0050	64.1	300.0	21.4	22686.6	49.4
125	0055	224.9	300.0	75.0	22911.5	49.6
126	0100	225.9	300.0	75.3	23137.4	49.8
127	0106	185.6	360.0	51.5	23323.0	49.8
128	0115	190.5	540.0	35.3	23513.5	49.6
129	0120	64.1	300.0	21.4	23577.6	49.4
130	0125	225.9	300.0	75.3	23803.5	49.6
131	0135	809.1	600.0	134.9	24612.6	50.6
132	0140	377.3	300.0	125.8	24990.0	51.1
133	0145	.0	300.0	.0	24990.0	50.8
134	0157	196.1	720.0	27.2	25186.1	50.5
135	0203	63.5	360.0	17.6	25249.6	50.2
136	0210	.0	420.0	.0	25249.6	49.8
137	0230	185.6	1200.0	15.5	25435.2	49.0
138	0235	.0	300.0	.0	25435.2	48.7
139	0240	.0	300.0	.0	25435.2	48.4
140	0245	197.1	300.0	65.7	25632.3	48.5
141	0250	196.3	300.0	65.4	25828.6	48.6
142	0300	224.9	600.0	37.5	26053.5	48.5
143	0315	489.5	900.0	54.4	26542.9	48.6
144	0320	225.2	300.0	75.1	26768.1	48.8
145	0325	63.5	300.0	21.2	26831.7	48.6
146	0335	196.9	600.0	32.8	27028.6	48.4
147	0340	63.5	300.0	21.2	27092.1	48.3
148	0345	196.3	300.0	65.4	27288.4	48.4
149	0350	196.1	300.0	65.4	27484.5	48.5
150	0400	424.8	600.0	70.8	27909.3	48.7
151	0405	191.1	300.0	63.7	28100.4	48.8
152	0420	683.0	900.0	75.9	28783.4	49.2
153	0425	185.6	300.0	61.9	28969.0	49.3

	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELAPSED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
154	0435	196.1	600.0	32.7	29165.1	49.1
155	0440	63.5	300.0	21.2	29228.6	49.0
156	0445	196.1	300.0	65.4	29424.7	49.0
157	0450	.0	300.0	.0	29424.7	48.8
158	0455	63.5	300.0	21.2	29488.2	48.7
159	0500	224.9	300.0	75.0	29713.1	48.8
160	0535	1475.2	2100.0	70.2	31188.4	49.5
161	0540	190.5	300.0	63.5	31378.9	49.6
162	0545	.0	300.0	.0	31378.9	49.3
163	0550	186.4	300.0	62.1	31565.3	49.4
164	0555	196.1	300.0	65.4	31761.4	49.5
165	0600	63.5	300.0	21.2	31825.0	49.3
166	0605	.0	300.0	.0	31825.0	49.1
167	0610	.0	300.0	.0	31825.0	48.9
168	0615	196.1	300.0	65.4	32021.1	49.0
169	0620	63.5	300.0	21.2	32084.6	48.8
170	0625	267.0	300.0	89.0	32351.6	49.0
171	0630	186.4	300.0	62.1	32538.0	49.1
172	0635	.0	300.0	.0	32538.0	48.9
173	0640	.0	300.0	.0	32538.0	48.6
174	0645	63.5	300.0	21.2	32601.5	48.5
175	0650	190.5	300.0	63.5	32792.0	48.6
176	0655	186.4	300.0	62.1	32978.4	48.6
177	0700	.0	300.0	.0	32978.4	48.4
178	0705	64.1	300.0	21.4	33042.5	48.3
179	0710	127.6	300.0	42.5	33170.2	48.3
180	0715	368.3	300.0	122.8	33538.5	48.6
181	0720	197.1	300.0	65.7	33735.6	48.7
182	0725	.0	300.0	.0	33735.6	48.5
183	0730	196.1	300.0	65.4	33931.7	48.5
184	0739	318.2	540.0	58.9	34249.9	48.6
185	0743	127.0	240.0	52.9	34376.9	48.6
186	0745	.0	120.0	.0	34376.9	48.6
187	0750	127.6	300.0	42.5	34504.5	48.5
188	0755	315.6	300.0	105.2	34820.1	48.8
189	0800	63.5	300.0	21.2	34883.6	48.7
190	0810	315.1	600.0	52.5	35198.7	48.7
191	0815	63.5	300.0	21.2	35262.2	48.6
192	0820	315.6	300.0	105.2	35577.8	48.8
193	0825	191.1	300.0	63.7	35768.9	48.9
194	0850	482.3	1500.0	32.2	36251.2	48.5
195	0855	127.6	300.0	42.5	36378.8	48.5
196	0900	424.8	300.0	141.6	36803.6	48.9
197	0905	63.5	300.0	21.2	36867.1	48.8

	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELAPSED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
1	1235	.0	.0	.0	.0	.0
2	1240	224.9	300.0	75.0	224.9	75.0
3	1245	267.0	300.0	89.0	491.9	82.0
4	1250	393.1	300.0	131.0	884.9	98.3
5	1310	785.5	1200.0	65.5	1670.4	79.5
6	1315	186.4	300.0	62.1	1856.8	77.4
7	1320	377.3	300.0	125.8	2234.2	82.7
8	1325	63.5	300.0	21.2	2297.7	76.6
9	1330	393.2	300.0	131.1	2690.9	81.5
10	1340	557.5	600.0	92.9	3248.5	83.3
11	1345	64.1	300.0	21.4	3312.6	78.9
12	1350	196.1	300.0	65.4	3508.7	78.0
13	1355	267.0	300.0	89.0	3775.7	78.7
14	1400	266.0	300.0	88.7	4041.6	79.2
15	1405	197.1	300.0	65.7	4238.7	78.5
16	1410	224.9	300.0	75.0	4463.6	78.3
17	1415	.0	300.0	.0	4463.6	74.4
18	1425	186.4	600.0	31.1	4650.0	70.5
19	1440	1131.3	900.0	125.7	5781.3	77.1
20	1445	196.9	300.0	65.6	5978.3	76.6
21	1450	191.1	300.0	63.7	6169.4	76.2
22	1455	63.5	300.0	21.2	6232.9	74.2



	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELAPSED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
1	1025	.0	.0	.0	.0	.0
2	1030	196.9	300.0	65.6	196.9	65.6
3	1035	196.1	300.0	65.4	393.1	65.5
4	1040	63.5	300.0	21.2	456.6	50.7
5	1045	393.1	300.0	131.0	849.6	70.8
6	1050	372.0	300.0	124.0	1221.6	81.4
7	1110	641.9	1200.0	53.5	1863.5	69.0
8	1115	186.4	300.0	62.1	2049.9	68.3
9	1120	225.2	300.0	75.1	2275.1	68.9
10	1125	186.4	300.0	62.1	2461.5	68.4
11	1130	63.5	300.0	21.2	2525.0	64.7
12	1135	63.5	300.0	21.2	2588.6	61.6
13	1140	266.4	300.0	88.8	2854.9	63.4
14	1145	127.6	300.0	42.5	2982.6	62.1
15	1150	196.9	300.0	65.6	3179.5	62.3
16	1155	127.0	300.0	42.3	3306.5	61.2
17	1200	63.5	300.0	21.2	3370.0	59.1
18	1205	315.1	300.0	105.0	3685.1	61.4
19	1210	186.4	300.0	62.1	3871.5	61.5
20	1215	.0	300.0	.0	3871.5	58.7
21	1220	.0	300.0	.0	3871.5	56.1
22	1230	393.1	600.0	65.5	4264.6	56.9
23	1235	196.1	300.0	65.4	4460.7	57.2
24	1240	.0	300.0	.0	4460.7	55.1
25	1245	224.9	300.0	75.0	4685.6	55.8
26	1250	127.0	300.0	42.3	4812.6	55.3
27	1255	197.1	300.0	65.7	5009.7	55.7
28	1300	.0	300.0	.0	5009.7	53.9
29	1305	185.6	300.0	61.9	5195.3	54.1
30	1310	197.1	300.0	65.7	5392.4	54.5
31	1315	127.6	300.0	42.5	5520.0	54.1
32	1320	185.6	300.0	61.9	5705.6	54.3
33	1325	196.9	300.0	65.6	5902.5	54.7
34	1330	127.0	300.0	42.3	6029.5	54.3
35	1335	63.5	300.0	21.2	6093.1	53.4
36	1340	225.2	300.0	75.1	6318.3	54.0
37	1345	63.5	300.0	21.2	6381.8	53.2
38	1350	.0	300.0	.0	6381.8	51.9
39	1355	.0	300.0	.0	6381.8	50.6
40	1400	63.5	300.0	21.2	6445.3	50.0
41	1405	196.9	300.0	65.6	6642.2	50.3
42	1410	64.1	300.0	21.4	6706.3	49.7
43	1540	4239.7	5400.0	78.5	10946.1	57.9
44	1700	1855.3	4800.0	38.7	12801.4	54.0
45	1710	1112.1	600.0	185.4	13913.5	57.3
46	1720	424.8	600.0	70.8	14338.3	57.6
47	1735	266.4	900.0	29.6	14604.6	56.6
48	1745	196.9	600.0	32.8	14801.6	56.1
49	1805	2413.8	1200.0	201.2	17215.4	62.4
50	1810	225.9	300.0	75.3	17441.3	62.5
51	1815	63.5	300.0	21.2	17504.8	62.1

	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELASPED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
52	1820	127.0	300.0	42.3	17631.8	61.9
53	1830	266.4	600.0	44.4	17898.2	61.5
54	1835	196.1	300.0	65.4	18094.3	61.5
55	1840	64.1	300.0	21.4	18158.4	61.1
56	1845	63.5	300.0	21.2	18221.9	60.7
57	1850	225.6	300.0	75.2	18447.5	60.9
58	1855	127.6	300.0	42.5	18575.1	60.7
59	1900	127.0	300.0	42.3	18702.2	60.5
60	1905	63.5	300.0	21.2	18765.7	60.1
61	1910	64.1	300.0	21.4	18829.8	59.8
62	1915	224.9	300.0	75.0	19054.7	59.9
63	1920	127.6	300.0	42.5	19182.3	59.8
64	1925	63.5	300.0	21.2	19245.8	59.4
65	1930	63.5	300.0	21.2	19309.3	59.0

	TIME (LCL)	DISTANCE FROM PREVIOUS POSIT. (m)	ELAPSED TIME (seconds)	SWIMMING SPEED (cm/sec)	CUMULATIVE DISTANCE (m)	AVERAGE SPEED (cm/sec)
1	1059	.0	.0	.0	.0	.0
2	1101	.0	120.0	.0	.0	.0
3	1110	191.1	540.0	35.4	191.1	29.0
4	1115	127.0	300.0	42.3	318.2	33.1
5	1120	191.1	300.0	63.7	509.3	40.4
6	1125	63.5	300.0	21.2	572.8	36.7
7	1130	.0	300.0	.0	572.8	30.8
8	1135	196.1	300.0	65.4	768.9	35.6
9	1140	197.1	300.0	65.7	966.0	39.3
10	1145	196.1	300.0	65.4	1162.2	42.1
11	1150	127.0	300.0	42.3	1289.2	42.1
12	1155	.0	300.0	.0	1289.2	38.4
13	1200	197.1	300.0	65.7	1486.3	40.6
14	1205	190.5	300.0	63.5	1676.9	42.3
15	1210	185.6	300.0	61.9	1862.4	43.7
16	1215	63.5	300.0	21.2	1925.9	42.2
17	1220	64.1	300.0	21.4	1990.0	40.9
18	1225	225.6	300.0	75.2	2215.6	42.9
19	1230	.0	300.0	.0	2215.6	40.6
20	1235	196.1	300.0	65.4	2411.7	41.9
21	1240	185.6	300.0	61.9	2597.3	42.9
22	1245	254.6	300.0	84.9	2851.9	44.8
23	1250	63.5	300.0	21.2	2915.4	43.8
24	1255	225.9	300.0	75.3	3141.4	45.1
25	1300	63.5	300.0	21.2	3204.9	44.1
26	1305	127.6	300.0	42.5	3332.5	44.1
27	1310	196.1	300.0	65.4	3528.6	44.9
28	1315	63.5	300.0	21.2	3592.1	44.0
29	1320	196.9	300.0	65.6	3789.1	44.8
30	1325	225.2	300.0	75.1	4014.3	45.8
31	1330	127.0	300.0	42.3	4141.3	45.7
32	1335	225.9	300.0	75.3	4367.2	46.7
33	1340	127.0	300.0	42.3	4494.2	46.5
34	1345	127.6	300.0	42.5	4621.8	46.4
35	1350	196.1	300.0	65.4	4818.0	47.0
36	1355	.0	300.0	.0	4818.0	45.6
37	1400	190.5	300.0	63.5	5008.5	46.1
38	1405	225.9	300.0	75.3	5234.4	46.9
39	1420	424.4	900.0	47.2	5658.8	46.9
40	1425	127.6	300.0	42.5	5786.4	46.8
41	1430	266.6	300.0	88.9	6053.0	47.8
42	1435	197.1	300.0	65.7	6250.1	48.2
43	1440	186.4	300.0	62.1	6436.5	48.5
44	1445	63.5	300.0	21.2	6500.0	47.9
45	1520	1160.2	2100.0	55.2	7660.2	48.9