

# FEAR ~ EEA

# Shell OCSS Development


December 1, 1978  
Volume III

Prepared By:

United States Geological Survey  
Department of the Interior  
300 North Los Angeles Street  
Los Angeles, California 90013

State Lands Commission  
1807 13th Street  
Sacramento, California 95814

Port of Long Beach  
925 Harbor Plaza  
Long Beach, California 90801



EIR/EA  
Shell OCS Beta Unit Development

---

Technical Appendix

Volume III

December 1, 1978

Prepared By:

State Lands Commission  
1807 13th Street  
Sacramento, California 95814

Port of Long Beach  
925 Harbor Plaza  
Long Beach, California 90801

United States Geological Survey  
Department of the Interior  
300 North Los Angeles Street  
Los Angeles, California 90013

Technical Assistance:  
WESTEC Services, Inc.  
180 East Main Street, Suite 150  
Tustin, California 92680

APPENDIX  
TABLE OF CONTENTS

<u>APPENDIX</u>	<u>SUBJECT</u>	<u>PAGE</u>
I	Geotechnical Materials USGS Analysis	I-1
II	Marine Biology Species Lists	II-1
III	Navigation Aids/Marine Traffic	III-1
IV	Offshore Structure Collision Data	IV-1
V	Oil Spill Simulation Equations	V-1
VI	List of Coastal Beaches/Parks	VI-1
VII	Utility/Service Agency Correspondence	VII-1
VIII	Oceanographic Survey Results	VIII-1

APPENDIX I  
GEOTECHNICAL MATERIALS  
INCLUDING USGS ANALYSIS  
AND SUPPORT MATERIALS



**U.S. DEPARTMENT OF COMMERCE**  
**National Oceanic and Atmospheric Administration**  
 National Marine Fisheries Service  
 Southwest Region  
 300 South Ferry Street  
 Terminal Island, California 90731

November 17, 1977

FSW33/RSH

Mr. F. J. Schambeck  
 Department of Interior, Geological Survey  
 300 N. Los Angeles Street  
 Los Angeles, California 90012

NOTED - SCHAMBECK

NOTED - LAVELLE


Dear Mr. Schambeck:


NOTED - ADAMS

Subject: Platforms Ellen and Elly, OCS-P 0300 and OCS-P 0301,  
 San Pedro Bay, off Southern California

We do not feel that the proposed platforms will significantly affect those resources for which we have a responsibility. However, the location of the platforms could alter the normal migration route of the California gray whale (Eschrichtius robustus) and therefore should be considered in the environmental analysis.

Sincerely,

  
 Gerald V. Howard  
 Regional Director



United States Department of the Interior

FISH AND WILDLIFE SERVICE

500 NE MULTNOMAH STREET SUITE 1692  
PORTLAND OREGON 97232

NOTED - DUNAWAY

November 28, 1977

NOTED - ADAMS

To: Oil and Gas Supervisor, Pacific Area, USGS, Los Angeles, California

From: Regional Director, FWS, Portland, Oregon

Subject: Plan of Development, Beta Unit, OCS Leases P-0296, 0300, 0301, 0306, and Unleased Tract 0255

*File*

We have reviewed the subject Plan of Development (POD), Beta Platform Site Evaluations, and Report Pipeline Route Survey, San Pedro Shelf, California. We do not object to the concept as described in the POD.

We wish to review installation methods, pipeline corridors, and other material submitted in the future relative to the development of the subject platform(s).


We have noted that the general area is one of intense commercial fishing. Some impacts upon fish populations and fishing may occur in the event of an oil spill. Impacts upon fish and wildlife populations may be severe if an unforeseen spill reaches the few remaining lagoons and estuaries in the coastal area.

Our response has been submitted in accordance with the provisions of Section 4(f) 3 of Secretarial Order No. 2974 (revised) of January 19, 1977. Attached is the Plan of Development, site evaluation, and pipeline route survey for the Beta Unit.

Thank you for the opportunity to review this material and to provide our comments.

Attachments

*J. W. Teeter*  
Acting Regional Director





UNITED STATES  
DEPARTMENT OF THE INTERIOR  
GEOLOGICAL SURVEY

7211 FEDERAL BUILDING  
300 NO. LOS ANGELES STREET  
LOS ANGELES, CALIFORNIA 90012

An environmental review for the following activity has been conducted in accordance with Section 402.04 of Part 402, Chapter IV, Endangered Species Act of 1973 (16 U.S.C. 1531 et seq.):

SHELL OIL COMPANY AS OPERATOR  
PLAN OF OPERATION (DEVELOPMENT)  
BETA UNIT, PARCELS COVERED BY LEASES

OCS-P 0296 Block 37W, 34N  
OCS-P 0300 Block 37W, 33N  
OCS-P 0301 Block 36W, 33N  
OCS-P 0306 Block 36W, 32N  
and

Unleased Tract 255 Block 36W, 34N  
SAN PEDRO BAY OFF CALIFORNIA

The following determination has been made for this activity to identify if it may jeopardize the existence of any endangered species or result in the destruction or adverse modification of critical habitat.

1. The above activity will not jeopardize the continued existence of any endangered species or result in the destruction or adverse modification of critical habitat.
2. The above activity may jeopardize the continued existence of any endangered species or result in the destruction or adverse modification of critical habitat and a consultation is recommended with Fish and Wildlife Service and/or National Marine Fisheries Service.

F. J. Schrambeck  
Oil and Gas Supervisor

9/12/78  
Date

### 3.4 ARCHEOLOGICAL AND HISTORICAL SITE CLEARANCE DETERMINATION

The Bureau of Land Management was contacted for comments initially on November 29, 1977, and subsequently on July 31, 1978. Responses appear on the following pages.



## memorandum


1780.11 OCS P-0296, 0300, 0301, 0306

DATE: JAN 5 1978

REPLY TO  
ATTN OF: Manager, Pacific OCS OfficeSUBJECT: *File*  
Beta Unit Prospect, OCS Leases P-0296, -0300, -0301, -0306, and Unleased Tract 255, San Pedro Bay

JAN 5 1978

TO: Oil and Gas Supervisor, Pacific Area

We have reviewed Shell's Plan of Development Concept ~~LOS ANGELES~~ Comments are:

1. No legal conflicts nor encumbrances in the leases except that the "Beta Unit" is being proposed and that the Operator has not been designated. We have no legal objection to Shell's Plan of Development Concept.
2. No comment on Oil Spill Contingency Plan.
3. No impacts on biological resources.
4. Comments on cultural resources as follows:

NOTED - DUNAWAY

NOTED - SCHAMBECK

a. Platform: We have reviewed the reports submitted and the remote sensing data obtained by BB&N. An archeological assessment was made by Dr. Hole for the original drill sites proposed and he found "no indication of anything of archeological interest near the proposed drill sites in Block 262". He also found, with the exception of some anomalies on the north end of OCS P-0300, "...no other anomalies in either block." Dr. Hole did, however, state that "The magnetometry is technically good, but generally ineffective at the depths over most of this survey. The sensor head was never more than about 100' deep; thus, over much of the area it was too far above the bottom to record even very large anomalies."

Our review does, however, indicate numerous anomalies based on both the magnetometry and sidescan data. The deployment of the magnetometer high in the water column and on a 305 meter grid, may have also resulted in a failure to pick up other anomalies. From the data available, in our opinion, these anomalies cannot be reliably determined to be of a non-significant cultural resource nature. For this reason and the fact that the data quality precludes an effective assessment of cultural resources potential, we question whether these blocks can be considered clear.

b. Pipeline Routes: Cursory surveys of two of the three routes proposed indicated that numerous anomalies are present. To be confident that these are not culturally related, their source should be identified as geological, as non-significant cultural resources, or avoided. To avoid them, in many cases, would require more than a single line of data to locate them accurately. Also, these pipeline routes were not evaluated



Buy U.S. Savings Bonds Regularly on the Payroll Savings Plan

by a qualified archaeologist, therefore the necessary professional expertise was not utilized. In addition, there has been no determination as to whether the planned pipeline would be a "GS gathering line" or BLM permitted pipeline.

c. Conclusion: Due to the recent temporary BLM-GS agreement obliging us to utilize the existing cultural resources NTL for these reviews, we will not recommend withholding approval of the conceptual plan of development as it relates to platform sites and associated operations. However, we do believe you should be aware that, in our opinion, the cultural resources potential on these leases and pipeline routes has not been adequately evaluated. As to the pipeline, we are asking our Washington Office for guidance pertaining to the BLM/GS jurisdictional question. We expect to have additional opportunities to comment as more specific plans are finalized.

5. Comments on camouflage stipulation as follows:

This stipulation applies to both leases with platform proposals. Reported visibility ranges for the San Pedro Bay-Catalina Channel area exceed 10 nautical miles 22% of the time in January and 31% of the time in July. Thus, the platforms will probably be visible from shore for significant time periods. We look forward to working with you on color or pattern selections. It might be beneficial for Shell to employ the services of a professional in the design arts to aid in both the structural design and color selection to lessen visual impact.

We recommend approval of Shell's Beta Unit Plan of Development Concept.

We are returning the following information which was forwarded to us with your memorandum of November 29, 1977, and additional information which was subsequently received informally from you:

1. Beta Unit Plan of Development
2. Oil Spill Contingency Plan
3. Critical Operations and Curtailment Plans
4. Report Pipeline Route Survey
5. Geoseismic Investigation of Palos Verde Fault Zone
6. An Environmental Design Study
7. Soil Boring, Sampling & Testing Program Tract 261
8. Soil Boring, Sampling & Testing Program Tract 262
9. Beta Platform Site Evaluations
10. BB&N Remote Sensing Data and Report

Enclosures

*William E. Grant*

## memorandum

DATE: AUG 07 1978

REPLY TO  
ATTN OF: Manager, Pacific OCS Office

1780-11

OCS P-0296, 0300, 0301, 0306



SUBJECT: Cultural Resource Survey, Proposed Beta Unit  
Plan of DevelopmentTO:  
Oil and Gas Supervisor, Pacific Area

We have reviewed the enclosed Dames and Moore's report concerning the section on cultural resource surveys and our comments are:

The report indicates that the cultural survey study along the proposed pipeline route from the proposed production platform Elly to Port of Long Beach was conducted by specialists in marine archaeology and history. They have concluded that no potential sites or features of possible cultural values are located in the survey area. Based on their conclusion we recommend approval of the section on cultural survey.

We are returning the following report which was forwarded to us with your memorandum of July 31, 1978: Dames and Moore. July 1978. Report Marine Geophysical and Cultural Resource Surveys Proposed Pipeline Route Offshore Long Beach, California for Shell Oil Company.

Enclosure



I-7

Buy U.S. Savings Bonds Regularly on the Payroll Savings Plan



### 3.5 REVIEW COMMENTS AND RELATED CORRESPONDENCE

Cooperative proposal review and environmental report planning has been an ongoing process with the State of California since early October 1977. Shell's Special Report to the Governor of California of September 29, 1977 was received by the Governor's office on October 3, 1977. In accordance with 30 CFR 250.34, a copy of the Plan of Development (POD) was transmitted to the Governor. Mr. Allan Lind of the Office of Planning and Research (OPR) requested an additional 20 copies for State Agency review, which was honored. By letter of January 25, 1978, Mr. Bill Press, the Director of OPR, transmitted State comments to Shell Oil Company and the USGS. Responding State Agencies were:

- OPR
- Air Resources Board
- State Lands Commission
- Department of Fish and Game
- Department of Navigation and Ocean Development
- Department of Conservation
  - Division of Mines and Geology
  - Division of Oil and Gas
- State Water Resources Control Board

On March 23, 1978, Mr. William F. Northrup, Executive Officer of the State Lands Commission, sent letters by certified mail to those agencies which were determined, under the provisions of the California Environmental Quality Act of 1970, as amended, and subsequent guidelines; to be "responsible agencies" in the Shell Beta Project. By this letter, the State Lands Commission and the Port of Long Beach requested, pursuant to Public Resources Code Section 21080.4(a), a written statement of the scope and content of the environmental information in connection with the proposed project which was germane to each agency's statutory responsibilities and which was to be included in the EIR. Attached were a proposed outline for the subject environmental document, and a project description. Responses were received from:

- United States Coast Guard
- California Department of Transportation
- California Regional Water Quality Control Board,  
Los Angeles Region
- California Division of Oil and Gas

- California Coastal Commission
- South Coast Air Quality Management District
- City of Huntington Beach
- State Water Quality Resources Control Board
- Los Angeles County Flood Control District
- City of Long Beach

Department of Fire  
Department of Planning and Building

- California Air Resources Board

Additionally, comments were received from:

- Office of Historic Preservation, Department of Parks and Recreation

### 3.6 MAPS AND ILLUSTRATIONS

Maps and illustrations appear as appropriate throughout the document and are not segregated in a single appendix.



UNITED STATES  
DEPARTMENT OF THE INTERIOR  
GEOLOGICAL SURVEY

7760 Federal Building  
300 North Los Angeles Street  
Los Angeles, California 90012

September 8, 1978

MEMORANDUM


To: Oil and Gas Supervisor, Pacific Area  
From: District Geologist, Los Angeles  
Subject: Geologic Hazard Evaluation for Beta Platforms Ellen and Elly

Shell's proposal to site a development drilling platform (Ellen) at  $x=1,428,175$   $y=520,220$  and a production platform (Elly) at  $x=1,428,310$   $y=520,580$  (California Zone 6) was reviewed by Scott Hamlin, Don Krotser, Richard Tudor, and Roger Nielson.

The site proposed for these platforms is relatively well chosen within this geologically complex area, between two surface faults and just inshore from the current and paleo shelf breaks. There is indication of slumping beyond the shelf break, but none in the more gradual gradient inshore. Possible indications of gas seeps have been identified as close as 500 feet away from the site(s) in contractor interpretations, but are not confirmable in the perhaps incomplete geophysical data profiles submitted.

Foundation stability is discussed in a report to Shell by Robert Pyke, in which reference is made to a final geotechnical report by Woodward Clyde Consultants dated April 1978. This reference was not submitted, so evaluation of Pyke's report of negligably small risk of instability is left unsubstantiated, though plausible.

Drilling from Platform Ellen will be subject to hazards of subsurface gas and faulting indicated in geophysical data and in previous drilling.

  
for Keith A. Yenne

KAY/DJK/sk

U.S. GEOLOGICAL SURVEY  
CONSERVATION DIVISION

To: District Engineer, Ventura

From: Los Angeles District Office

Subject: Geologic Hazards Analysis of APD

Operator: Shell

Lease Number: OCS P-0300

Well Number: Platforms Ellen and Elly

Surface Location: Ellen - X = 1,428,175 Y = 520,220 (Drill)  
Elly - X = 1,428,310 Y = 520,580 (Production)

Deviated Location: \_\_\_\_\_

Elly 255 ft.

Water Depth: Ellen 265 ft.,

Total Well Depth: \_\_\_\_\_

A. Surface Hazards:

1. Slope stability

Slope at the site is  $2^{\circ}$  SE. Two areas of disturbed recent sediments lie 2000 ft. SW and 1500 ft. SE of the proposed site below the paleo shelf break. Geophysical profiles show evidence of slump movement in these areas.

2. Slumping

The possibility of slumping at the site appears to be minor.

3. Surface faulting

The site lies adjacent to the Palos Verdes Fault Zone, 500 ft. NE of the Beta Fault, a NE branch of the Palos Verdes Fault.

4. Seeps

Two questionable areas of oil and/or gas seeps lie 500 ft. east and west of the site and uncertain seeps are linked with the disturbed sediments to the SE in the contracted hazard interpretation. No seeps were observed in the submitted geophysical records.

5. Seismicity

The site is located along the seismically active Palos Verdes Fault Zone. The maximum credible earthquake magnitude for this area is 7. At the site the associated bedrock acceleration would equal 0.35 g.

B. Subsurface Hazards:1. Potential oil or gas reservoir, depth

The reservoir occupies seven subdivisions from 2,700 to 4,700 ft. subsea within the Lower Pliocene and Upper Miocene correlating to shales and sandstones of the Repetto and Monterey formations. The reservoir is composed of Delmontian sands, shales, and silts.

2. Possible shallow gas zones, depth

Shallow gas zones are expected to be encountered at various depths (see P-0300 #1 drilling record). Geophysical profiles show numerous bright spot anomalies from 25 to 500 ft. subsea.

3. Possible surface cutting faults which may be encountered in drilling, approximate depths

None seen in the geophysical profiles.

4. Possible subsurface faults which may be encountered in drilling, approximate depths

The Beta and Palos Verdes faults form updip boundaries of the reservoir between 2500 and 5000 ft. below the sea bottom. These faults are likely to be encountered at depth in subsequent development drilling for production.



Lease Number: OCS P-0300

Well Number: Platform Ellen and Elly

5. Possible fresh or salt water flows, depth

Unknown

6. Possible lost circulation zones, depth

Some of the previous wells drilled in OCS leases P-0296, 0300, and 0301 encountered temporary lost circulation. Recovery occurred spontaneously or was accomplished by increasing the drilling fluid viscosity in these wells.

7. Possible abnormal pressure zones, depth

As indicated from review of the drilling records of leases P-0300, P-0296, and 0301, no abnormal pressures are expected.

8. Remarks

Foundation conditions:

We are unable to evaluate the geotechnical character of the platform locations for the following reasons:

1. Data was not submitted for coreholes in the immediate vicinity of the platform sites.
2. The submitted geotechnical reports (Site Stability Studies and Soil Boring, Sampling, and Laboratory Testing Program) are based upon coreholes located outside of the immediate platform site area.

Date: Sept. 11, 1978

Signed: *P. B. Ballantyne I*

Initials: SNH, DJK, RBT, RGN

Rev. 3-78 Hazard Unit

*Acting Dist. Geol.*

APPENDIX II  
MARINE BIOLOGY SPECIES LISTS

TABLE I

SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ALAMITOS BAY AT EACH STATION DURING QUARTERLY TRAWLING SURVEYS, 1971-72 (FROM ENVIRONMENTAL QUALITY ANALYSTS, INC., AND MARINE BIOLOGICAL CONSULTANTS, INC., 1973b).

Species	Stations													
	RW7	RW8	RW9	RW10	RW12	RW13	Total	RW7	RW8	RW9	RW10	RW12	RW13	Total
	October 1971							May 1972						
CARCHARHINIDAE														
<u>Mustelus henlei</u>												1		1
SQUALIDAE														
<u>Squalus acanthias</u>												1		1
RHINOBATIDAE														
<u>Platyrrhinoidis triseriata</u>														
<u>Rhinobatos productus</u>	1						1					10		10
DASYATIDAE														
<u>Urolophus halleri</u>														
MYLIOBATIDAE														
<u>Myliobatis californica</u>			1				1	1				4		5
ENGRAULIDAE														
<u>Anchoa compressa</u>						1	1							
<u>A. delicatissima</u>														
<u>Engraulis mordax</u>	244	7			26	1	278	2		21			9	32
BATRACHOIDIDAE														
<u>Porichthys myriaster</u>													2	2
<u>P. notatus</u>								2		3	2			7
OPHIDIDAE														
<u>Otophidium scrippsae</u>						1	1							
<u>O. taylori</u>													1	1
SYNGNATHIDAE														
<u>Syngnathus griseolineatus</u>													1	1
<u>Syngnathus sp.</u>														
SERRANIDAE														
<u>Paralabrax nebulifer</u>					1	1	2							
POMADASYIDAE														
<u>Anisotremus davidsoni</u>														
SCIAENIDAE														
<u>Genyonemus lineatus</u>			6		2	11	19	167	14	418	38	51	210	898
<u>Menticirrhus undulatus</u>		8				1	9	1				9		10
<u>Seriphus politus</u>					1		1	61	48	113	24	741	39	1026
<u>Umbrina roncadore</u>														
EMBIOTOCIDAE														
<u>Amphistichus argenteus</u>	1	6					7			2		13	5	20
<u>Cymatogaster aggregata</u>			12		16	12	40	106	6	371	86	6	199	774

TABLE 1 (Cont.)  
 SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ALAMITOS BAY AT EACH STATION DURING QUARTERLY  
 TRAWLING SURVEYS, 1971-72 (FROM ENVIRONMENTAL QUALITY ANALYSTS, INC., AND MARINE BIOLOGICAL  
 CONSULTANTS, INC., 1973b).

Species	October 1971							May 1972						
	RW7	RW8	RW9	RW10	RW12	RW13	Total	RW7	RW8	RW9	RW10	RW12	RW13	Total
EMBIOTOCIDAE (Cont)														
<u>Embiotoca jacksoni</u>										9	1	2	2	14
<u>Hyperprosopon argenteum</u>		20				36	56	120	6	96	5	47	22	296
<u>Phanerodon furcatus</u>	1	19	20	35	28	102	205	420	24	269	172	87	135	1107
<u>Rhacochilus vacca</u>														
CLINIDAE														
<u>Heterostichus rostratus</u>														
STROMATEIDAE														
<u>Peprilus simillimus</u>								1					1	
COTTIDAE														
<u>Hemilepidotus spinosus</u>														
<u>Leptocottus armatus</u>	2	1					3	3					3	
BOTHIDAE														
Bothidae, unid.					2		2							
<u>Citharichthys stigmaeus</u>	56	16	131	13	35	145	346	13		92	67	9	16	197
<u>Hippoglossina stomata</u>	1						1							
<u>Paralichthys californicus</u>	10	28			17	2	57			2		7	1	10
PLEURONECTIDAE														
<u>Hypsopsetta guttulata</u>			1		2	1	4			2		2	1	5
<u>Parophrys vetulus</u>											2			2
<u>Pleuronichthys decurrens</u>										3	2			5
<u>P. ritteri</u>														
<u>P. verticalis</u>	2	1	6	4	3	15	31	6		20	30		3	59
CYNOGLOSSIDAE														
<u>Symphurus atricauda</u>	1	1	1	1	2	11	17			30	26		12	68
Number of Individuals	319	107	178	53	135	340	1132	908	98	1451	455	990	658	4555
Number of Species	10	10	8	4	12	14	21	13	5	15	12	15	16	26
Diversity Index	1.11	2.77	1.40	1.28	2.76	2.27		2.21	1.90	2.68	2.60	1.51	2.44	

TABLE 1 (CONT.)

SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ALAMITOS BAY AT EACH STATION DURING QUARTERLY TRAWLING SURVEYS, 1971-72 (FROM ENVIRONMENTAL QUALITY ANALYSTS, INC., AND MARINE BIOLOGICAL CONSULTANTS, INC., 1973b).

Species	Stations													Grand Total	
	RW7	RW8	RW9	RW10	RW12	RW13	Total	RW7	RW8	RW9	RW10	RW12	RW13		Total
	September 1972						November 1972								
CARCHARHINIDAE															
<u>Mustelus henlei</u>															1
SQUALIDAE															
<u>Squalus acanthias</u>															1
RHINOBATIDAE															
<u>Platyrrhinoidis triseriata</u>	1					1	2		5	1				6	8
<u>Rhinobatos productus</u>	11	8	6		3	11	39								50
DASYATIDAE															
<u>Urolophus halleri</u>					2		2					1	3	4	6
MYLIOBATIDAE															
<u>Myliobatis californica</u>			4			1	5						2	2	13
ENGRAULIDAE															
<u>Anchoa compressa</u>	16	12	19	5		10	62	8	161	79	6		52	306	369
<u>A. delicatissima</u>										41			3	44	44
<u>Engraulis mordax</u>	130	226	285	47	384	277	1349	7	73	343	1807	385	48	2663	4322
BATRACHOIDIDAE															
<u>Porichthys myriaster</u>		2				4	6								8
<u>P. notatus</u>													1	1	8
OPHIDIDAE															
<u>Otophidium scrippsae</u>															1
<u>O. taylori</u>								2			2			4	5
SYNGNATHIDAE															
<u>Syngnathus griseolineatus</u>															1
<u>Syngnathus sp.</u>			1				1		2					2	3
SERRANIDAE															
<u>Paralabrax nebulifer</u>			1				1								3
POMADASYIDAE															
<u>Anisotremus davidsoni</u>			1				1								1
SCIAENIDAE															
<u>Genyonemus lineatus</u>	47	30	78	147	1	77	380	69	54	93	206	4	113	539	1836
<u>Menticirrhus undulatus</u>	3	1	50		1	2	57		14			5		19	95
<u>Seriphus politus</u>	3	14		814		173	1004	9	41	240	574	2	199	1065	3096
<u>Umbrina roncadore</u>										1				1	1
EMBIOTOCIDAE															
<u>Amphistichus argenteus</u>	4	11	2			1	18		17					17	62
<u>Cymatogaster aggregata</u>	60	12	272	9	5	1	359	3	3	3	5	1	33	48	1221

TABLE 1 (Cont.)  
 SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ALAMITOS BAY AT EACH STATION DURING QUARTERLY  
 TRAWLING SURVEYS, 1971-72 (FROM ENVIRONMENTAL QUALITY ANALYSTS, INC., AND MARINE BIOLOGICAL  
 CONSULTANTS, INC., 1973b).

Species	September 1972							November 1972							Grand Total
	RW7	RW8	RW9	RW10	RW12	RW13	Total	RW7	RW8	RW9	RW10	RW12	RW13	Total	
EMBIOTOCIDAE (Cont)															
<u>Embiotoca jacksoni</u>	4	13	2	6			25				5		1	6	45
<u>Hyperprosopon argenteum</u>	34	24	61	16			135	2	30	4	5		23	64	551
<u>Phanerodon furcatus</u>			95	66	4	1	166	2	22	2	2	52	131	211	1689
<u>Rhacochilus vacca</u>									1					1	1
CLINIDAE															
<u>Heterostichus rostratus</u>									1					1	1
STROMATEIDAE															
<u>Peprilus simillimus</u>				22			22						7	7	30
COTTIDAE															
<u>Hemilepidotus spinosus</u>				1			1								1
<u>Leptocottus armatus</u>	2	2	1	1			6	2	1					3	15
BOTHIDAE															
Bothidae, unid.															2
<u>Citharichthys stigmaeus</u>	6	5	16	25		2	54		10	3	18	2	28	61	708
<u>Hippoglossina stomata</u>															1
<u>Paralichthys californicus</u>	12	3	7		14	7	43	1	4			14	2	21	131
PLEURONECTIDAE															
<u>Hypsopsetta guttulata</u>	1		3	1	1		6				2			2	17
<u>Parophrys vetulus</u>															2
<u>Pleuronichthys decurrens</u>													1	1	6
<u>P. ritteri</u>									1					1	1
<u>P. verticalis</u>	1	1	11	42			55		1		14	1	9	25	170
CYNOGLOSSIDAE															
<u>Symphurus atricauda</u>				137			137			2	81		17	100	322
Number of Individuals	335	364	915	1339	415	568	3936	105	441	812	2727	467	673	5225	14848
Number of Species	16	15	19	15	9	14	26	10	18	12	13	10	18	29	41
Diversity Index	2.75	2.20	2.72	2.08	0.56	1.90		1.89	2.91	2.11	1.48	0.99	2.89		

TABLE 2

SPECIES AND COMMON NAMES, FREQUENCY OF CAPTURE, AND RELATIVE ABUNDANCE OF FISHES CAPTURED BY TRAWLING OFFSHORE OF HUNTINGTON BEACH, 1976-78 (AFTER MARINE BIOLOGICAL CONSULTANTS, INC., 1977, 1978)

Species	Common Name	Number Captured	Frequency of Capture	Relative Abundance (%)
<u>August 1976</u>				
<u>Seriphus politus</u>	queenfish	1822	12	42.3
<u>Genyonemus lineatus</u>	white croaker	1773	12	41.1
<u>Engraulis mordax</u>	northern anchovy	356	12	8.3
<u>Hyperprosopon argenteum</u>	walleye surfperch	145	9	3.4
<u>Peprilus simillimus</u>	Pacific butterflyfish	68	7	1.6
<u>Phanerodon furcatus</u>	white surfperch	59	9	1.4
<u>Amphistichus argenteus</u>	barred surfperch	18	7	0.4
<u>Citharichthys stigmaeus</u>	speckled sanddab	14	6	0.3
<u>Symphurus atricauda</u>	Pacific tonguefish	10	6	0.2
<u>Cymatogaster aggregata</u>	shiner surfperch	7	4	0.2
<u>Paralichthys californicus</u>	Pacific halibut	7	5	0.2
<u>Embiotoca jacksoni</u>	black surfperch	6	2	0.1
<u>Menticirrhus undulatus</u>	California corbina	5	4	0.1
<u>Syngnathus leptorhynchus</u>	bay pipefish	5	4	0.1
<u>Pleuronichthys verticalis</u>	hornyhead turbot	5	3	0.1
<u>Chilara taylori</u>	spotted cusk-eel	3	3	0.07
<u>Rhinobatos productus</u>	shovelnose guitarfish	2	2	0.05
<u>Mustelus henlei</u>	brown smoothhound	1	1	0.02
<u>Torpedo californica</u>	Pacific electric ray	1	1	0.02
<u>Otophidium scrippsi</u>	basketweave cusk-eel	1	1	0.02
<u>Damalichthys vacca</u>	pile surfperch	1	1	0.02
Total Number Captured	4309			
Number of Species	21			
<u>August 1977</u>				
<u>Genyonemus lineatus</u>	white croaker	3743	12	46.4
<u>Engraulis mordax</u>	northern anchovy	3126	11	38.7
<u>Phanerodon furcatus</u>	white surfperch	275	11	3.4
<u>Hyperprosopon argenteum</u>	walleye surfperch	254	12	3.1
<u>Amphistichus argenteus</u>	barred surfperch	206	11	2.5
<u>Seriphus politus</u>	queenfish	134	12	1.7
<u>Citharichthys stigmaeus</u>	speckled sanddab	85	9	1.1
<u>Paralichthys californicus</u>	California halibut	80	11	1.0
<u>Cymatogaster aggregata</u>	shiner surfperch	62	9	0.8
<u>Syngnathus californiensis</u>	kelp pipefish	39	8	0.5
<u>Symphurus atricauda</u>	California tonguefish	11	5	0.1
<u>Platyrrhinoidis triseriata</u>	thornback	9	2	0.1
<u>Leptocottus armatus</u>	staghorn sculpin	8	5	0.1
<u>Pleuronichthys verticalis</u>	hornyhead turbot	6	3	0.07
<u>Synodus lucioceps</u>	California lizardfish	5	3	0.06
<u>Porichthys myriaster</u>	specklefin midshipman	4	3	0.05
<u>Mustelus henlei</u>	brown smoothhound	3	2	0.04

TABLE 2 (Cont.)  
 SPECIES AND COMMON NAMES, FREQUENCY OF CAPTURE, AND RELATIVE ABUNDANCE  
 OF FISHES CAPTURED BY TRAWLING OFFSHORE OF HUNTINGTON BEACH, 1976-78  
 (AFTER MARINE BIOLOGICAL CONSULTANTS, INC., 1977, 1978) (Cont)

Species	Common Name	Number Captured	Frequency of Capture	Relative Abundance (%)
<u>August 1977 (Cont)</u>				
<u>Rhinobatos productus</u>	shovelnose guitarfish	3	2	0.04
<u>Menticirrhus undulatus</u>	California corbina	3	3	0.04
<u>Xystreurus liolepis</u>	fantail sole	3	2	0.04
<u>Pleuronichthys ritteri</u>	spotted turbot	2	1	0.04
<u>Otophidium scrippsi</u>	basketweave cusk-eel	1	1	0.01
<u>Sebastes paucispinis</u>	bocaccio	1	1	0.01
<u>Girella nigricans</u>	opaleye	1	1	0.01
<u>Damalichthys vacca</u>	pile surfperch	1	1	0.01
<u>Heterostichus rostratus</u>	giant kelpfish	1	1	0.01
<u>Peprilus simillimus</u>	Pacific butterflyfish	1	1	0.01
<u>Hypsosetta guttulata</u>	diamond turbot	1	1	0.01
Total Number Captured	8068			
Number of Species	28			
<u>February 1978</u>				
<u>Genyonemus lineatus</u>	white croaker	6030	14	74.8
<u>Seriphus politus</u>	queenfish	1272	13	15.8
<u>Engraulis mordax</u>	northern anchovy	220	13	2.7
<u>Citharichthys stigmaeus</u>	speckled sanddab	94	12	1.2
<u>Anchoa compressa</u>	deepbody anchovy	92	10	1.1
<u>Syngnathus exilis</u>	pipefish	70	11	0.87
<u>Cymatogaster aggregata</u>	shiner surfperch	58	11	0.72
<u>Amphistichus argenteus</u>	barred surfperch	34	4	0.42
<u>Otophidium scrippsi</u>	basketweave cusk-eel	32	6	0.40
<u>Pleuronichthys verticalis</u>	hornyhead turbot	27	7	0.34
<u>Paralichthys californicus</u>	California halibut	22	11	0.27
<u>Menticirrhus undulatus</u>	California corbina	19	7	0.24
<u>Symphurus atricauda</u>	California tonguefish	17	6	0.21
<u>Phanerodon furcatus</u>	white surfperch	14	7	0.17
<u>Synodus lucioceps</u>	California lizardfish	14	5	0.17
<u>Hyperprosopon argenteum</u>	walleye surfperch	13	3	0.16
<u>Rhinobatos productus</u>	shovelnose guitarfish	5	5	0.06
<u>Xystreurus liolepis</u>	faintail sole	5	3	0.06
<u>Pleuronichthys ritteri</u>	spotted turbot	4	2	0.05
<u>Myliobatis californica</u>	bat ray	3	3	0.04
<u>Platyrrhinoidis triseriata</u>	thornback	3	3	0.04
<u>Urolophus halleri</u>	round stingray	3	3	0.04
<u>Paralabrax nebulifer</u>	barred sand bass	2	2	0.03
<u>Scorpaena guttata</u>	California scorpion	2	2	0.03
<u>Squalus acanthias</u>	spiny dogfish	1	1	0.01



TABLE 2 (Cont.)  
 SPECIES AND COMMON NAMES, FREQUENCY OF CAPTURE, AND RELATIVE ABUNDANCE  
 OF FISHES CAPTURED BY TRAWLING OFFSHORE OF HUNTINGTON BEACH, 1976-78  
 (AFTER MARINE BIOLOGICAL CONSULTANTS, INC., 1977, 1978) (Cont)

Species	Common Name	Number Captured	Frequency of Capture	Relative Abundance (%)
<u>February 1978 (Cont)</u>				
<u>Atractoscion nobilis</u>	white sea bass	1	1	0.01
<u>Umbrina roncadore</u>	yellowfin croaker	1	1	0.01
<u>Xenistius californiensis</u>	salema	1	1	0.01
<u>Lepidogobius lepidus</u>	bay goby	1	1	0.01
<u>Peprilus simillimus</u>	Pacific butterflyfish	1	1	0.01
<u>Sebastes serranoides</u>	live rockfish	1	1	0.01
Total Number Captured	8062			
Number of Species	31			

TABLE 3  
SPECIES AND COMMON NAMES, FREQUENCY OF CAPTURE, AND RELATIVE ABUNDANCE  
OF FISHES CAPTURED BY TRAWLING OFFSHORE OF ALAMITOS BAY, FEBRUARY 1978  
(AFTER MARINE BIOLOGICAL CONSULTANTS, INC., 1978)

Species	Common Name	Number Captured	Frequency of Capture	Relative Abundance (%)
<u>Genyonemus lineatus</u>	white croaker	3782	8	70.6
<u>Symphurus atricauda</u>	California tonguefish	791	6	14.8
<u>Seriphus politus</u>	queenfish	345	8	6.4
<u>Engraulis mordax</u>	northern anchovy	114	7	2.1
<u>Anchoa compressa</u>	deepbody anchovy	98	8	1.8
<u>Citharichthys stigmaeus</u>	speckled sanddab	36	7	0.67
<u>Cymatogaster aggregata</u>	shiner surfperch	32	7	0.60
<u>Tilapia mossambica</u>	tilapia	31	3	0.58
<u>Otophidium scrippsi</u>	basketweave cusk-eel	23	4	0.43
<u>Syngnathus exilis</u>	pipefish	22	1	0.41
<u>Phanerodon furcatus</u>	white surfperch	13	6	0.24
<u>Paralichthys californicus</u>	California halibut	11	6	0.21
<u>Hypsopsetta guttulata</u>	diamond turbot	8	6	0.15
<u>Pleuronichthys verticalis</u>	hornyhead turbot	8	5	0.15
<u>Menticirrhus undulatus</u>	California corbina	7	5	0.13
<u>Sebastes dallii</u>	callico rockfish	6	2	0.11
<u>Embiotoca jacksoni</u>	black surfperch	6	3	0.11
<u>Synodus lucioceps</u>	California lizardfish	5	7	0.09
<u>Syngnathus sp.</u>	pipefish	4	2	0.07
<u>Hyperprosopon argenteum</u>	walleye surfperch	4	3	0.07
<u>Porichthys myriaster</u>	specklefin midshipman	3	2	0.06
<u>Chilara taylori</u>	spotted cusk-eel	3	2	0.06
<u>Platyrhinoidis triseriata</u>	thornback	1	1	0.02
<u>Rhinobatos productus</u>	shovelnose guitarfish	1	1	0.02
<u>Atherinops affinis</u>	topsmelt	1	1	0.02
<u>Pleuronichthys ritteri</u>	spotted turbot	1	1	0.02
Total Number Captured		5356		
Number of Species		26		

TABLE 4  
 SPECIFIC AND COMMON NAMES, NUMBER, RANKED ABUNDANCE, AND FREQUENCY OF  
 CAPTURE OF FISHES TAKEN BY TRAWLING SEPTEMBER, 1975 (FROM LONG BEACH  
 HARBOR CONSULTANTS, 1976)

Species	Common Name	Number Captured	Relative Abundance (%)	Frequency of Capture
<u>Genyonemus lineatus</u>	white croaker	3,437	55.5	21
<u>Engraulis mordax</u>	northern anchovy	993	16.0	18
<u>Phanerodon furcatus</u>	white surfperch	445	7.2	17
<u>Symphurus atricauda</u>	California tonguefish	353	5.7	18
<u>Cymatogaster aggregata</u>	shiner surfperch	282	4.6	11
<u>Sebastes dallii</u>	calico rockfish	261	4.2	11
<u>Seriphus politus</u>	queenfish	229	3.7	13
<u>Citharichthys stigmaeus</u>	speckled sanddab	81	1.3	8
<u>Pleuronichthys verticalis</u>	hornyhead turbot	38	0.6	10
<u>Lepidogobius lepidus</u>	bay goby	24	0.4	10
<u>Sebastes auriculatus</u>	brown rockfish	12	0.2	4
<u>Embiotoca jacksoni</u>	black surfperch	7	0.1	5
<u>Citharichthys sordidus</u>	Pacific sanddab	5	0.1	2
<u>Porichthys myriaster</u>	specklefin midshipman	4	0.1	3
<u>Hyperprosopon argenteum</u>	walleye surfperch	4	0.1	1
<u>Paralichthys californicus</u>	California halibut	3	0.1	2
<u>Squalus acanthias</u>	spiny dogfish	2	0.03	1
<u>Chilara taylori</u>	spotted cusk-eel	2	0.03	1
<u>Otophidium scrippsi</u>	basketweave cusk-eel	2	0.03	2
<u>Paralabrax clathratus</u>	kelp bass	2	0.03	2
<u>Rhinobatos productus</u>	shovelnose guitarfish	1	0.02	1
<u>Torpedo californica</u>	Pacific electric ray	1	0.02	1
<u>Damalichthys vacca</u>	pile surfperch	1	0.02	1
<u>Paralabrax nebulifer</u>	barred sandbass	1	0.02	1
<u>Peprilus simillimus</u>	Pacific butterfish	1	0.02	1
<u>Sebastes paucispinis</u>	bocaccio	1	0.02	1
<u>Artedius lateralis</u>	smoothhead sculpin	1	0.02	1
<u>Leptocottus armatus</u>	staghorn sculpin	1	0.02	1
<u>Odontopyxis trispinosa</u>	pygmy poacher	1	0.02	1
<u>Lepidopsetta bilineata</u>	rock sole	1	0.02	1
<u>Parophrys vetulus</u>	English sole	1	0.02	1
Total Number Captured		6,197		

TABLE 5  
SPECIES, NUMBER, AND RANK OF FISHES CAPTURED BY  
GILL NETTING IN LONG BEACH HARBOR, 1974-78 (FROM  
ENVIRONMENTAL QUALITY ANALYSTS AND MARINE  
BIOLOGICAL CONSULTANTS, INC., 1978).

Species Name	Number Captured	Rank
<u>Genyonemus lineatus</u>	1,811	1
<u>Seriphus politus</u>	1,102	2
<u>Cymatogaster aggregata</u>	901	3
<u>Phanerodon furcatus</u>	632	4
<u>Engraulis mordax</u>	554	5
<u>Hyperprosopon argenteum</u>	314	6
<u>Embiotoca jacksoni</u>	218	7
<u>Porichthys myriaster</u>	127	8
<u>Atherinops affinis</u>	108	9
<u>Damalichthys vacca</u>	89	10
<u>Trachurus symmetricus</u>	70	11
<u>Menticirrhus undulatus</u>	61	12
<u>Synodus lucioceps</u>	53	13
<u>Cheilotrema saturnum</u>	50	14
<u>Paralabrax nebulifer</u>	30	15
<u>Mustelus henlei</u>	26	16
<u>Paralichthys californicus</u>	23	17
<u>Mustelus californicus</u>	22	18
<u>Sardinops sagax caeruleus</u>	22	18
<u>Girella nigricans</u>	20	20
<u>Squalus acanthias</u>	13	21
<u>Sphyræna argentea</u>	13	21
<u>Medialuna californiensis</u>	13	21
<u>Rhacochilus toxotes</u>	13	21
<u>Sebastes seranoides</u>	12	25
<u>S. paucispinis</u>	11	26
<u>Roncador sternsii</u>	8	27
<u>Atherinopsis californiensis</u>	7	28
<u>Paralabrax clathratus</u>	7	28
<u>Anisotremus davidsoni</u>	7	28
<u>Sarda chiliensis</u>	7	28
<u>Paralabrax maculatofasciatus</u>	6	32
<u>Scomber japonicus</u>	6	32
<u>Heterodontus francisii</u>	5	34
<u>Urolophus halleri</u>	5	34
<u>Myliobatis californica</u>	5	34
<u>Amphistichus argenteus</u>	5	34
<u>Peprilus simillimus</u>	5	34
<u>Symphurus atricauda</u>	5	34
<u>Triakis semifasciata</u>	4	40
<u>Paralabrax clathratus</u>	4	40

TABLE 5 (Cont.)  
 SPECIES, NUMBER, AND RANK OF FISHES CAPTURED BY  
 GILL NETTING IN LONG BEACH HARBOR, 1974-78 (FROM  
 ENVIRONMENTAL QUALITY ANALYSTS AND MARINE  
 BIOLOGICAL CONSULTANTS, INC., 1978) (Cont)

Species Name	Number Captured	Rank
<u>Atractoscion nobilis</u>	4	40
<u>Brachyistius frenatus</u>	4	40
<u>Hyperprosopon anale</u>	4	40
<u>Atherinopsis californiensis</u>	3	45
<u>Scorpaena guttata</u>	3	45
<u>Pleuronichthys verticalis</u>	3	45
<u>Sebastes gordini</u>	2	48
<u>Heterostichus rostratus</u>	2	48
<u>Hypsopsetta guttulata</u>	2	48
<u>Anchoa compressa</u>	1	51
<u>Sebastes melanops</u>	1	51
<u>S. mystinus</u>	1	51
<u>S. rastrelliger</u>	1	51
<u>Arteichius lateralis</u>	1	51
<u>Stereolepis gigas</u>	1	51
<u>Umbrina roncadore</u>	1	51
<u>Hermosilla azurea</u>	1	51
<u>Hyperprosopon ellipticum</u>	1	51
<u>Micrometrus minimus</u>	1	51
<u>Chromis punctipinnis</u>	1	51
<u>Mugil cephalus</u>	1	51
<u>Acanthogobius flavimanus</u>	1	51
<u>Citharichthys stigmaeus</u>	1	51
<u>Sebastes sp.</u>	1	
Number of Individuals	6436	
Number of Species	64	

TABLE 6

SPECIES, NUMBER, AND RANK OF FISHES CAPTURED BY  
TRAWLING IN LONG BEACH HARBOR, 1974-78 (FROM  
ENVIRONMENTAL QUALITY ANALYSTS AND MARINE  
BIOLOGICAL CONSULTANTS, INC., 1978)

Species Name	Number Captured	Rank
<u>Genyonemus lineatus</u>	68,547	1
<u>Engraulis mordax</u>	28,898	2
<u>Seriphus politus</u>	15,601	3
<u>Lepidogobius lepidus</u>	8,159	4
<u>Symphurus atricauda</u>	5,483	5
<u>Phanerodon furcatus</u>	3,693	6
<u>Cymatogaster aggregata</u>	2,756	7
<u>Peprilus simillimus</u>	1,869	8
<u>Sebastes dallii</u>	1,742	9
<u>Anchoa compressa</u>	748	10
<u>Synodus lucioceps</u>	664	11
<u>Citharichthys stigmaeus</u>	654	12
<u>Porichthys myriaster</u>	511	13
<u>Pleuronichthys verticalis</u>	158	14
<u>Chilara taylori</u>	144	15
<u>Embiotoca jacksoni</u>	120	16
<u>Paralichthys californicus</u>	68	17
<u>Paralabrax nebulifer</u>	56	18
<u>Syngnathus sp.</u>	50	19
<u>Damalichthys vacca</u>	50	19
<u>Sebastes miniatus</u>	34	21
<u>Hyperprosopon argenteum</u>	26	22
<u>Otophidium scrippsi</u>	25	23
<u>Porichthys notatus</u>	19	24
<u>Sebastes auriculatus</u>	19	24
<u>Odontopyxis trispinosa</u>	16	26
<u>Sebastes paucispinis</u>	11	27
<u>Leptocottus armatus</u>	9	28
<u>Trachurus symmetricus</u>	9	28
<u>Citharichthys sordidus</u>	8	30
<u>Xystreurus liolepis</u>	7	31
<u>Squalus acanthias</u>	5	32
<u>Sebastes mystinus</u>	4	33
<u>Myliobatos californica</u>	3	34
<u>Urolophus halleri</u>	3	34
<u>Scorpaena guttata</u>	3	34
<u>Sebastes saxicola</u>	3	34
<u>S. serranoides</u>	3	34
<u>Rhacochilus toxotes</u>	3	34
<u>Hippoglossina stomata</u>	3	34
<u>Rhinobatos productus</u>	2	41

TABLE 6 (Cont.)

SPECIES, NUMBER, AND RANK OF FISHES CAPTURED BY  
 TRAWLING IN LONG BEACH HARBOR, 1974-78 (FROM  
 ENVIRONMENTAL QUALITY ANALYSTS AND MARINE  
 BIOLOGICAL CONSULTANTS, INC., 1978) (Cont)

Species Name	Number Captured	Rank
<u>Anchoa delicatissima</u>	2	41
<u>Atherinops affinis</u>	2	41
<u>Artedius lateralis</u>	2	41
<u>Neoclinus uninotatus</u>	2	41
<u>Pleuronichthys decurrens</u>	2	41
<u>Mustelus henlei</u>	1	47
<u>Leuresthes tenuis</u>	1	47
<u>Sebastes semicinctus</u>	1	47
<u>Atroctoscion nobilis</u>	1	47
<u>Cheilotrema saturnum</u>	1	47
<u>Heterostichus rostratus</u>	1	47
<u>Clevelandia ios</u>	1	47
<u>Ilypnus gilberti</u>	1	47
<u>Parophrys vetulus</u>	1	47
Engraulididae, unid.	13	
Sebastes sp.	10	
Embiotocidae, unid.	4	
Gobiidae, unid.	2	
Number of Individuals	104,235	
Number of Species	55	

TABLE 7

QUALITATIVE LIST OF THE INTERTIDAL ANIMALS COLLECTED FROM THE  
FOUR OFFSHORE OIL ISLANDS, SPRING 1976  
(FROM SOUTHERN CALIFORNIA OCEAN STUDIES CONSORTIUM, 1977)

Species	Grissom	White	Chaffee	Freeman
PORIFERA				
<u>Leucandra heathi</u>			X	
<u>Leucosolenia eleanor</u>			X	
<u>Xestospongia vanilla</u>			X	
COELENTERATA				
HYDROZOA				
<u>Obelia</u> sp.	X		X	X
<u>Plumularia</u> sp.			X	X
<u>Sertularia</u> sp.	X			
<u>Stauridiosarsia</u>			X	
<u>Syncoryne</u> sp.	X			X
<u>Tubularia</u> sp.	X			
ANTHOZOA				
<u>Anthopleura elegantissima</u>	X	X	X	X
<u>A. xanthogrammica</u>	X	X	X	X
<u>Metridium senile</u>				X
PLATYHELMINTHES				
TURBELLARIA				
polyclads, unid.	X		X	X
NEMERTEA				
<u>Emplectonema gracile</u>	X	X		X
<u>Lineus ruber</u>				X
<u>Nemertopsis gracilis</u>	X			
<u>Paranemertes peregrina</u>				X
Nemertea, unid.	X		X	X
ANNELIDA				
POLYCHAETA				
<u>Anaitides williamsi</u>		X		
<u>Arabella iricolor</u>	X	X	X	X
<u>A. semimaculata</u>	X	X	X	X
<u>Armandia brevis</u>		X		
<u>Autolytus</u> sp.				X
<u>Axiothella rubrocincta</u>			X	
<u>Boccardia proboscidea</u>	X	X		X
<u>Boccardia</u> sp.		X	X	X
<u>Capitella capitata</u>		X		
<u>Chaetozone setosa</u>				X
<u>Chrysopetalum occidentale</u>				X
<u>Cirratulus cirratus</u>		X		
<u>Cirriformia luxuriosa</u>	X	X	X	X
<u>C. spirabrancha</u>	X	X	X	X



TABLE 7 (Cont.)

QUALITATIVE LIST OF THE INTERTIDAL ANIMALS COLLECTED FROM THE  
FOUR OFFSHORE OIL ISLANDS, SPRING 1976  
(FROM SOUTHERN CALIFORNIA OCEAN STUDIES CONSORTIUM, 1977)

Species	Grissom	White	Chaffee	Freeman
ANNELIDA				
POLYCHAETA (Cont)				
<u>Dodecaceria fewkesi</u>				X
<u>Dorvillea rudolphi</u>			X	
<u>Eudistylia polymorpha</u>			X	
<u>Eulalia quadrioculata</u>	X	X	X	
<u>Eulalia sp.</u>		X	X	X
<u>Eumida bifoliata</u>	X	X		X
<u>E. sanguinea</u>				X
<u>Eupomatus gracilis</u>	X	X	X	X
<u>Exogone uniformis</u>			X	
<u>Genetyllis castanea</u>			X	X
<u>Halosydna brevisetosa</u>	X	X	X	X
<u>Langerhansia heterochaeta</u>			X	
<u>Lumbrineris zonata</u>		X		X
<u>Naineris dendritica</u>	X	X	X	X
<u>Nereis grubei</u>	X	X	X	X
<u>N. lateascens</u>	X	X		X
<u>Odontosyllis phosphorea</u>			X	X
<u>Ophiodromus pugettensis</u>			X	X
<u>Paleanotus bellis</u>			X	X
<u>Phragmatopoma californica</u>		X	X	X
<u>Platynereis bicanaliculata</u>		X	X	
<u>Polydora sp.</u>	X	X	X	X
<u>Polyophthalmus pictus</u>		X	X	X
<u>Pseudopotamilla ocellata</u>		X		X
<u>Sabella crassicornis</u>			X	
<u>S. media</u>		X	X	X
<u>Sabellaria cementarium</u>		X	X	
<u>Serpula vermicularis</u>		X		
<u>Sigambra bassi</u>				X
<u>Spirobinae, unid.</u>			X	X
<u>Steggoa californiensis</u>			X	
<u>Syllis gracilis</u>			X	
<u>Tharyx sp.</u>	X			X
<u>Thelepus setosus</u>			X	
<u>Trypanosyllis gemmipara</u>		X		
<u>Typosyllis aciculata</u>				X
<u>T. fasciata</u>	X	X	X	
<u>Typosyllis sp.</u>				X
SIPUNCULOIDEA				
<u>Phascolosoma agassizi</u>			X	

TABLE 7 (Cont.)

QUALITATIVE LIST OF THE INTERTIDAL ANIMALS COLLECTED FROM THE  
FOUR OFFSHORE OIL ISLANDS, SPRING 1976  
(FROM SOUTHERN CALIFORNIA OCEAN STUDIES CONSORTIUM, 1977)

Species	Grissom	White	Chaffee	Freeman
MOLLUSCA				
POLYPLACOPHORA				
<u>Cyanoplax dentiens</u>			X	
<u>C. hartwegii</u>				X
<u>Lepidozona cooperi</u>			X	
<u>L. mertensii</u>				X
<u>Mopalia ciliata</u>				X
<u>M. hindsii</u>		X	X	X
<u>M. lowei</u>			X	
<u>M. mucosa</u>	X	X	X	X
<u>Nuttallina fluxa</u>	X	X	X	X
<u>Stenoplax conspicua</u>			X	
<u>S. fallax</u>			X	
GASTROPODA				
PROSOBRANCHIATA				
<u>Acanthina spirata</u>			X	
<u>Amphissa versicolor</u>			X	
<u>Ceratostoma nuttalli</u>			X	X
<u>Collisella conus</u>				X
<u>C. digitalis</u>	X	X	X	X
<u>C. limatula</u>	X	X	X	X
<u>C. pelta</u>				X
<u>C. scabra</u>	X	X	X	X
<u>C. strigatella</u>	X		X	X
<u>Crepidula onyx</u>	X	X	X	X
<u>Crepidatella lingulata</u>	X	X	X	X
<u>Diodora aspera</u>				X
<u>Haliotis cracherodii</u>	X	X	X	X
<u>H. fulgens</u>				X
<u>Iselica fenestrata</u>		X		
<u>Littorina planaxis</u>	X	X	X	X
<u>L. scutulata</u>	X		X	
<u>Lottia gigantea</u>	X	X	X	X
<u>Megatebennus bimaculatus</u>			X	X
<u>Megathura crenulata</u>		X	X	X
<u>Mitrella carinata</u>	X	X	X	X
<u>M. tuberosa</u>			X	
<u>Nassarius mendicus</u>		X		
<u>N. tegula</u>			X	
<u>Norrisia norrisi</u>		X		
<u>Notoacmaea fenestrata</u>	X			
<u>N. insessa</u>	X	X	X	
<u>N. persona</u>	X			
<u>Ocenebra poulsoni</u>	X	X		X
<u>Serpulorbis squamigerous</u>		X	X	X
<u>Tegula brunnea</u>			X	
<u>T. eiseni</u>	X	X	X	X
<u>T. funebris</u>	X	X	X	X
<u>T. gallina</u>		X	X	X

TABLE 7 (Cont.)  
 QUALITATIVE LIST OF THE INTERTIDAL ANIMALS COLLECTED FROM THE  
 FOUR OFFSHORE OIL ISLANDS, SPRING 1976  
 (FROM SOUTHERN CALIFORNIA OCEAN STUDIES CONSORTIUM, 1977)

Species	Grissom	White	Chaffee	Freeman
MOLLUSCA				
GASTROPODA				
OPISTHOBRANCHIATA				
<u>Coryphella trilineata</u>		X		X
<u>Dendronotus frondosus</u>		X		
<u>D. iris</u>		X		
<u>Hemissenda crassicornis</u>		X		
<u>Navanax inermis</u>			X	
<u>Surilla oliviae</u>			X	
PELECYPODA				
<u>Chama pellucida</u>		X	X	X
<u>Chione undatella</u>	X			
<u>Chlamys hastata</u>			X	X
<u>Cooperella subdiaphana</u>	X			
<u>Cumingia californica</u>			X	
<u>Hiatella arctica</u>	X	X	X	X
<u>Hinnites multirugosus</u>		X	X	X
<u>Leptopecten latiauratus</u>		X		
<u>Modiolus demissus</u>		X		
<u>Mytilus californianus</u>		X	X	X
<u>M. edulis</u>	X	X	X	X
<u>Ostrea lurida</u>	X		X	
<u>Penitella penita</u>		X		
<u>Petricola californiensis</u>			X	
<u>Pitar newcombianus</u>				X
<u>Pododesmus cepio</u>	X			X
<u>Protothaca staminea</u>	X	X	X	X
<u>Saxidomus nuttalli</u>			X	
<u>Tresus nuttallii</u>			X	
CEPHALOPODA				
<u>Octopus bimaculatus</u>	X		X	
ARTHROPODA				
CRUSTACEA				
COPEPODA				
<u>Tigriopus californicus</u>		X	X	
CIRRIPEDIA				
<u>Balanus amphitrite</u>	X			
<u>B. crenatus</u>	X	X	X	X
<u>B. glandula</u>	X	X	X	X
<u>B. tintinnabulum</u>	X	X	X	X
<u>Chthamalus dalli</u>	X	X		X
<u>C. fissus</u>	X	X	X	X
<u>Pollicipes polymerus</u>	X	X	X	X
<u>Tetraclita squamosa rubescens</u>	X	X	X	X

TABLE 7 (Cont.)  
 QUALITATIVE LIST OF THE INTERTIDAL ANIMALS COLLECTED FROM THE  
 FOUR OFFSHORE OIL ISLANDS, SPRING 1976  
 (FROM SOUTHERN CALIFORNIA OCEAN STUDIES CONSORTIUM, 1977)

Species	Grissom	White	Chaffee	Freeman
ARTHROPODA				
CRUSTACEA				
MALACOSTRACA				
AMPHIPODA				
<u>Ampithoe humeralis</u>		X		X
<u>A. plumulosa</u>	X	X	X	
<u>Caprella californica</u>	X			
<u>C. equilibra</u>	X		X	
<u>Ceradocus spinicaudus</u>			X	
<u>Corophium acherusicum</u>	X	X	X	X
<u>Elasmopus rapax</u>	X		X	X
<u>Ericthonius brasiliensis</u>			X	X
<u>Hyale anceps</u>	X	X		
<u>H. frequens</u>				X
<u>Jassa falcata</u>	X		X	X
<u>Najna sp.</u>				X
<u>Orchestia traskiana</u>		X		X
<u>Parallorchestes ochotensis</u>			X	
<u>Paramoera mohri</u>	X			
<u>Parapleustes nautilus</u>	X			
<u>Photis brevipes</u>			X	
<u>Podocerus brasiliensis</u>			X	X
ISOPODA				
<u>Cirolana hardfordi</u>		X	X	X
<u>Dynamenella sp.</u>	X	X		
<u>Ianiropsis kincaidi kincaidi</u>			X	
<u>I. minuta</u>				X
<u>Idothea sp.</u>				X
<u>Jaeropsis dubia paucispinis</u>			X	
<u>Ligia occidentalis</u>	X	X	X	X
<u>Limnoria tripunctata</u>	X	X		
<u>Paracerceis sp.</u>	X	X		X
TANAIDACEA				
<u>Anatanais normani</u>		X	X	X
<u>Leptochelia dubia</u>		X	X	X
<u>Pancolus californiensis</u>	X			
CUMACEA				
Cumacean, unid.			X	
DECAPODA				
<u>Alpheus sp.</u>			X	
<u>Bataeus longidactylus</u>			X	
<u>Cancer antennarius</u>			X	
<u>C. productus</u>		X		
<u>Hemigrapsus oregonensis</u>	X			
<u>Lecythorhynchus sp.</u>		X		
<u>Lophopanopeus bellus</u>			X	

TABLE 7 (Cont.)  
 QUALITATIVE LIST OF THE INTERTIDAL ANIMALS COLLECTED FROM THE  
 FOUR OFFSHORE OIL ISLANDS, SPRING 1976  
 (FROM SOUTHERN CALIFORNIA OCEAN STUDIES CONSORTIUM, 1977)

Species	Grissom	White	Chaffee	Freeman
ARTHROPODA				
CRUSTACEA				
MALACOSTRACA				
DECAPODA				
<u>Opisthopus transverus</u>		X		
<u>Pachycheles rudis</u>		X	X	X
<u>Pachygrapsus crassipes</u>	X	X	X	X
<u>Pagurus hirsutiusculus</u>			X	
<u>P. samuelis</u>	X	X	X	X
<u>Pelia tumida</u>			X	
<u>Petrolisthes cinctipes</u>	X	X	X	X
<u>Pugettia producta</u>	X		X	X
<u>P. richii</u>			X	
PYCNOGONIDA				
Pycnogonids, unid.		X	X	
ENTOPROCTA				
<u>Barentsia sp.</u>		X		X
ECTOPROCTA				
<u>Bowerbankia gracilis</u>	X		X	X
<u>Bugula californica</u>	X			X
<u>B. neritina</u>		X	X	X
<u>Crisia occidentalis</u>			X	X
<u>Crisulipora occidentalis</u>	X	X	X	X
<u>Cryptosula pallasiana</u>	X	X	X	X
<u>Filicrisia franciscana</u>			X	X
<u>Membranipora membranacea</u>	X	X	X	
<u>M. tuberculata</u>	X		X	
<u>Schizoporella unicornis</u>	X	X		X
<u>Thalamoporella californica</u>		X	X	
<u>Victorella pavida</u>				X
ECHINODERMATA				
ASTEROIDEA				
<u>Asterometris sertulifera</u>				X
<u>Patiria miniata</u>		X	X	X
<u>Pisaster brevispinus</u>		X	X	
<u>P. giganteus</u>	X	X	X	
<u>P. ochraceus</u>	X	X	X	X
ECHINOIDEA				
<u>Strongylocentrotus franciscanus</u>		X	X	X
<u>S. purpuratus</u>		X	X	X
OPHIUROIDEA				
<u>Amphipholis squamata</u>			X	
<u>Ophiactis simplex</u>		X	X	
<u>Ophionereis annulata</u>				X
HOLOTHUROIDEA				
<u>Cucumaria miniata</u>			X	
<u>Stichopus californicus</u>			X	X

TABLE 7 (Cont.)  
 QUALITATIVE LIST OF THE INTERTIDAL ANIMALS COLLECTED FROM THE  
 FOUR OFFSHORE OIL ISLANDS, SPRING 1976  
 (FROM SOUTHERN CALIFORNIA OCEAN STUDIES CONSORTIUM, 1977)

Species	Grissom	White	Chaffee	Freeman
CHORDATA				
UROCHORDATA				
<u>Aplidium californicum</u>	X			
<u>Archidistoma ritteri</u>	X			
<u>Ascidia ceratodes</u>	X			X
<u>Botrylloides diegensis</u>	X			X
<u>Botryllus sp.</u>	X			X
<u>Ciona intestinalis</u>	X			
<u>Diplosoma pizoni</u>	X			
<u>Distaplia occidentalis</u>	X			
<u>Perophora annectens</u>				X
<u>Pyura haustor</u>	X			
<u>Styela plicata</u>	X			X

TABLE 8  
 NUMBER OF SPECIES AND INDIVIDUALS OF INTERTIDAL ORGANISMS COLLECTED FROM GRISSOM  
 ISLAND ON JANUARY 17 THROUGH 19, 1977.  
 (FROM SOUTHERN CALIFORNIA OCEAN STUDIES CONSORTIUM, 1977)

Species	Station 1			Station 2			Station 3			Station 4		
	H*	M	L	H	M	L	H	M	L	H	M	L
PORIFERA												
<u>Incerta sp.</u>		S*	S			S						C
<u>Leucilla nuttingi</u> sponge, unid.						S						S
sponge, unid.									S			
COELENTERATA												
HYDROZOA												
<u>Obelia sp.</u>			S									S
<u>Phialidium sp.</u>			S									
<u>Tubularia sp.</u>			C						S	S		C
ANTHOZOA												
<u>Anthopleura elegantissima</u>						A						
<u>A. xanthogrammica</u>			S			S		S	S			S S
PLATYHELMINTHES												
<u>Polycladia, unid.</u>		S				8			7			4 56
NEMERTEA												
<u>Nemertean, unid.</u>			156			8		22				4
NEMATODA												
<u>Nematode, unid.</u>					160		4	52			4	320
ANNELIDA												
POLYCHAETA												
<u>Armandia bioculata</u>			4			48						
<u>Capitella capitata</u>			80			112						4
<u>Capitita ambiseta</u>						8						
<u>Cirriiformia spirabrancha</u>									1			
<u>Cirriiformia sp.</u>			12									
<u>Eulalia avioulaseta</u>									1			
<u>Eupomatus gracilis</u>	S	C	4		3	C		10	A		30	14
<u>Halosydna brevisetosa</u>						8					4	2
<u>H. johnsoni</u>						40		2	S			
<u>Lumbrineris sp.</u>						8						
<u>Naineris dendritica</u>			8			8			1			12
<u>Nereis grubea</u>					4	16						
<u>Nereis sp.</u>												2
<u>Notomasus tenuis</u>						40						
<u>Ophryotrocha sp.</u>						40						
<u>Paleonotus bellis</u>						32			3			
<u>Pionosyllis gigantea</u>						16			1			
<u>Polydora limicola</u>						64			5		2	28
<u>Polyopthalmus pictus</u>						24			12			4
<u>Sabellidae, unid.</u>			4									2
<u>Syllidae, unid.</u>						8						
<u>Tharyx spp.</u>						56						
<u>Typosyllis fasciata</u>						760			64			

TABLE 8 (Cont.)  
 NUMBER OF SPECIES AND INDIVIDUALS OF INTERTIDAL ORGANISMS COLLECTED FROM GRISSOM  
 ISLAND ON JANUARY 17 THROUGH 19, 1977.  
 (FROM SOUTHERN CALIFORNIA OCEAN STUDIES CONSORTIUM, 1977)

Species	Station 1			Station 2			Station 3			Station 4		
	H	M	L	H	M	L	H	M	L	H	M	L
ARTHROPODA												
CRUSTACEA												
OSTRACODA												
<u>Cyldroleberis mariae</u>			4									
COPEPODA												
Harpacticoida, unid.			C			A		A	C			C
CIRRIPIEDIA												
<u>Balanus glandula</u>	18	918	C	270	378	C	36	117	C	36	6	8
<u>B. tintinnabulum</u>								2				
<u>Chthamalus fissus</u>	3340	267	C	7625	792	C	1296	486	C	1332	1935	C
<u>Tetraclita squamosa</u>		S	C		17	C		3	C	S	3	
MALOCOSTRACA												
CUMACEA												
<u>Cumella</u> sp.			20			40		14				14
TANAIDACEA												
<u>Anatanais normani</u>			28		6				2			374
ISOPODA												
<u>Ianiropsis tridens</u>			136									
<u>Ligia occidentalis</u>										A		A
<u>Limnoria tripunctata</u>												14
<u>Munna</u> sp.						16						
AMPHIPODA												
<u>Ampithoe</u> sp.												1
<u>Caprella equilibra</u>			4									
<u>C. verrucosa</u>			4									
<u>Corophium acherusicum</u>								1			6	13
<u>Elasmopus rapax</u>			112			2		15				
<u>Gammaropsis thompsoni</u>											2	
<u>Jassa falcata</u>											6	16
<u>Maera simiule</u>			28					1				
<u>Podocerus</u> sp.												4
<u>Stenothoe valida</u>					2	2						
DECAPODA												
<u>Pachygrapsus crassipes</u>	S	A	A		A	A	C	A	2	S	A	A
<u>Petrolithes cinctipes</u>			C						C			
LABIATA												
INSECTA												
Coleoptera, unid.								2			2	
PYCNOGONIDA												
Pycnogonida, unid.											4	S
MOLLUSCA												
POLYPLACOPHORA												
<u>Mopalia hindsii</u>			1			S						
<u>M. muscosa</u>		S				S		S	C			S



TABLE 8 (Cont.)  
 NUMBER OF SPECIES AND INDIVIDUALS OF INTERTIDAL ORGANISMS COLLECTED FROM GRISSOM  
 ISLAND ON JANUARY 17 THROUGH 19, 1977.  
 (FROM SOUTHERN CALIFORNIA OCEAN STUDIES CONSORTIUM, 1977)

Species	Station 1			Station 2			Station 3			Station 4		
	H	M	L	H	M	L	H	M	L	H	M	L
MOLLUSCA (Cont)												
GASTROPODA												
<u>Acanthina spirata</u>										S		
<u>Bankia setacea</u>												2
<u>Collisella digitalis</u>	C				C		9			3		
<u>C. limatula</u>	S	C	C		17	C	S	36		S	36	
<u>C. persona</u>											3	
<u>C. scabra</u>	C	150	C	160	233	C	54	387		72	207	S
<u>Crepidula onyx</u>												2
<u>Crepidula sp.</u>			4									
<u>Crepidatella lingulata</u>		S	S								2	
<u>Diodora aspera</u>			S									
<u>Littorina planaxis</u>	21			20			3			A	C	
<u>Lottia gigantea</u>								3				
<u>Megathura crenulata</u>		S	S									S
<u>Mitrella carinata</u>								S	1			14
<u>Norrisia norrisii</u>		S	S									
<u>Ocenebra poulsoni</u>								S	S			
<u>Serpulorbis squamigerus</u>												S
<u>Tegula funebris</u>	S	S			S					S		
PELECYPODA												
<u>Chama pellucida</u>			S									
<u>Cumingia californica</u>						16						
<u>Hiatella arctica</u>		3	12			32		6	3		12	14
<u>Kellia laperousii</u>												14
<u>Leptopecten sp.</u>			S									S
<u>Mytilus edulis</u>	S	24	311		40	852		4	16	S	16	56
<u>Pelecypod, unid.</u>			4									4
<u>Petricola sp.</u>						40						2
<u>Protothaca sp.</u>						64						
ECTOPROCTA												
<u>Bugula neritina</u>												A
<u>Crisulipora occidentalis</u>											S	
<u>Cryptosula pallasiana</u>		A	A		A	A		A	A		A	A
<u>Membranipora sp.</u>			S								C	
ENTOPROCTA												
<u>Barentsia sp.</u>												A
												A
ECHINODERMATA												
ECHINOIDEA												
<u>Strongylocentrotus purpuratus</u>			C			S				S		C
ASTEROIDEA												
<u>Pisaster ochraceous</u>			C			C		C	A			C
OPHIUROIDEA												
<u>Amphipholis squamata</u>			44							1		8
<u>Ophionereis annulata</u>											2	

TABLE 8 (Cont.)

NUMBER OF SPECIES AND INDIVIDUALS OF INTERTIDAL ORGANISMS COLLECTED FROM GRISSOM ISLAND ON JANUARY 17 THROUGH 19, 1977.  
(FROM SOUTHERN CALIFORNIA OCEAN STUDIES CONSORTIUM, 1977)

Species	Station 1			Station 2			Station 3			Station 4		
	H	M	L	H	M	L	H	M	L	H	M	L
CHORDATA												
UROCHORDATA												
ASCIDIACEA												
<u>Botryllus</u> sp.												
<u>Ciona intestinalis</u>				S							S	C
VERTEBRATA												
OSTEICHTHYS												
<u>Gobiesox eugrammus</u>							S					
CHLOROPHYTA												
<u>Ulva</u> sp.							A		A			A
PHAEOPHYTA												
<u>Egregia</u> sp.							C		A			S
RHODOPHYTA												
<u>Gelidium</u> sp.							C					
<u>Gigartina</u> sp.				S								A
<u>Polysiphonia</u> sp.												S
Total Number of Specimens	3379	1362	980	8075	1492	2528	1398	1115	172	1443	2306	988
Total Number of Species	10	17	46	4	14	50	7	29	34	11	35	46
* H = High Tide Zone	M = Mid Tide Zone			L = Low Tide Zone								
S = Sparse (1-10)	C = Common (11-500)			A = Abundant (>50)								

TABLE 9

SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch
Trawl Date: 2/26/75									
<u>Chilara taylori</u>		1		1	1	3	1	78	0.21
<u>Chitonotus pugetensis</u>		1	27	1	1	41	103	174	5.13
<u>Citharichthys sordidus</u>		20		181	3	37	121	362	10.67
<u>C. stigmaeus</u>	40	355	244		142	41	198	1020	30.06
<u>C. xanthostigma</u>					7			7	0.21
<u>Cymatogaster aggregata</u>	1	2						3	0.09
<u>Embiotoca jacksoni</u>	1							1	0.03
<u>Genyonemus lineatus</u>	2							2	0.06
<u>Hippoglossina stomata</u>		4	1	1	9	7	4	26	0.77
<u>Hydrolagus colliei</u>							1	1	0.03
<u>Icelinus quadriseriatus</u>		4	64	80	198	421	396	984	29.00
<u>Lepidogobius lepidus</u>			3		1			4	0.12
<u>Microstomus pacificus</u>		108		6		1		115	3.39
<u>Odontopyxis trispinosa</u>			3		2	5	4	14	0.41
<u>Ophiodon elongatus</u>						1		1	0.03
<u>Paralichthys californicus</u>	5				1			6	0.18
<u>Parophrys vetulus</u>	6	222		3		1		232	6.84
<u>Phanerodon furcatus</u>	3							3	0.09
<u>Pleuronichthys decurrens</u>	1	1				6		8	0.24
<u>P. verticalis</u>	16	6	4	3	1	12	2	44	1.30
<u>Porichthys myriaster</u>	1			1				2	0.06
<u>P. notatus</u>		2	4	36	9			51	1.50
<u>Rathbunella hypoplecta</u>						5		5	0.15
<u>Scorpaena guttata</u>		6	4		1		5	16	0.47
<u>Sebastes dalli</u>				1				1	0.03
<u>S. jordani</u>				1		9		10	0.29
<u>S. saxicola</u>		3	5	2		6	4	20	0.59
<u>S. semicinctus</u>		5				2		7	0.21
<u>Symphurus atricauda</u>	7	43	27	13	48	3	8	149	4.39

TABLE 9 (Cont.)  
 SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY  
 TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch
Trawl Date: 2/26/75 (Cont)									
<u>Synodus lucioceps</u>	3	1		2				6	0.18
<u>Zalembeus rosaceus</u>		2	5	3		35	7	52	1.15
<u>Zaniolepis frenata</u>				6		15	1	22	0.65
<u>Z. latipinnis</u>		23	4	4	1	1	5	38	1.12
Number of Individuals	86	809	395	545	246	652	860	3393	
Percent	2.5	23.8	11.6	10.2	7.3	19.2	25.3		
Trawl Date: 5/12/75									
<u>Anoplopoma fimbria</u>						3		3	0.07
<u>Caulolatilus princeps</u>							1	1	0.02
<u>Chilara taylori</u>		1		2	2	4		9	0.22
<u>Chitonotus pugetensis</u>		21	23	9		17	33	103	2.48
<u>Citharichtys fragilis</u>				12		1		13	0.31
<u>C. sordidus</u>		57	12	314		38	48	469	11.28
<u>C. stigmaeus</u>	15	113	184		88	26	28	454	10.92
<u>C. xanthostigma</u>			2		1		1	4	0.10
<u>Cymatogaster aggregata</u>					2			2	0.05
<u>Embiotoca jacksoni</u>	2							2	0.05
<u>Engraulis mordax</u>	2					4		6	0.14
<u>Genyonemus lineatus</u>	42		109	1	4	204	9	369	8.88
<u>Glyptocephalus zachirus</u>				7				7	0.17
<u>Hippoglossina stomata</u>			20		1	2	6	29	0.70
<u>Icelinus quadriseriatus</u>		52	264	45	1	49	169	580	13.95
<u>Lepidogobius lepidus</u>		2	1				1	4	0.10
<u>Lycodopsis pacifica</u>				66		6		72	1.73
<u>Lyopsetta exilis</u>				45				45	1.08
<u>Microstomus pacificus</u>	1	83	17	160	3	31	2	297	7.14
<u>Odontopyxis trispinosa</u>			2	2		2	2	8	0.19
<u>Otophidium scrippsae</u>	2							2	0.05
<u>Paralichthys californicus</u>	2					1		3	0.07
<u>Parophrys vetulus</u>		8	2	4	4	2	9	29	0.70
<u>Phanerodon furcatus</u>	1							1	0.02

TABLE 9 (Cont.)  
 SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY  
 TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch
Trawl Date: 5/12/75 (Cont)									
<u>Pleuronichthys decurrens</u>				1				1	0.02
<u>P. verticalis</u>	2		19	3	10		3	37	0.89
<u>Porichthys notatus</u>		9	9	157	1	7	3	186	4.47
<u>Scorpaena guttata</u>						3	5	8	0.19
<u>Sebastes crameri</u>				1				1	0.02
<u>S. dalli</u>			1	1				2	0.05
<u>S. goodei</u>	1		3			4	15	23	0.55
<u>S. jordani</u>	1		1	1		2	3	8	0.19
<u>S. levis</u>				10		1		11	0.26
<u>S. miniatus</u>	2	2	23		3		8	38	0.91
<u>S. rosenblatti</u>		1						1	0.02
<u>S. rubrivinctus</u>						1		1	0.02
<u>S. saxicola</u>	1	296	111	282	3	187	74	954	22.95
<u>S. semicinctus</u>		5		6		14		25	0.60
<u>Seriphus politus</u>						1		1	0.02
<u>Symphurus atricauda</u>	9	36	35	14	39		6	139	3.34
<u>Zalembeus rosaceus</u>		5	89	6		16	13	129	3.10
<u>Zaniolepis frenata</u>				11		1		12	0.29
<u>Z. latipinnis</u>		21	13	22		6	6	68	1.63
Number of Individuals	83	712	940	1182	162	633	445	4157	
Trawl Date: 7/9/75									
<u>Anoplopoma fimbria</u>		1		1				2	0.04
<u>Chilara taylori</u>		4	1	2				7	0.15
<u>Chitonotus pugetensis</u>		7	4	7		21	24	63	1.31
<u>Citharichthys sordidus</u>		89	16	258		165	173	701	14.35
<u>C. stigmaeus</u>	22	265	102		67	72	74	602	12.48
<u>C. xanthostigma</u>							1	1	0.02
<u>Coryphopterus nicholsi</u>							1	1	0.02
<u>Cymatogaster aggregata</u>	2		4		27			33	0.68
<u>Embiotoca jacksoni</u>	3							3	0.06
<u>Engraulis mordax</u>	34					2		36	0.75

TABLE 9 (Cont.)  
 SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY  
 TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch
Trawl Date: 7/9/75 (Cont)									
<u>Genyonemus lineatus</u>	126		106		42	12	102	388	8.04
<u>Hippoglossina stomata</u>		1	3	3		2	1	10	0.21
<u>Hydrolagus colliei</u>							2	2	0.04
<u>Hyperprosopon argenteum</u>	1							1	0.02
<u>Hyperprosopon argenteum</u>	1							1	0.02
<u>Icelinus quadriseriatus</u>		25	149	111		117	54	456	9.45
<u>Lepidogobius lepidus</u>							1	1	0.02
<u>Lycodopsis pacifica</u>				9				9	0.19
<u>Microstomus pacificus</u>	1	45	8	42	1			97	2.01
<u>Odontopyxis trispinosa</u>		2	3	3		4		12	0.25
<u>Otophidium scrippsae</u>	1							1	0.02
<u>Paralabrax nebulifer</u>			1					1	0.02
<u>Paralichthys californicus</u>	4						1	5	0.10
<u>Parophrys vetulus</u>		29	1	7	21	3	3	64	1.33
<u>Phanerodon furcatus</u>	17		1		4			22	0.46
<u>Pleuronichthys verticalis</u>	13		12		6		1	32	0.66
<u>Porichthys notatus</u>		11	2	23			1	37	0.77
<u>Rathbunella sp. A</u>		1		1				2	0.04
<u>Scorpaena guttata</u>							1	1	0.02
<u>Sebastes dalli</u>	67	744	289	219	96	212	349	1976	40.96
<u>S. goodei</u>				1				1	0.02
<u>S. hopkinsi</u>					1			1	0.02
<u>S. miniatus</u>			1				1	2	0.04
<u>S. rosenblatti</u>				1				1	0.02
<u>S. saxicola</u>		16		3		1		20	0.41
<u>S. semicinctus</u>				23				23	0.48
<u>S. umbrosus</u>		2						2	0.04
<u>Seriphus politus</u>	2		1					3	0.06
<u>Symphurus atricauda</u>	22	45	27	15	20	1	5	135	2.80
<u>Syngnathus californiensis</u>	1							1	0.02
<u>Synodus lucioceps</u>						1	1	2	0.04

TABLE 9 (Cont.)  
 SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY  
 TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch
Trawl Date: 7/9/75 (Cont)									
<u>Torpedo californica</u>				1				1	0.02
<u>Zalembius rosaceus</u>			10	8				18	0.37
<u>Zaniolepis frenata</u>				1				1	0.02
<u>Z. latipinnis</u>		23		24				47	0.97
Number of Individuals	316	1310	741	763	285	613	796	4824	
Percent	6.6	27.2	15.4	15.8	5.9	12.7	16.5		
Trawl Date: 10/8/75									
<u>Chilara taylori</u>		4			**	*	4	8	0.19
<u>Chitonotus pugetensis</u>		45	10	18			30	185	4.41
<u>Citharichthys sordidus</u>		63		336		54	219	903	21.56
<u>C. stigmaeus</u>	8	182	210	58	68	177	27	700	16.71
<u>Engraulis mordax</u>	71							71	1.69
<u>Genyonemus lineatus</u>	250			1		2		254	6.06
<u>Hippoglossina stomata</u>		5	1				2	8	0.38
<u>Hyperprosopon argenteum</u>	6							6	0.14
<u>Icelinus quadriseriatus</u>		186	59	399		12	209	100	23.04
<u>Lepidogobius lepidus</u>				1		2		3	0.07
<u>Microstomus pacificus</u>		28		8				2	38
<u>Odontopyxia trispinosa</u>		1	6	4		11	5	7	34
<u>Paralichthys californicus</u>	3							3	0.07
<u>Parophrys vetulus</u>		16	1	8	3	7		1	33
<u>Phanerodon furcatus</u>	5					3		8	0.19
<u>Pleuronichthys decurrens</u>							2	4	6
<u>P. verticalis</u>	17	1	3		4	6	2	2	31
<u>Porichthys notatus</u>		19	5	11		2	1	38	0.90
<u>Scorpaena guttata</u>			1			1	1	5	8

\* Retrawled station nT4 (10-13-75) numbers are included in total and % total catch because large piece of wire may have fouled trawl and resulted in low numbers on 10-8-75 that was not representative of station.

\*\*Station nT4 (10-8-75) numbers not represented in total or % total catch.

TABLE 9 (Cont.)  
 SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY  
 TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch	
Trawl Date: 10/8/75 (Cont)										
<u>Sebastes crameri</u>		1						1	0.02	
<u>S. dalli</u>		79	1	72		1	120	103	376	8.98
<u>S. jordani</u>		1							1	0.02
<u>S. miniatus</u>		10							10	0.24
<u>S. rubrivinctus</u>						1			1	0.02
<u>S. saxicola</u>		25							25	0.60
<u>S. semicinctus</u>		6							6	0.14
<u>S. (Sebastomus) sp. UI</u>						1			1	0.02
<u>Seriphus politus</u>	31								31	0.74
<u>Symphurus atricauda</u>	35	41	45	27		94	3	46	291	6.95
<u>Syngnathus californiensis</u>						1	1		2	0.05
<u>S. exilus</u>					2					
<u>Xeneretmus triacanthus</u>		2							2	0.05
<u>Zalembeus rosaceus</u>		14		3		14		10	41	0.98
<u>Zaniolepis latipinnis</u>		55		16		14		5	90	2.15
Number of Individuals	426	784	342	962	77	401	628	645	4188	
Trawl Date: 1/14/76										
<u>Cheilotrema saturnum</u>	1								1	0.03
<u>Chilara taylori</u>		1		1					2	0.06
<u>Chitonotus pugetensis</u>		5	17	2		1	9	18	52	1.50
<u>Citharichthys fragilis</u>				2					2	0.06
<u>C. sordidus</u>		45	18	328		49	82	104	626	18.11
<u>C. stigmaeus</u>	11	114	100	1	111	4	34		375	10.85
<u>C. xanthostigma</u>			8		3	1			12	0.35
<u>Cymatogaster aggregata</u>		18		4					22	0.64
<u>Engraulis mordax</u>	1								1	0.03
<u>Genyonemus lineatus</u>	45			36					81	2.34
<u>Hippoglossina stomata</u>		1	1	1		4		6	13	0.38
<u>Hyperprosopon argenteum</u>	1								1	0.03
<u>Hypsopsetta guttulata</u>	1								1	0.03
<u>Icelinus quadriseriatus</u>		95	22	239		345	215		916	26.50
<u>Lepidogobius lepidus</u>			1			1			2	0.0



TABLE 9 (Cont.)  
 SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY  
 TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch
Trawl Date: 1/14/76 (Cont)									
<u>Lycodopsis pacifica</u>						1		1	0.03
<u>Lyopsetta exilis</u>						1		1	0.03
<u>Menticirrhus undulatus</u>	3							3	0.09
<u>Microstomus pacificus</u>		17		2		2		21	0.61
<u>Odontopyxis trispinosa</u>		1		2		1		4	0.12
<u>Paralichthys californicus</u>	4		1	1				6	0.17
<u>Parophrys vetulus</u>		22		16	6		1	45	1.30
<u>Phanerodon furcatus</u>	7							7	0.20
<u>Pleuronichthys verticalis</u>	9	1	8	9	3		4	34	0.98
<u>Porichthys myriaster</u>	1							1	0.03
<u>P. notatus</u>				183		1		184	5.32
<u>Rhacochilus vacca</u>		1						1	0.03
<u>Scorpaena guttata</u>		209	37	126		51	53	476	13.77
<u>Sebastes saxicola</u>		5				5		10	0.29
<u>S. semicinctus</u>		5						5	0.14
<u>Seriphus politus</u>	35			28				63	1.82
<u>Symphurus atricauda</u>		70	25	20	47	38	10	210	6.07
<u>Syngnathus californiensis</u>						1		1	0.03
<u>Zalemnius rosaceus</u>		192				2		194	5.61
<u>Zaniolepis frenata</u>						1		1	0.03
<u>Z. latipinnis</u>		40	2	24		2	5	73	2.11
Number of Individuals	119	842	243	1026	221	553	453	3457	
Percent	3.4	24.4	7.0	29.7	6.4	16.0	13.1		

Trawl Date: 4/7/76									
<u>Argentia sialis</u>						1		1	0.04
<u>Chilara taylori</u>				1				1	0.04
<u>Chitonotus pugetensis</u>			11	3		4	23	41	1.82
<u>Citharichthys sordidus</u>		14	26	279	11	14	46	390	17.33
<u>C. stigmaeus</u>	10	142	82	5	40	3	10	292	12.97
<u>C. xanthostigma</u>			7		2			9	0.40
<u>Cymatogaster aggregata</u>	4				3		11	18	0.80

TABLE 9 (Cont.)  
 SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY  
 TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch
Trawl Date: 4/7/76 (Cont)									
<u>Embiotoca jacksoni</u>	3							3	0.13
<u>Genyonemus lineatus</u>	63			40	1		1	105	4.67
<u>Hippoglossina stomata</u>		2	4	1			6	13	0.58
<u>Icelinus quadriseriatus</u>	2	8	99	47	1	67	103	327	14.53
<u>Microstomus pacificus</u>		215	3	12		3	2	235	10.44
<u>Odontopyxis trispinosa</u>			2	1				3	0.13
<u>Paralichthys californicus</u>	3							3	0.13
<u>Parophrys vetulus</u>		64	7	11	7	3	2	94	4.18
<u>Phanerodon furcatus</u>					2			2	0.09
<u>Pleuronichthys verticalis</u>	8		9	3	6	1		27	1.20
<u>Porichthys notatus</u>		1		10	4	1		17	0.76
<u>Rhacochilus vacca</u>					1	1		2	0.09
<u>Scorpaena guttata</u>		1	1			1	5	8	0.36
<u>Sebastes dalli</u>		44	15	209		128	7	403	17.91
<u>S. miniatus</u>						1		1	0.04
<u>S. saxicola</u>			6			15	7	28	1.24
<u>S. semicinctus</u>		1		2				3	0.13
<u>Sebastes sp.</u>					1			1	0.04
<u>Seriphus politus</u>	5							5	0.22
<u>Symphurus atricauda</u>	10	5	27	6	11	8	9	76	3.38
<u>Syngnathus californiensis</u>	1			1				2	0.09
<u>Zalembeius rosaceus</u>		27	1	8	4	18	15	73	3.24
<u>Zaniolepis latipinnis</u>		13		19		6	29	67	2.98
Number of Individuals	109	537	300	658	94	275	277	2250	
Percent	4.8	23.9	13.3	29.2	4.2	12.2	12.3		
Trawl Date: 7/28/76									
<u>Agonopsis sterletus</u>							1	1	0.03
<u>Argentina sialis</u>				1				1	0.03
<u>Chilara taylori</u>		1		4				5	0.16
<u>Chitonotus pugetensis</u>		19	15	6		6	15	61	1.93

TABLE 9 (Cont.)

SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch
Trawl Date: 7/28/76 (Cont)									
<u>Citharichthys fragilis</u>				3				3	0.09
<u>C. sordidus</u>		57	2	136		82	104	381	12.06
<u>C. stigmaeus</u>	2	83	102		67		67	321	10.16
<u>Coryphopterus nicholsi</u>				1				1	0.03
<u>Cymatogaster aggregata</u>	10							10	0.32
<u>Engraulis mordax</u>	79							79	2.50
<u>Genyonemus lineatus</u>	692		103					795	25.17
<u>Hippoglossina stomata</u>			5	3	1	1	6	16	0.51
<u>Hyperprosopon argenteum</u>	14							14	0.44
<u>Icelinus quadriseriatus</u>		136	152	86		24	15	413	13.07
<u>Icichthys lockingtoni</u>		1						1	0.03
<u>Lepidogobius lepidus</u>			2					2	0.06
<u>Menticirrhus undulatus</u>	1							1	0.03
<u>Microstomus pacificus</u>		26	2	11		1		40	1.27
<u>Odontopyxis trispinosa</u>		5	3	3		2	4	17	0.54
<u>Paralichthys californicus</u>	6							6	0.19
<u>Parophrys vetulus</u>		7	1	16	2	2	2	30	0.95
<u>Peprilus simillimus</u>		1	1					2	0.06
<u>Phanerodon furcatus</u>	20							20	0.63
<u>Pleuronichthys decurrens</u>				1				1	0.03
<u>P. verticalis</u>	7		1	6	10		1	25	0.79
<u>Porichthys myriaster</u>	1				1			2	0.06
<u>P. notatus</u>		66		102		4	2	174	5.51
<u>Sebastes dalli</u>		25		349				374	11.84
<u>S. diploproa</u>		1						1	0.03
<u>S. goodei</u>				1				1	0.03
<u>S. rubrivinctus</u>				1				1	0.03
<u>S. saxicola</u>		4		1				5	0.16
<u>S. semicinctus</u>				42				42	1.33
<u>S. ?paucispinis</u>						1		1	0.03
<u>Seriphus politus</u>	98		1					99	3.13

TABLE 9 (Cont.)  
 SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY  
 TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch
Trawl Date: 7/28/76 (Cont)									
<u>Symphurus atricauda</u>	34	30	1	10	12	1	21	109	3.45
<u>Synodus lucioceps</u>					3		1	4	0.13
<u>Zalembeus rosareus</u>		11		3			3	17	0.54
<u>Zaniolepis frenata</u>				1				1	0.03
<u>Z. latipinnis</u>		68		14				82	2.60
Number of Individuals	964	541	391	801	96	124	242	3159	
Percentage	30.5	17.1	12.4	25.4	3.0	3.9	7.7		
Trawl Date 10/6/76									
<u>Argentina sialis</u>							3	3	0.19
<u>Chilara taylori</u>		1					1	2	0.13
<u>Chitonotus pugetensis</u>		32		3			5	40	2.54
<u>Citharichthys sordidus</u>		84	2	286		188	29	589	37.37
<u>C. stigmaeus</u>	1	130	73		1	20	136	361	22.91
<u>C. xanthostigma</u>			5			1	11	17	1.08
<u>Embiotoca jacksoni</u>	10							10	1.08
<u>Genyonemus lineatus</u>					2			2	0.13
<u>Hippoglossina stomata</u>		4	4	1		4		13	0.82
<u>Icelinus quadriseriatus</u>		76		210		57	1	344	21.83
<u>Microstomus pacificus</u>		2		2				4	0.25
<u>Odontopyxis trispinosa</u>				2				2	0.13
<u>Oxylebius pictus</u>	1							1	0.06
<u>Parophrys vetulus</u>		50	2	4	7	2		65	4.12
<u>Phanerodon furcatus</u>	2							2	0.13
<u>Pleuronichthys verticalis</u>	1		5	4		5	2	17	1.08
<u>Porichthys notatus</u>		2	4			1		7	0.44
<u>Scorpaena guttata</u>			7	1				8	0.51
<u>Sebastes serranoides</u>	1							1	0.06
<u>Symphurus atricauda</u>		56	3	1		22		82	5.20
<u>Syngnathus exilis</u>							1	1	0.06

TABLE 9 (Cont.)  
 SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY  
 TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch
Trawl Date: 10/6/76 (Cont)									
<u>Synodus lucioceps</u>				1	1		1	3	0.19
<u>Xystreurus liolepis</u>						2		2	0.13
Number of Individuals	16	437	105	515	11	311	181	1576	
Percent	1.0	27.7	6.7	32.7	0.7	19.7	11.5		
Trawl Date: 1/5/77									
<u>Chitonotus pugetensis</u>		3		2		16		21	1.30
<u>Citharichthys fragilis</u>		4	4	1		13		22	1.36
<u>Citharichthys sordidus</u>		13		115		47	2	177	10.98
<u>C. stigmaeus</u>	14	92	175	16	37	5	111	450	27.92
<u>C. xanthostigma</u>			5	33			5	43	2.67
<u>Embiotoca jacksoni</u>							1	1	0.06
<u>Genyonemus lineatus</u>					1			1	0.06
<u>Hyppoglossina stomata</u>			4	1		4	2	11	0.68
<u>Icelinus tenuis</u>						1		1	0.06
<u>I. quadriseriatus</u>		227	4	121		308	1	661	41.00
<u>Lepidogobius lepidus</u>						1		1	0.06
<u>Lyopsetta exilis</u>							1	1	0.06
<u>Odontopyxis trispinosa</u>				1		1		2	0.12
<u>Paralichthys californicus</u>	3				2			5	0.31
<u>Parophrys vetulus</u>				5				5	0.31
<u>Pleuronichthys decurrens</u>						1		1	0.06
<u>P. verticalis</u>	2	2	9	2	1	2	2	20	1.24
<u>Porichthys notatus</u>		4	1	17		20		42	2.61
<u>Scorpaena guttata</u>		1		2		4		7	0.43
<u>Sebastes dalli</u>						3	4	7	0.43
<u>S. miniatus</u>							1	1	0.06
<u>S. rubrivinctus</u>							1	1	0.06
<u>S. vexillaris</u>						1		1	0.06
<u>Symphurus atricauda</u>	1	38	21	8	17	25	3	113	7.01
<u>Syngnathus exilis</u>					2		1	3	0.19

TABLE 9 (Cont.)  
 SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY  
 TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch
Trawl Date: 1/5/77 (Cont)									
<u>Synodus lucioceps</u>		2		5			1	8	0.50
<u>Xystreureys liolepis</u>			3		1		1	5	0.31
<u>Zalembeius rosaceus</u>						1		1	0.06
Number of Individuals	20	386	226	329	61	453	137	1612	
Trawl Date: 5/2/77									
<u>Chilara taylori</u>	1		1	1	1	4		8	0.32
<u>Chitonotus pugetensis</u>			8	2		6	4	20	0.79
<u>Citharichthys fragilis</u>		1	14	20		3	1	39	1.54
<u>C. sordidus</u>		31	2	286	4	22	19	364	14.40
<u>C. stigmaeus</u>	8	102	279	63	102	387	91	1032	40.84
<u>C. xanthostigma</u>			14	11	13	6	3	47	1.86
<u>Genyonemus lineatus</u>	7						4	11	0.44
<u>Hippoglossina stomata</u>				3		1	1	5	0.20
<u>Icelinus quadriseriatus</u>		12	67	127	29	64	4	303	11.99
<u>Lepidogobius lepidus</u>			1					1	0.04
<u>Microstomus pacificus</u>		7	1	29				37	1.46
<u>Odonotopyxis trispinosa</u>		1	1	4	1	2		9	0.36
<u>Paralichthys californicus</u>	1						1	2	0.08
<u>Parophrys vetulus</u>		29		15	4		4	52	2.06
<u>Phanerodon furcatus</u>	3							3	0.19
<u>Pleuronichthys decurrens</u>							1	1	0.04
<u>P. verticalis</u>	2		9	2	2			15	0.59
<u>Porichthys notatus</u>		12	1	275	20			308	12.19
<u>Scorpaena guttata</u>			4		1	6	1	12	0.47
<u>Scorpaenichthys marmoratus</u>					1			1	0.04
<u>Sebastes dalli</u>		5		48		1	1	55	2.18
<u>S. goodei</u>						1		1	0.04
<u>S. miniatus</u>			3		1			4	0.16
<u>S. saxicola</u>		7	21	2	4	5	14	53	2.08
<u>S. serranoides</u>						1		1	0.04

TABLE 9 (Cont.)

SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF ORANGE COUNTY AT EACH STATION DURING QUARTERLY TRAWLING SURVEYS, 1975-77 (FROM CSDOC, 1975, 1976, 1977)

Species	nT0	nT1	nT2	nT3	nT4	nT5	nT6	Total	Percent of Total Catch
Trawl Date: 5/2/77 (Cont)									
<u>Symphurus atricauda</u>	2	16	12	9	21	7		67	2.65
<u>Synodus lucioceps</u>	1	2	26				2	31	1.23
<u>Zalembeus rosaceus</u>			6	11	3		5	25	0.99
<u>Zaniolepis latipinnis</u>		6	1	11		2		20	0.79
Number of Individuals	25	231	471	919	207	518	156	2527	

TABLE 10

SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF HUNTINGTON BEACH AT EACH STATION DURING QUARTERLY TRAWLING SURVEYS, 1971-72 (FROM ENVIRONMENTAL QUALITY ANALYSTS, INC., AND MARINE BIOLOGICAL CONSULTANTS, INC., 1973a).

Species	October 1971										May 1972									
	A	B	C	D	E	F	G	H	O	Total	A	B	C	D	E	F	G	H	O	Total
<u>Galeorhinus zyopterus</u>																				
<u>Triakis semifasciata</u>																				
<u>Squalus acanthias</u>																				
<u>Platyrrhinoidis triseriata</u>																				
<u>Rhinobatos productus</u>								1		1										
<u>Torpedo californica</u>																		1		1
<u>Myliobatis californica</u>																				
<u>Anchoa compressa</u>	1							1		2										
<u>Engraulis mordax</u>									1	1										
<u>Porichthys myriaster</u>																				
<u>P. notatus</u>																				
<u>Otophidium scrippsae</u>																				
<u>O. taylori</u>																				
<u>Syngnathus californiensis</u>														1		1		2		4
<u>Paralabrax maculatofasciatus</u>																				
<u>P. nebulifer</u>																				
<u>Chelotrema saturnum</u>																				
<u>Cynoscion nobilis</u>														1						1
<u>Genyonemus lineatus</u>	19			3	2		2	20	14	60	1	9			10				18	38
<u>Menticirrhus undulatus</u>								2		2										
<u>Seriphus politus</u>	102			1	1			36	794	934	1				7					8
<u>Amphistichus argenteus</u>							34	1		35	1	2	1				12	7	1	24
<u>Cymatogaster aggregata</u>	13	1	40	24	65		21	1		165		9	4	2	12	3	61	3	23	17
<u>Embiotoca jacksoni</u>																				
<u>Hyperprosopon argenteum</u>	18	2		7			4	11	1	43					1		1			2
<u>Phanerodon furcatus</u>	23	4	33	35	50		5			150	4	11	8		38	1	6		40	108
<u>Rhacochilus toxotes</u>																				
<u>Zalembeus rosaceus</u>															2					2
<u>Heterostichus rostratus</u>																				
<u>Peprius similimus</u>								1		1										
<u>Scorpaena guttata</u>															2	1				3
<u>Leptocottus armatus</u>	2		1							3		1					1			2
<u>Citharichthys sordidus</u>					5		10	1		16					2					2
<u>C. stigmaeus</u>	26	1	33	3	93	9	8	1		174	20	29	31	46	70	20	114	14	28	372



TABLE 10 (Cont.)

SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF HUNTINGTON BEACH AT EACH STATION DURING QUARTERLY TRAWLING SURVEYS, 1971-72 (FROM ENVIRONMENTAL QUALITY ANALYSTS, INC., AND MARINE BIOLOGICAL CONSULTANTS, INC., 1973a).

Species	Stations										Stations										Grand Total
	A	B	C	D	E	F	G	H	O	Total	A	B	C	D	E	F	G	H	O	Total	
	October 1971										May 1972										
<u>Hippoglossina stomata</u>							3	2		5				1	4						5
<u>Paralichthys californicus</u>			1	1					2	4		1			2			1	1		5
<u>Hypsopsetta guttulata</u>																	1	1			2
<u>Microstomus pacificus</u>														2	2						4
<u>Parophrys vetulus</u>					1					1	2	2		7	6	4					21
<u>Pleuronichthys decurrens</u>																1	1	1			3
<u>P. verticalis</u>	3		3		11	1				18		2	2	4	8	2	1	1		1	20
<u>Symphurus atricauda</u>	1		1	1	7	1	1			12		3		5	11			1			20
Number of Individuals	208	8	112	75	235	11	94	75	809	1627	29	69	48	67	177	33	199	31	111	764	
Number of Species	10	4	7	8	9	3	13	10	3	19	6	10	7	7	15	8	10	9	6	22	
Diversity Index	1.69	1.12	1.48	1.62	1.47	0.83	2.64	2.08	0.30		1.53	2.53	1.68	1.97	2.82	2.35	1.61	2.75	2.05		
	September 1972										December 1972										
<u>Galeorhinus zyopterus</u>								1		1											1
<u>Triakis semifasciata</u>									2	2											2
<u>Squalus acanthias</u>		1								1											1
<u>Platyrhinoideis triseriata</u>							3	1		4	2							3			5
<u>Rhinobatos productus</u>	3		1				3	2		9		1									1
<u>Torpedo californica</u>																					1
<u>Myliobatis californica</u>														1							1
<u>Anchoa compressa</u>	3						3	16		22	3	2	2								7
<u>Engraulis mordax</u>	69	85		208	126	72	48	62	124	794	21	50	24		4				46	145	940
<u>Porichthys myriaster</u>					2					2											2
<u>P. notatus</u>								2		2											2

TABLE 10 (Cont.)

SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF HUNTINGTON BEACH AT EACH STATION DURING QUARTERLY TRAWLING SURVEYS, 1971-72 (FROM ENVIRONMENTAL QUALITY ANALYSTS, INC., AND MARINE BIOLOGICAL CONSULTANTS, INC., 1973a).

Species	September 1972										December 1972										Grand Total	
	A	B	C	D	E	F	G	H	O	Total	A	B	C	D	E	F	G	H	O	Total		
<u>Otophidium scrippsae</u>				1		13				14		3								3	17	
<u>O. taylori</u>					6					6											6	
<u>Syngnathus californiensis</u>								1		1					1	2				1	4	9
<u>Paralabrax maculatofasciatus</u>								2		2											2	
<u>P. nebulifer</u>																1					1	1
<u>Chellotrema saturnum</u>			1							1											1	
<u>Cynoscion nobilis</u>																					1	
<u>Genyonemus lineatus</u>	26	127	10	30	394	197	7	160	7	958	19	76	17		5					6	123	1179
<u>Menticirrhus undulatus</u>	1		3					2	4	10	2	1	2						3		8	20
<u>Seriphus politus</u>	73	137		203	116	217	126	84	30	986	136	349	68		3					88	644	2572
<u>Amphistichus argenteus</u>	6	4	10		1		41	34	1	97	31	1	7					20		1	60	216
<u>Cymatogaster aggregata</u>	8	11	12	9	43	16	62	39	4	204	1	4	6		7					3	21	507
<u>Embiotoca jacksoni</u>		4	10							14											14	
<u>Hyperprosopon argenteum</u>	19	16	2	11	10		21	13	4	96	6	3	44							16	69	210
<u>Phanerodon furcatus</u>	39	25	27	9	26		9	18	1	154	5	3	8	2	4					1	23	435
<u>Rhacochilus toxotes</u>		2								2											2	
<u>Zalemnius rosaceus</u>																					2	
<u>Heterostichus rostratus</u>								2		2		1		2						3	5	
<u>Peprilus similimus</u>	21	3	1			2	6	3	3	39											40	
<u>Scorpaena guttata</u>																					3	
<u>Leptocottus armatus</u>							2	10		12											17	
<u>Citharichthys sordidus</u>																					18	
<u>C. stigmaeus</u>	19	19	41	18	67	57	15	70		306		3	11	5	29	19			7	1	75	927
<u>Hippoglossina stomata</u>																					10	

TABLE 10 (Cont.)

SPECIES AND NUMBER OF FISHES CAPTURED OFFSHORE OF HUNTINGTON BEACH AT EACH STATION DURING QUARTERLY TRAWLING SURVEYS, 1971-72 (FROM ENVIRONMENTAL QUALITY ANALYSTS, INC., AND MARINE BIOLOGICAL CONSULTANTS, INC., 1973a).

Species	September 1972										December 1972										Grand	
	A	B	C	D	E	F	G	H	O	Total	A	B	C	D	E	F	G	H	O	Total	Total	Total
<u>Paralichthys californicus</u>			2		1		2	4		9				2				1		3	21	
<u>Hypsopsetta guttulata</u>							1	1		2		1						1		2	6	
<u>Microstomus pacificus</u>																					4	
<u>Parophrys vetulus</u>						1				1				2	2					4	27	
<u>Pleuronichthys decurrens</u>								1		1								1		1	5	
<u>P. verticalis</u>	1	2	2	5	14	39	1	3		67		1	2		13	1				17	122	
<u>Symphurus atricauda</u>	1	3	1	7	45	44				101	2	22	4	1	16	2				47	180	
Number of Individuals	289	439	123	501	851	658	355	532	174	3922	228	521	196	14	84	25		36	163	1267		
Number of Species	14	14	14	10	13	10	19	22	8	33	11	16	13	6	10	5		7	9	23		
Diversity Index	2.97	2.55	2.87	1.97	2.47	2.49	2.91	3.15	1.41		1.92	1.67	2.79	2.41	2.74	1.26		1.96	1.79			

APPENDIX III  
NAVIGATION AIDS  
AND  
MARINE TRAFFIC



DEPARTMENT OF TRANSPORTATION  
UNITED STATES COAST GUARD

MAILING ADDRESS:  
COMMANDER (oan)  
ELEVENTH COAST GUARD DISTRICT  
UNION BANK BLDG.  
400 OCEANGATE  
LONG BEACH, CA. 90822

*[Handwritten signatures and stamps]*

16575/PF  
Ser: oan 264-78  
AUG 08 1978

Mr. W. M. Marshall  
Division Production Manager  
Western Division  
Shell Oil Company  
P. O. Box 831  
Houston, Texas 77001

Dear Mr. Marshall:

Enclosed is the approved application for Private Aids to Navigation for the proposed platforms Ellen and Elly. Your attention is directed to the comment in Block 11.

Approval of this application does not constitute Coast Guard approval of the platforms or the proposed site. The Coast Guard's position on the proposal is that it is unacceptable without the institution of the shipping safety fairways as described in the Eleventh Coast Guard District Commander's letter of 31 March 1978 to the District Engineer, Los Angeles District, Corps of Engineers.


If I can be of any further assistance please contact me at (213) 590-2222.

Sincerely yours,

*[Handwritten signature of M. J. Danko]*

M. J. DANKO  
Lieutenant Commander, U. S. Coast Guard  
Chief, Aids to Navigation Branch  
By direction of the District Commander

Copy to:  
Corps of Engineers, Los Angeles District



# SHELL OIL COMPANY

P.O. BOX 831  
HOUSTON, TEXAS 77001

July 14, 1978

Department of Transportation  
United States Coast Guard  
Eleventh Coast Guard District  
400 Oceangate  
Long Beach, California 90822

Attention Chief, Aids to Navigation Branch

Attached is our revised "Application for Class I Private Aids to Navigation" for our planned platforms Ellen and Elly located in OCS waters about eight miles southwest of Huntington Beach, California.

This application has been revised to comply with the requirements outlined in your letter 16575/PF, Ser: oan 205-78. The eight lanterns, all of which will be located 40 feet above mean high water, will each produce at least 6,500 candela. The two two-mile fog signal emitters, mounted at the same elevation on opposite corners of the complex, are directional and are synchronized. The HALS 15 lighting system for the top of the derrick remains unchanged. The entire navigational aid system will be connected to the emergency stand-by generator buss.

If you have any questions or need additional information, please let me know.

Yours very truly,

For: W. M. Marshall  
Division Production Manager  
Western Division

BLF:DC

## Attachment

cc - United States Geological Survey  
Los Angeles, California

Governor's Office of Planning & Research  
Sacramento, California

Port of Long Beach  
Long Beach, California

State Lands Commission  
Sacramento, California





TREASURY DEPARTMENT U. S. COAST GUARD CG-4143 (5-62)		APPLICATION FOR CLASS 1 PRIVATE AIDS TO NAVIGATION ON ARTIFICIAL ISLANDS AND FIXED STRUCTURES (Please read instructions on reverse)		Form Approved Budget Bureau No. 48-R383	
1. NAME AND ADDRESS OF CORPORATION OR PERSON MAKING APPLICATION Shell Oil Company 1200 Milam St. Houston, Texas 77002 Attn; Mr. R.C. Visser			2. ACTION REQUESTED FOR PRIVATE AIDS TO NAVIGATION A. <input checked="" type="checkbox"/> ESTABLISH AND MAINTAIN      E. <input type="checkbox"/> DISCONTINUE B. <input type="checkbox"/> CHANGE OWNERSHIP                      F. DATE OF ACTION C. <input type="checkbox"/> CHANGE EQUIPMENT D. <input type="checkbox"/> MOVE		
3. POSITION					
A. GENERAL LOCALITY AND GRID AREA		B. LATITUDE		C. LONGITUDE	
OCS San Pedro Bay, Ca. Zone 6		33° 34' 56.2867"N		118° 07' 39.2990"	
D. BLOCK NUMBER	E. SIGN	F. LEASE NUMBER		G. WELL NUMBER	
261	Ellen and Elly	OCS-P-0300		N/A	
4. LIGHT					
A. CHARACTERISTICS		COLOR	WHITE <input type="checkbox"/>	B. NUMBER INSTALLED	
FLASH 0.4 SECONDS		RED <input type="checkbox"/>		9	
ECLIPSE 0.6 SECONDS				C. ILLUMINANT (Check)	
				<input checked="" type="checkbox"/> ELECTRICITY <input type="checkbox"/> GAS <input type="checkbox"/> OIL	
				<input type="checkbox"/> OTHER (Specify)	
D. HEIGHT ABOVE MEAN HIGH WATER	E. VOLTS	F. AMPERES	G. INSIDE DIAMETER		H. CANDLEPOWER (If known)
+40 (8)	120 (8)	4.17 (8)	LENS	GLOBE	6,700 (8) EFI
+236 (1)	120 (1)	8.34 (1)	250MM	G-40	14,000 (1)
5. FOG SIGNAL (Characteristic will be one two-second blast every twenty seconds)					
A. CLASS		B. MANUFACTURED BY		C. MODEL NUMBER	
<input checked="" type="checkbox"/> A (2-Mile)		Automatic Power, Inc. Houston, Texas		ELG-500/02 (2)	
<input type="checkbox"/> B (1/2-Mile)					
6. STRUCTURE					
A. COLOR		B. HEIGHT ABOVE MEAN HIGH WATER		C. DEPTH OF WATER BELOW MEAN LOW WATER	
To be determined		Platform +39 Derrick		+236      -258	
7. AUTHORIZED BY CORPS OF ENGINEERS, U. S. ARMY, PERMIT NO.					
8. PERSON IN DIRECT CHARGE OF AID					
A. NAME			C. ADDRESS		
Mr. R. F. Honer			196 S. Fir St.		
B. TELEPHONE NUMBER			Ventura, Calif. 93001		
805-648-2751 Home 805-642-1665					
9. The applicant agrees to save the Coast Guard harmless with respect to any claim or claims that may result arising from the alleged negligence of the operation of the approved aids.					
Attached to this application are:					
A. <input checked="" type="checkbox"/> LOCATION PLAT		B. <input checked="" type="checkbox"/> PRINT OF STRUCTURE		C. <input checked="" type="checkbox"/> AIDS TO NAVIGATION EQUIPMENT LIST	
D. <input type="checkbox"/> CERTIFICATE REQUIRED BY 33 CFR 67.10-1(4)					
DATE			SIGNATURE		
July 13, 1978			R. C. Visser		
			TITLE		
			Project Manager, Western Division Production		
FOR COAST GUARD USE					
10. FROM: Commander Eleventh Coast Guard District (oan)					
A. THE ACTION DESCRIBED ABOVE IS			B. NOTICE TO MARINERS		
<input checked="" type="checkbox"/> APPROVED			<input checked="" type="checkbox"/> WILL BE ISSUED		
<input checked="" type="checkbox"/> APPROVED SUBJECT TO THE COMMENTS IN BLOCK 11 ON REVERSE			<input type="checkbox"/> WILL NOT BE ISSUED		
C. CHARTS AFFECTED			D. NAME OF AID(S)		
18746 (C&GS 5142)			Ellen and Elly Structure		
18747 (C&GS 5142-SC)					
E. DATE		F. SIGNATURE (By direction in accordance with 33 CFR 67)			
24 July 1978		M. J. DANKO, LCDR, USCG			

25791 TREAS. CGHQ. WASH., D.C.

Block 11:

All obstruction lights shall be operated to flash in unison  
(33 CFR 67.05-5)





- ▷ ELG 500-02 FOGHORNS (BOTH AT SAME ELEV.)
- FA 250 LANTERNS (ALL AT SAME ELEV.)
- ☆ HALS-15, 15 MILE LIGHTING SYSTEM

DATUM: MLLW

ADJACENT PROPERTY OWNERS:

- ① STATE OF CALIFORNIA
- ② U. S. DEPT. OF THE INTERIOR, OCS

LOCATION OF NAVIGATION AIDS ON PROPOSED DRILLING PLATFORM AND PRODUCTION PLATFORM

ON OCS LEASE P-0300 SAN PEDRO BAY, CALIF.

APPLICATION BY SHELL OIL CO. OPERATOR OF THE BETA UNIT PARCELS COVERED BY LEASES OCS P-0296, 0300, 0301, 0306 AND UNLEASED TRACT 255

ZF-2790

EQUIPMENT LIST

for

PLATFORMS ELLY AND ELLEN

(OCS LEASE P-0300)

SAN PEDRO BAY, CALIFORNIA

QUANTITY

DESCRIPTION

- |   |  |
|---|--|
| 1 | CG-1000 Fog Signal inverter with remote control switch and two ELG-500/02 emitters. Blast characteristic is 2 sec. ON and 18 sec. OFF 120/240 VAC 60Hz power source required.  |
| 1 | SF-4000 Light Controller and Monitor with photocell 120 VAC, 60Hz power source required.   |
| 8 | Dual Ventilated FA-250 Lanterns with 120 volt AC, 500 watt lamp with mounting stand to operate as a master and standby system. Flash characteristic is 0.4 sec. ON, 0.6 sec. OFF, 7000 effective candelas, 120 VAC, 60Hz power source required.    |
| 1 | HALS 15, 15 Mile Derrick Light. Dual ventilated FA-250 lantern with mounting stand and both lanterns operating simultaneously 15,000 effective candelas. Flash characteristic 1.0 sec. ON, 2.0 sec. OFF, 220 volt AC, 60Hz, power source required. |

REFERENCE EQUIPMENT LAYOUT PER DRAWING L-1205-C

ASSEMBLY -- 10th session  
Agenda item 8(b)



IMCO

RESOLUTION A.379(X)  
adopted on 14 November 1977

Distr.  
GENERAL

A X/Res.379  
15 December 1977

Original: ENGLISH

RECEIVED  
U.S. COAST GUARD

JAN 11 1978

WASHINGTON, D.C.  
GAIA

ESTABLISHMENT OF SAFETY ZONES AND FAIRWAYS OR ROUTEING  
SYSTEMS IN OFF-SHORE EXPLORATION AREAS

THE ASSEMBLY,

NOTING Article 16(i) of the Convention on the Inter-Governmental  
Maritime Consultative Organization concerning the functions of the  
Assembly,

RECOGNIZING the need for ensuring unencumbered exploitation of sea-  
bed resources as well as safety at sea,

RECOGNIZING FURTHER that the congestion of navigable waters by off-  
shore platforms or other similar structures could result in ships  
colliding with such structures thereby causing loss of life, pollution  
of the marine environment and economic loss,

RECALLING Resolution A.340(IX) by which it adopted a Recommendation  
on Establishment of fairways through off-shore exploration areas,

NOTING that in accordance with Article 5 of the 1958 Convention on  
the Continental Shelf, Governments may establish safety zones, extending  
to a maximum distance of 500 m around continental shelf installations or  
other devices, which should be respected by ships of all nationalities,

BEING INFORMED of the frequent infringements of safety zones by  
ships,

HAVING CONSIDERED the Recommendation adopted by the Maritime Safety  
Committee at its thirty-sixth session,

RECOMMENDS that Governments:

- (a) ensure that the exploitation of sea-bed resources does not seriously obstruct sea approaches and shipping routes;
- (b) study the pattern of shipping traffic through off-shore resource exploration areas at an early stage so as to be able to assess potential interference with marine traffic passing close to or through such areas at all stages of exploitation;
- (c) where proliferation of oil installations or changes of traffic pattern warrants it, consider as appropriate the designation of safety zones around off-shore platforms and other similar structures or the establishment and charting of fairways or routeing systems through exploration areas,

URGES Governments:

- (a) to take all necessary steps to ensure that ships under their flags, unless specifically authorized, do not enter or pass through duly designated safety zones;
- (b) to promulgate by all appropriate means details of designated safety zones and established fairways or routeing systems, taking into account Resolution A.341(IX) on the Dissemination of Information, Charting and Manning of Drilling Rigs, Production Platforms and Other Similar Structures,

REVOKES Resolution A.340(IX).

APPENDIX IV

OFFSHORE STRUCTURE COLLISION DATA

TABLE 11

RAMMING INCIDENTS INVOLVING VESSELS IN EXCESS OF  
500 GROSS TONS AND ARTIFICIAL ISLANDS,  
GULF OF MEXICO      JULY 1, 1962 - JUNE 30, 1977

CASE	<u>DATE</u> TIME OF DAY	<u>VES. TYPE</u> G. TONNAGE	<u>VISIBILITY</u> WIND SPEED	PRINCIPAL CAUSE	<u>DAMAGE - STRUCTURE</u> <u>DAMAGE - VESSEL</u>
1.	<u>11/9/63</u> Night	<u>Cargo</u> 5 - 10,000	<u>8 mi</u> 30 kts	Failure to make proper allowance for leeway in high wind.	<u>300,000</u> 5,000
2.	<u>5/4/64</u> Day	<u>Tug &amp; Tow</u> 1 - 5,000	<u>2 mi</u> 20 kts	Personnel error - unlicensed/un- certified crew member - passed too close to windward.	<u>180,000</u> <1,000
3.	<u>10/4/65</u> Night	<u>Cargo</u> 5 - 10,000	<u>1 - 3 mi</u> 20 kts	Lack of proper look out in reduced visibility (fog).	<u>50,000</u> 191,000
4.	<u>12/15/65</u> Day	<u>Cargo</u> 1 - 5,000.	<u>Unknown</u> Unknown	Unknown/insufficient information.	<u>200,000</u> <1,000

TABLE 11 (Cont.)

CASE	DATE	VES. TYPE	VISIBILITY	PRINCIPAL CAUSE	DAMAGE - STRUCTURE
	TIME OF DAY	G. TONNAGE	WIND SPEED		DAMAGE - VESSEL
5.	<u>9/12/67</u> Night	<u>Unknown</u> Unknown	<u>Unknown</u> Unknown	Damaged by collision with unidentified vessel.	<u>100,000</u> Unknown
6.	<u>10/30/67</u> Night	<u>Cargo</u> 10 - 15,000	<u>Poor in</u> <u>heavy rain</u> 45 kts	Use of uncorrected charts. Platform not identified on radar - sea return.	<u>1,100,000</u> 12,000
7.	<u>8/27/69</u> Night	<u>Cargo</u> 10 - 15,000	<u>18 mi</u> 5 kts	Failed to identify platform lights.	<u>500,000</u> 10,000
8.	<u>10/10/70</u> Night	<u>Tanker</u> >15,000	<u>5 mi</u> 7 kts	Lack of proper lookout.	<u>865,000</u> 60,000

TABLE 11 (Cont.)

<u>CASE</u>	<u>DATE</u> <u>TIME OF DAY</u>	<u>VES. TYPE</u> <u>G. TONNAGE</u>	<u>VISIBILITY</u> <u>WIND SPEED</u>	<u>PRINCIPAL</u> <u>CAUSE</u>	<u>DAMAGE - STRUCTURE</u> <u>DAMAGE - VESSEL</u>
9.	<u>May '74</u> <u>Night</u>	<u>Cargo</u> <u>1,5000</u>	<u>2+ miles</u> <u>slight</u>	Vessel operator carelessness/ inattention.	<u>850,000</u> <u>500,000</u>
10.	<u>August '75</u> <u>Night</u>	<u>Tanker</u> <u>&gt;15,000</u>	<u>2+ miles</u> <u>4 - 10 kts</u>	Failed to post lookout.	(total > <u>10,000,000</u> loss) > <u>10,000,000</u> (total loss)

IV-3

Source: References 3, 4.



APPENDIX V

OIL SPILL SIMULATION EQUATIONS

APPENDIX B

SIMULATED OIL SPILL EQUATIONS

Oil Spreading Equations

(1) gravity-inertia regime:

$$r_i(t) = K_{2i} (\Delta g V t^2)^{1/4} \text{ for } 0 \leq t \leq t_{iv}$$

$$\text{where } t_{iv} = (K_{2v}/K_{2i})^4 (V/\Delta g \nu_w)^{1/3}$$

and in particular,

$$r_i(t) = 1.192 V^{1/4} t^{1/2} \text{ for } 0 \leq t \leq 243.4 V^{1/3}$$

(2) gravity-viscous regime:

$$r_v(t) = K_{2v} (\Delta g V^2 t^{3/2} / \nu_w^{1/2})^{1/6} \text{ for } t_{iv} < t \leq t_{vt}$$

$$\text{where } t_{vt} = (K_{2v}/K_{2t})^2 (\rho_w/\sigma) (\Delta g \nu_w)^{1/3} V^{2/3}$$

and in particular,

$$r_v(t) = 4.709 V^{1/3} t^{1/4} \text{ for } 243.4 V^{1/3} < t \leq 175.2 V^{2/3}$$

(3) surface tension-viscous regime:

$$r_t(t) = K_{2t} (\sigma^2 t^3 / \rho_w \nu_w)^{1/4} \text{ for } t_{vt} < t \leq t_f$$

$$\text{where } t_f = ((10^5/\pi)^{1/2} (\sigma^2 / \rho_w \nu_w)^{-1/4} / K_{2t})^{4/3} V^{1/2}$$

and in particular,

$$r_t(t) = .35569 t^{3/4} \text{ for } 175.2 V^{2/3} < t \leq 3985 V^{1/2}$$

$r(t)$  = radius of slick in meters (m) at time  $t$  in seconds (sec)

$K_{2i} = 1.14$ ;  $K_{2v} = 1.45$ ;  $K_{2t} = 2.30$

$\rho_o$  = density of oil =  $900 \text{ Kg/m}^3$

$\rho_w$  = density of water =  $1025 \text{ Kg/m}^3$

$\Delta = \frac{\rho_w - \rho_o}{\rho_o} = 0.12195$  (dimensionless)

$g$  = acceleration of gravity =  $9.80665 \text{ m/sec}^2$

$V$  = initial spill volume ( $\text{m}^3$ )

$\nu_w$  = kinematic viscosity of water =  $1.04 \times 10^{-6} \text{ m}^2/\text{sec}$

$\sigma$  = spreading coefficient due to surface tension =  $.025 \text{ Kg/sec}^2 \text{ (N/m)}$

#### Vector Equations for Wind and Current Induced Drift

Calculate components of the wind velocity parallel and perpendicular to the surface current velocity:

$$W_p = W \cos \theta$$

$$W_v = W \sin \theta$$

where  $W$  = wind speed,

$W_p$  = speed component parallel to  $\vec{C}$ ,

$W_v$  = speed component perpendicular to  $\vec{C}$ ,

$\theta$  = angle between  $\vec{W}$  and  $\vec{C}$ ,

and  $\vec{C}$  = surface current velocity

If the wind is producing co-current drift, i.e.,  $W_p C > 0$ , and  $W_p > 20 C$  where  $C$  is the current speed, then a current drift-factor of 96.7% and a wind drift-factor of 3.3% can be used to give the total slick speed parallel to the surface current:

$$T_p = 0.967 C + 0.033 W_p$$

If the wind-induced drift is co-current ( $W_p C > 0$ ) but  $W_p \leq 20 C$ , then Tsahalis

showed that:

$$\begin{aligned}
T_p = C + C \left[ &-.00011 + 0.0015 \left(\frac{W_p}{C}\right) - 0.00229 \left(\frac{W_p}{C}\right)^2 \right. \\
&+ 0.0011 \left(\frac{W_p}{C}\right)^3 - 0.000102 \left(\frac{W_p}{C}\right)^4 \\
&+ 0.00000426 \left(\frac{W_p}{C}\right)^5 - (4.26 \times 10^{-6}) \left(\frac{W_p}{C}\right)^6 \\
&\left. + (6.78 \times 10^{-10}) \left(\frac{W_p}{C}\right)^7 \right]
\end{aligned}$$

If the wind is producing counter-current drift ( $W_p C < 0$ ) and if  $W_p < -35C$ , i.e.,  $W_p$  larger in absolute value than  $35C$ , the current and wind drift-factors of 96.7% and 3.3%, respectively, can be used:

$$T_p = 0.967 C + 0.033 W_p$$

If the wind-induced drift is counter-current ( $W_p C < 0$ ) but if  $W_p \geq 35C$ , i.e.,  $W_p$  is smaller in absolute value than  $35C$ , the total slick speed parallel to the surface current is:

$$\begin{aligned}
T_p = C + C \left[ &-.0027 + 0.0071 \left(\frac{W_p}{C}\right) - 0.00391 \left(\frac{W_p}{C}\right)^2 \right. \\
&- 0.000551 \left(\frac{W_p}{C}\right)^3 - 0.0000379 \left(\frac{W_p}{C}\right)^4 \\
&- 0.00000122 \left(\frac{W_p}{C}\right)^5 - (1.85 \times 10^{-8}) \left(\frac{W_p}{C}\right)^6 \\
&\left. - (1.06 \times 10^{-10}) \left(\frac{W_p}{C}\right)^7 \right]
\end{aligned}$$

In both the co-current and counter-current wind-induced drift cases, the total slick speed perpendicular to the surface current is given by a wind drift-factor of 3.3% applied to the wind-speed component perpendicular to the surface current:

$$T_v = 0.033 W_v$$

The resultant slick speed is then:

$T = \sqrt{T_p^2 + T_v^2}$ , with direction in degrees relative to the surface  
current direction:

$$\beta = \tan^{-1} \frac{T_v}{T_p}$$

---

References:

- Fay, James A. "Physical Processes in the Spread of Oil on a Water Surface". Proceedings of Joint Conference on Prevention and Control of Oil Spills. (1971), pp. 463-467.
- Premack, Joel and George A. Brown. "Prediction of Oil Slick Motions in Narragansett Bay". Proc Joint Conf. Prevention and Control of Oil Spills. (1973), pp. 531-540.
- Tsahalis, D. T. "Theoretical and Experimental Study of Wind and Wave Induced Drift". preprint from J. Phys. Ocean.

APPENDIX VI  
LIST OF COASTAL  
BEACHES AND PARKS

APPENDIX VI

COASTAL BEACH AND PARK FACILITIES  
(Long Beach to San Clemente, inclusive)

State

Bolsa Chica  
Huntington  
Doheny  
San Onofre  
San Clemente  
Corona del Mar (State owned, but maintained  
by Newport Beach)

County

Los Angeles  
Santa Ana River

Orange  
Sunset  
Sunset Aquatic  
Newport Harbor  
Laguna Niguel  
Dana Point Harbor  
Aliso Beach  
Dana Cove (maintained but not owned  
by County)

City

Huntington  
City Beach  
Huntington Harbor (joint jurisdiction -  
patrolled by County, maintained  
by City)

Laguna Beach  
Crescent Way Bay Beach  
Shaw Cove Beach  
Boat Canyon Beach  
Diver's Cove Beach  
Picnic Beach  
Rock Pile Beach  
Heisler Park  
Main Beach  
Sleepy Hollow Beach  
Saint Ann Beach  
Thalia Street Beach  
Oak Street Beach

Laguna Beach, continued

Mountain Road Beach  
Bluebird Street Beach  
Pearl Street Beach  
Wood's Cove Beach  
Moss Street Beach  
Victoria Street Beach  
\* Blue Lagoon  
\* Aliso Beach and Pier  
\* Camille  
\* Laguna Niguel Royal  
\* West Street  
\*\* Irvine Cove  
\*\* Emerald Bay  
\*\* Three Arch Bay

Long Beach

Alamitos Park  
Bixby Park  
Bluff Park  
Marine Park  
Overlook Park  
Bay Shore Beach (swim)  
Dana Place to 68th Place (swim and boat)  
Peninsula Bay Beach (swim and boat)  
Alamitos Bay (boat)  
Marine Stadium (motor boats)  
1st Place to 72nd Place (swim)  
Colorado Lagoon (swim)  
Long Beach Marina  
Long Beach Harbor  
Los Cerritos Channel  
San Gabriel River  
Los Angeles River Channel  
Shoreline Aquatic Park (proposed)

Newport Beach

Newport Pier  
Newport Beach  
Newport Dunes Aquatic Park  
Newport Bay Wildlife Preserve  
West Newport Park  
North Star Beach  
Bluffs Parks  
Cliff Drive View Park  
Balboa Pier  
Balboa Peninsula Park  
Balboa Island Beaches  
West Jetty Park  
Las Arenas Park  
Lido Park  
Lido Isle Beaches

\* Owned by County but guarded by Laguna Beach

\*\* Privately owned and maintained; guarded by City



San Clemente  
City Beach  
San Clemente Pier


Seal Beach  
Eisenhower Park  
Seal Beach Municipal Pier

Commercial Recreational Facilities Privately Run

Sportsfishing  
Seal Beach Sportsfishing  
Davey's Locker (Newport Beach)  
Art's Landing (Balboa)  
Dana Wharf (Dana Point)

Boating  
Alamitos Bay Yacht Club (Long Beach)  
Long Beach Yacht Club  
Basin Marina  
Newport Dunes Aquatic Park  
Art's Landing  
Embarcadero Marina (Dana Point)

APPENDIX VII  
UTILITY/SERVICE AGREEMENTS



POLICE DEPARTMENT  
City of Huntington Beach

P.O. BOX 70 • 2000 MAIN STREET, HUNTINGTON BEACH, CA. 92648 • TEL: (714) 536-5311

EARLE ROBITAILLE  
Chief of Police

September 20, 1978

WESTEC SERVICES, INC.  
Applied Sciences  
180 East Main Street  
Tustin, California 92680

Attention: Nina Gruver

Dear Ms. Gruver:

A number of your organizations telephoned us about ten days ago and a discussion was held regarding the project "Shell OCS Beta Unit Development". To reiterate that discussion, and answer the questions in your letter, we consider the police service we provide to be among the best and most technically advanced in the nation.

We do not anticipate any adverse effect your project will have upon our level of service under normal operations. There may be some concern regarding the protection of automobiles which may be parked for extended periods. Because of the transient population using the beaches in our city, there may be a temptation for auto theft, auto burglary, theft from autos or malicious mischief in your parking lots.

The traffic problem appears to be insignificant at this time. However, the one area of concern regarding our service is major disasters such as major oil leaks flooding our beaches, aircraft or vessel collisions, or accidents and personal accidents requiring immediate or emergency services.


It is difficult to project the additional personnel needed to service the area as a result of your project. It will definitely create a need for some additional police man hours, however, the number cannot be anticipated now.

The response time for emergency service averages under five minutes and for non-emergency calls, the average time is approximately twenty two minutes with the average overall time at below thirteen minutes.

We are pleased to respond to your request. If you have further questions, please contact us.

Sincerely,

EARLE W. ROBITAILLE  
Chief of Police

  
ROBERT E. FICKLE, Sergeant  
Special Operations Division

EWR:REF:skd

— Address all communications to the Chief of Police —

# City of Seal Beach

## FIRE DEPARTMENT

September 18, 1978

Nina Gruver  
Environmental Analyst  
Westec Services, Inc.  
180 East Main Street  
Tustin, CA 92680


Dear Ms. Gruver:

This is in response to your September 11, 1978 correspondence concerning a proposed Shell Beta Project and the use of the Shell Oil marine fueling facility for crew boats. We will answer your questions in the same order as presented.

1. We consider the current level of service for the community in relation to fire protection as "good."
2. We do not at this time contemplate any adverse effects upon the level of service should the project being researched be approved. We do, however, have one question, and that would be concerning any rupture of the underwater petroleum line between the proposed platforms and Pier J in the Port of Long Beach. A rupture of this pipeline could result in some problems relating to beach pollution, water pollution, and what could be combustible petroleum products in the water and on the sand.
3. We do not anticipate any additional personnel or cost to the Fire Department as a result of this project.
4. The only special services that we could forecast would be in relation to a rupture of the underwater pipe.
5. The Fire Department has an average response time of approximately five minutes to an emergency and five to ten minutes for non-emergency calls.


I hope that this response will assist you in preparation of the environmental impact necessary for the Shell Beta Project.

Yours truly,

  
R. E. Adams, Fire Chief  
SEAL BEACH FIRE DEPARTMENT

REA:mc

cc: City Manager  
Planning Department



# *The Port of Long Beach*

"America's most Modern Port"

P. O. BOX 570 • LONG BEACH, CALIFORNIA 90801 • TELEPHONES: (213) 437-0041 • (213) 775-3469 • TELEX: 65-6452 PORTOBEACH LGB

September 15, 1978

Ms. Nina Gruver  
Environmental Analyst  
Westec Services, Inc.  
180 East Main St.  
Tustin, Calif. 92680

Dear Ms. Gruver:

This is in reply to your letter of September 11, 1978, requesting written response to certain specified security aspects of the proposed Shell Beta Project. This will also confirm our responses to verbal inquiry tendered earlier.

Our current level of manned security services in the Port of Long Beach is considered to be good. Harbor Department civil service Security Officers maintain 24-hour days, every day in the year, continual harbor patrol. Security officers use black and white police type vehicles with two-way radio communication to command base. Security guard service for individual facilities, if required, must be furnished by the facility operator.

There is no presently recognized adverse effect the Shell Beta project might have on Port security service, and no additional port security personnel are contemplated as a result of the project.

No special security services on the part of the Port are anticipated. Project design and construction should provide any specific physical security measures considered appropriate.


PRESIDENT'S "E" AWARD  
FOR EXCELLENCE IN EXPORT



September 15, 1978  
Ms. Nina Gruver  
Page 2

Harbor Department Security force response time to an emergency averages two to five minutes, depending upon the nature of the emergency and its physical location relative to Port Security headquarters. Response time in non-emergency calls normally would vary from ten to thirty minutes-again depending upon the nature of the problem.

J. H. McJunkin  
General Manager

A handwritten signature in cursive script, appearing to read "H. H. Harnagel", written over the typed name.

H. H. Harnagel  
Director of Operations

HHH:njc



# PACIFIC HOSPITAL OF LONG BEACH

A NON-PROFIT CORPORATION

2776 PACIFIC AVENUE P.O. BOX 1268  
LONG BEACH, CALIF. 90801 (213) 595-1911

C. JOSEPH HEINZ  
Administrator

September 20, 1978


Ms. Nina Gruver  
Environmental Analyst  
Westec Services, Inc.  
Applied Sciences  
180 East Main Street  
Tustin, CA 92680

Dear Ms. Gruver:

We are in receipt of your letter of September 11th, outlining the proposed Shell Beta Project.

From the information submitted for our review, this product should have no adverse affect upon the level or quality of services provided by the Hospital, and we would be most willing to provide services if needed to employees and their families.

Sincerely,



C. Joseph Heinz

CJH:jl

LONG BEACH COMMUNITY HOSPITAL

OFFICE OF THE PRESIDENT

September 18, 1978


Ms. Nina Gruver  
Environmental Analyst  
Westec Services, Inc.  
Applied Sciences  
180 East Main Street  
Tustin, CA 92680

RE: Shell OCS Beta Unit Development

Dear Ms. Gruver:

In response to your September 11, 1978 letter, I cannot see that the proposed project would have any direct measurable effect on Long Beach Community Hospital.

Very truly yours,



Bruce R. Sanderson  
President

BRS/ep





# CITY OF LONG BEACH

## DEPARTMENT OF FIRE

400 WEST BROADWAY, ROOM 261

LONG BEACH, CALIFORNIA 90802

ADMINISTRATION  
436-2219

FIRE PREVENTION  
435-2458

FIRE TRAINING CENTER  
597-5488

TECHNICAL SERVICES  
599-3679

EMERGENCY PREPAREDNESS  
595-1751

September 15, 1978

Nina Gruver, Environmental Analyst  
Westec Services, Inc.  
Applied Sciences  
180 East Main Street  
Tustin, CA. 92680

Dear Ms. Gruver:

This is in reply to your letter of September 11, 1978 requesting information from our Department regarding our concerns about the Shell OCS Beta Unit development.

One of the problems in responding to your letter is that yours is one of several proposals involving the Harbor District. While individually a project may not adversely impact the level of service provided, collectively they are almost certain to do so.

However, we will predicate our response on the basis of no additional projects.

1. Our current level of service provided is excellent. Long Beach received a Class I rating for its Fire Department when rated in 1972 by the Insurance Services Office.
2. This project would not have significant impact on the level of service currently provided, but additional inspection responsibilities would be placed on our Fire Prevention Bureau.
3. No additional personnel would be required.
4. Special services required would consist of inspections by the Fire Prevention Bureau.


Shell OCS Beta Unit Development  
September 15, 1978

Page 2

5. Response time to an emergency call would be somewhere between three and four minutes, under normal conditions. Non-emergency response time would depend on the time of day and traffic conditions.

Should any additional information be required, please call me at (213) 436-2219.

Sincerely,



J. B. Souders, Deputy Chief - Administration  
Department of Fire

JBS:kd

APPENDIX VIII  
OCEANOGRAPHIC SURVEY RESULTS

# TABLE OF CONTENTS

	Page Number
TABLE OF CONTENTS . . . . .	VIII-1
LIST OF FIGURES . . . . .	VIII-3
LIST OF TABLES . . . . .	VIII-3
INTRODUCTION . . . . .	VIII-4
Objectives and Scope . . . . .	VIII-4
Characteristics of the Study Area . . . . .	VIII-4
EQUIPMENT AND PROCEDURES . . . . .	VIII-6
Study Areas and Sampling Locations . . . . .	VIII-6
Water Quality Measurements . . . . .	VIII-6
Current Measurements . . . . .	VIII-10
Current Meters . . . . .	VIII-10
Drogues . . . . .	VIII-10
Receiving Water Chemistry . . . . .	VIII-12
Nutrients . . . . .	VIII-12
Grease and Oil . . . . .	VIII-13
Trace Metals . . . . .	VIII-13
Coliform Bacteria . . . . .	VIII-13
Sediment Chemistry . . . . .	VIII-14
Grease and Oil . . . . .	VIII-14
Phenols . . . . .	VIII-14
Trace Metals . . . . .	VIII-15
Biochemical Oxygen Demand . . . . .	VIII-15
Data Reduction . . . . .	VIII-16
RESULTS . . . . .	VIII-16
Physical Oceanographic Measurements . . . . .	VIII-16
Temperature . . . . .	VIII-16
Salinity . . . . .	VIII-17
Density . . . . .	VIII-17
Dissolved Oxygen . . . . .	VIII-18
Hydrogen Ion Concentrations . . . . .	VIII-18
Light Transmittance . . . . .	VIII-18
Solar Irradiance . . . . .	VIII-19
Current Measurements . . . . .	VIII-19
Chemical Oceanographic Analyses . . . . .	VIII-38

	Page Number
Surface Receiving Waters . . . . .	VIII-38
Nutrients . . . . .	VIII-39
Grease and Oil . . . . .	VIII-40
Trace Metals . . . . .	VIII-40
Coliform Organisms . . . . .	VIII-40
Sediment Chemistry . . . . .	VIII-40
Organic Content . . . . .	VIII-42
Trace Metals . . . . .	VIII-42
Sediment Description . . . . .	VIII-44

## LIST OF FIGURES

		Page Number
Figure 1.	Shell Beta Study Area . . . . .	VIII-5
Figure 2.	Oceanographic Station Locations . . . . .	VIII-7
Figure 3.	Oceanographic Water Quality Data Acquisition System . . . . .	VIII-8
Figure 4.	Current Meter Mooring Arrangement. . . . .	VIII-11
Figure 5.	Biplane Drogue and Tethered Float . . . . .	VIII-12
Figure 6.	Shipek Bottom Grab Sampler. . . . .	VIII-14
Figure 7.	Surface, Mid-Depth, and Bottom Current Data for the Proposed Platform Site Station . . . . .	VIII-32
Figure 8.	Surface and Mid-Depth Current Data for Station P4 . . . . .	VIII-33
Figure 9.	Wind, Current, and Tide Data for Station T4 . . . . .	VIII-34
Figure 10.	Surface Drogue Tracks for Station S1 . . . . .	VIII-35
Figure 11.	Surface Drogue Tracks for Station T1 . . . . .	VIII-37

## LIST OF TABLES

Table 1.	Summary of Physical and Chemical Field Study Sampling Frequency . . . . .	VIII-9
Table 2.	Water Quality Profile Data . . . . .	VIII-20
Table 3.	Solar Irradiance Data . . . . .	VIII-30
Table 4.	Receiving Water Chemical Analysis Results (mg/l). . . . .	VIII-38
Table 5.	Receiving Water Examination for Coliform Organisms . . . . .	VIII-39
Table 6.	Sediment Chemical Analysis Results (mg/kg). . . . .	VIII-41

## INTRODUCTION

This report presents the results of a physical-chemical oceanographic field study conducted by the Environmental Sciences Division of Brown and Caldwell (B and C) in conjunction with the OCS Beta Unit EIR/EA for Shell Oil Company. This introductory section of the report describes the objectives and scope of the study, while the following sections present a description of the equipment and procedures used during the field effort, the data processing techniques, and the results of the survey.

### Objectives and Scope

The Shell Beta Unit Plan of Development envisions the construction of a 265 foot drilling platform (Ellen) and a 255 foot production platform (Elly). Future plans include a 700 foot drilling platform (Eureka) to produce the deep portions of the reservoir. A 16 inch pipeline will connect the production platform to a land site within Long Beach Harbor.

The estimated peak oil production rate from the 265 foot platform will be approximately 16,000 barrels per day (B/D) in 1982. The estimated peak oil rate from both the 265 and 700 foot platforms will be 26,000 B/D in 1984.

The purpose of this study was to gather supportive site specific data for the proposed Shell Beta platform site, the proposed pipeline corridor between the production platform and the Long Beach breakwater, and the proposed pipeline terminus corridor within Long Beach Harbor (Figure 1).

The primary objective of the study was to substantiate the validity of existing background information. A secondary objective of increasing the established data base was also accomplished. The scope of work included a single comprehensive examination of representative physical and chemical oceanographic parameters within the study area.

### Characteristics of the Study Area

The area of planned development is located within the Southern California Bight on the northeast shelf and slope of the San Pedro Basin, approximately nine miles offshore of Huntington Beach (Figure 1). The Beta accumulation (petroleum reservoir) is located at a structurally high position along the east side of the Palos Verdes Fault, extending along the fault approximately five miles. The beds dip generally to the northeast from 10 to 30 degrees. Oil-water contacts limit the width of the accumulation to approximately one mile.

Oil is currently produced from accumulations in folded structures associated with the Palos Verdes and Newport-Inglewood fault zones onshore and offshore in San Pedro Bay. There were no measureable quantities of  $H_2S$  in any of the




Figure 1. Shell Beta Study Area



produced gas samples from exploratory drilling. The sulfur content of the crude ranges from 3 to 4 percent with the estimated average produced gravity for both sites to be between 14 - 16° API.

## EQUIPMENT AND PROCEDURES

This section of the report describes sampling areas and sampling stations monitored, the survey equipment used, methodology employed, and the data reduction procedures followed during the Shell Beta field survey.

### Study Areas and Sampling Locations

Field surveys were conducted throughout the study area between July 18 and July 26, 1978. The Shell Beta study area was divided into three subareas as shown on Figure 1. The location of each of the 18 stations, including the positions from which drogues and current meters were deployed, are shown on Figure 2.

Table 1 presents a list of all parameters measured and the total number of samples taken at each site. Individual stations outside the breakwater were located using a Motorola miniranger electronic positioning system. Stations within the Long Beach breakwater were located by sextant and visual observation using landmarks and existing oil producing islands.

### Water Quality Measurements

Continuous vertical profile measurements of water quality parameters were measured during the July survey using Brown and Caldwell's water quality data acquisition system shown in Figure 3. The water quality data acquisition system consists of a Martek Mark III water quality analyzer and transmissometer and Brown and Caldwell's data processor. This system was used to record vertical profile measurements of temperature, conductivity, dissolved oxygen, pH, and light transmittance. As the underwater sensors were lowered from the surface to the bottom of each station, the analog signal produced by each sensor was converted to digital, displayed, and recorded at one-second intervals. The digital data processor recorded water quality profile information by printing the information on paper and also punching the information on mylar computer tape. The printed output was used as a hard copy for field check of the accuracy of parameters measured. The punched tape was used for computerized data reduction.

The survey procedure was the same at each water quality station. A label containing B and C's internal job number, date, station code, and time of the profile was recorded on punched tape by the digital data processor, and the station number was written directly on the printed output. The underwater sensors of the water quality data acquisition system were lowered over the side and held





Figure 2. Oceanographic Station Locations



OCEANOGRAPHIC WATER QUALITY DATA ACQUISITION SYSTEM

- |   |                                     |
|---|-------------------------------------|
| 1. Martek Mark III Underwater Sensors     | 4. Martek Transmissometer Deck Unit |
| 2. Martek XMS Transmissometer             | 5. Digital Data Processor           |
| 3. Martek Mark III Water Quality Analyzer | 6. Facit Paper Tape Punch           |

Parameter	Method of Measurement	Accuracy	Indicating and Recording Methods <sup>c</sup>
Depth	Pressure transducer	$\pm 1\%$ full scale, $\pm 1$ foot	Digital, printed paper tape, ASCII format mylar tape.
Temperature <sup>a</sup>	Transistor probe	$\pm 0.1\text{C}$ 0.01 C resolution	Digital, printed paper tape ASCII format mylar tape.
Specific electrical conductivity	Five-electrode guarded Kelvin cell	$\pm 0.02$ Millimho/cm	Digital, printed paper tape, ASCII format mylar tape.
pH	Glass and silver/silver chloride	$\pm 0.1$ pH unit, $\pm 0.01$ unit resolution	Digital, printed paper tape, ASCII format mylar tape.
Dissolved oxygen <sup>b</sup>	Auto-temperature compensated polarographic gold/silver electrode	$\pm 1\%$ full scale, $\pm 0.2$ part per million	Digital, printed paper tape ASCII format mylar tape.
Turbidity, light transmittance	Transmissometer, feedback-balanced photodiode detector	$\pm 1\%$ full scale	Digital, printed paper tape, ASCII format mylar tape.

<sup>a</sup> Temperature measuring systems are checked for accuracy with N.B.S. calibrated thermometers.

<sup>b</sup> Calibrated during the survey using the Winkler titration procedure.

<sup>c</sup> Accuracies listed are manufacturers specifications and are based on using analog meter outputs. Laboratory and field calibration indicates, however, that greater accuracy is achieved using digital recording outputs instead.

Table 1. Summary of Physical and Chemical Field Study Sampling Frequency

Parameters	PHYSICAL-CHEMICAL FIELD STUDIES					
	Receiving Water			Sediment		
	Site Area	Pipeline Area	Terminus Area	Site Area	Pipeline Area	Terminus Area
Temperature	10 <sup>a</sup>	24	24			
Salinity	10	24	24			
Dissolved Oxygen	10	24	24			
Hydrogen Ion Concentration	10	24				
Transmissivity	10	24	24			
Submarine Photometer	10	24	8			
Nitrate/Nitrite	3	7	8			
Phosphate	3	7	8			
Silicate	3	7	8			
Phenols				3		8
Coliform Bacteria	3					
Ammonia	3					
Grease and Oil	3	7	8	3	7	8
Cadmium				3		8
Copper				3		8
Lead				3		8
Silver				3		8
Zinc		7		3	7	8
Cobalt				3	3	8
Manganese				3		8
Arsenic				3		8
Boron				3		8
Nickel				3		8
Mercury				3		8
Total Chromium	3			3		8
Barium	3			3		8
Lead	3			3		8
Aluminum	3			3	4	3
Iron	3			3		8
BOD	3			3	7	8

<sup>a</sup> Numbers indicate total samples taken per area.

at the water's surface until they had equilibrated. The underwater sensors were then lowered manually by the electronic cable connecting the sensors to their readouts at approximately one-half foot per second as the digital data processor recorded all parameters simultaneously. After the sensors had been lowered all the way to the bottom (or to a depth of 280 feet) the digital data processor was turned off, and the sensors were retrieved. Each sensor was calibrated before and after each survey, and calibration checks were performed several times during each survey.

Light intensity was measured at all stations during daylight hours. A submarine photometer, measuring light within blue-green spectrum, was used to measure downwelling light intensity in milliamps at depths of surface, 6.5, 20, and 40 feet (surface, 2, 6, and 12 m). From these measurements, the percent of incident light remaining was calculated and converted to Langleys per second (ergs/cm<sup>2</sup>/sec).

## Current Measurements

The speed and direction of currents during the study period were determined by the use of both current meters and drogues.

Current Meters. Current measurements were taken at Stations S1 and P4 for a period of 25 hours. Three EG&G CT/3 electromagnetic current meters were moored using a dual string arrangement as shown in Figure 4. The meters were moored at near-surface (1 m depth), mid-depth, and within three meters of the bottom, moving vertically with the tide. Each assembly was moored with one danforth anchor on short scope to limit horizontal movement.

The EG&G meters are self-contained, in-situ recording instruments which orient themselves in the direction of the current flow by means of a large vane. Speed is detected using electromagnetic sensors. Direction is referenced to magnetic north by means of a flux gate compass. An internal program controlled by a crystal clock governs the rate of data recorded on a magnetic cassette tape recorder.

The meters have an accuracy of  $\pm 0.015$  m/sec in the speed range of 0.03 to 3.0 m/sec and a directional accuracy of  $\pm 5$  degrees. The three meters were programmed to record every six seconds over a two-minute period every fifteen minutes.

Current measurements within the Long Beach breakwater were obtained hourly at Station T4. A Bendix Model Q-15 ducted current meter was utilized to record both speed and direction at depths of 3, 6, 10, 15, 25, 30, and 40 feet. This current meter compensates for oscillatory wave motions so that only the net current speed is recorded. The current meter is oriented in the direction of the net current flow by a 10 foot long vane. Direction is referenced to magnetic north by a compass contained within the current meter housing. The ducted current meter system has an accuracy of  $\pm 0.03$  knots in the speed range of 0 to 1 knot and a directional accuracy of  $\pm 5$  degrees.

Drogues. Drogue releases were made during slack water between both flooding and ebbing tidal phases. The drogues were released twice at each station (S1, P4, and T4) and followed for a period of approximately six hours. Due to a positioning equipment malfunction, the data from Station P4 was determined unreliable and has not been included.

A typical drogue is shown in Figure 5. Each drogue consists of two 5' x 5' polyethylene panels attached at right angles to a frame of PVC pipe. The frame is rigid in the water but can be collapsed and rolled for ease of handling and storage. The drogue is attached to a surface float which in turn is tethered to a surface buoy. The drogue itself is weighted to maintain proper vertical orientation within five feet of the surface. The effective surface area ratios of drogue to marker buoy is approximately 100 to 1.




Figure 4. Current Meter Mooring Arrangement

## Receiving Water Chemistry

Receiving water samples were collected throughout the study area for subsequent laboratory analyses for nutrients, grease and oil, trace metals, and coliform organisms.

Nutrients. Two water samples were collected at each station and analyzed for ammonia, nitrate, nitrite, phosphate, and silica. The samples were taken from just below the water surface. One sample was filtered in the field through a  $0.45\mu$  membrane filter, then both the filtered and unfiltered samples were frozen to ensure preservation prior to laboratory analysis.

A Technicon Autoanalyzer was used to measure the ammonia by the automated Bertholet method (Standard Methods, 14th Edition, p. 616, 1975). Using this method, a blue-colored compound believed to be related to indophenol was formed when the sample was treated with sodium phenoxide, followed by the addition of sodium hypochlorite. Potassium sodium tartrate and sodium citrate were added to prevent the precipitation of




Figure 5. Biplane Drogue and Tethered Float

calcium and magnesium. Absorbance caused by the blue indophenol complex was measured at 630 nm and related to ammonia concentration by comparison with calibration standards.

Nitrite was measured using a Technicon Autoanalyzer by reaction with acidic sulfanilamide to form a diazo-complex. This complex was then coupled with N-1-naphthylenediamine dihydrochloride to form a reddish purple azo-dye which absorbs strongest at 550 nm. Standards and blanks were analyzed to relate absorbance to the nitrite concentration and correct for seawater matrix interference. This procedure is an adaptation of the nitrate nitrogen procedure (Standard Methods, 14th Edition, p. 620, 1975). In this case, the cadmium reduction column is eliminated.

Nitrates were reduced to nitrite using a cadmium reduction column (Standard Methods, 14th Edition, p. 620, 1975). Then total nitrite nitrogen (the sum of the original nitrite plus the reduce nitrate) was determined by the procedure described in the previous paragraph. The difference between this total nitrite nitrogen concentration and the nitrite nitrogen measured without the use of the reduction column provided the nitrate nitrogen concentration.

The automated determination of orthophosphate depends on the formation of a phosphomolybdenum blue-complex which absorbs strongly at 880 nm (Standard Methods, 14th Edition, p. 624, 1975). A reagent solution containing sulfuric acid, ammonium molybdate, ascorbic acid and antimony mixed with standards and samples and heated to 99.5 F (37.5 C) formed the colored complexes which allowed calculation of orthophosphate.

The automated procedure for the determination of soluble silicates is based on the reduction by ascorbic acid of an acidic silicomolybdate complex to molybdenum blue. Oxalic acid is introduced to prevent orthophosphate interference. The colored complexes produced from standards and samples were measured at 660 nm, allowing the calculation of the silicate concentrations (A Practical Handbook of Seawater Analysis, Strickland and Parsons, p. 65, 1968).

Grease and Oil. Water samples for grease and oil determinations were collected once at each station in 1-liter wide mouth glass containers and preserved by acidification. In the laboratory, dissolved or emulsified grease and oil was extracted from the water samples quantitatively with freon using the partition-gravimetric method (Standard Methods, 14th Edition, p. 515, 1975). Freon dissolves not only oil but also other organic substances. As no solvent can selectively dissolve only grease, the reported value is actually a measure of nonvolatile freon extractable material which is predominantly grease and oil. Each sample was transferred to a 2-liter separatory funnel, and the container was rinsed with 30 to 50 ml of freon, which was then transferred to the funnel. This mixture was shaken vigorously for several minutes and allowed to sit until layers of the solution separated. The organic layer containing grease was drained into a dried and tared boiling flask through freon moistened filter paper filled with anhydrous sodium sulfate. This extraction was repeated twice. The freon was then removed by vacuum evaporation, thus leaving the extracted material. The flask was dried at 221 F (105 C), cooled and weighed, and the freon extractable material was determined by weight difference.

Trace Metals. A one pint water sample was collected at each of the platform stations for laboratory analysis of total chromium, barium, lead, aluminum, and iron. The samples were fixed in the field with HNO<sub>3</sub>, cooled, and returned to the laboratory for analysis. All non-volatile metals samples were prepared for analysis by the digesting of 250 ml of sample to dryness after acidification with nitric acid. The sample was then redissolved into a matrix of nitric acid, hydrochloric acid and ammonium chloride, and filtered and diluted to 250 ml. The actual analyses by atomic absorption spectroscopy will be described in detail in the sediment metals section.

Coliform Bacteria. Receiving water was sampled for total coliform organisms at one foot below the surface at the three platform stations. Fermentation tubes containing sterile lactose broth were inoculated in the field with serial dilutions of 10, 1, 0.1, and 0.01 ml of sample and kept cool until returned to the laboratory for incubation at 95 F (35 C).



Tubes showing gas production at 24 or 48 hours were submitted to the confirmed test by aseptic transfer to fermentation tubes containing brilliant green bile lactose broth (BGB). Gas production within 48 hours of incubation at 95 F (35 C) in BGB constitutes a positive confirmed test for total coliform bacteria. Statistical tables allow determination of total coliform density as the most probable number (MPN) of organisms per 100 ml of sample.

### Sediment Chemistry

Sediment samples were collected at all the specified stations using a modified Shipek bottom sampler (Figure 6) and a one meter phlegar corer. The type of sampling device used depended primarily on the composition of the substratum being sampled. Approximately 14 ounces of sample were collected in glass jars for each station, the excess water was drained off, and the samples were cooled for transfer to the laboratory.

Analysis of sediment samples followed accepted procedures set forth by the Environmental Protection Agency, Chemistry Laboratory Manual, Bottom Sediments, Great Lakes Region Committee on Analytical Methods, 1969 (EPA, 1969). In the laboratory, portions of each sediment sample were dried at 22 F (105 C) to determine the percentage of solids so as to express all results on a dry weight basis.

Grease and Oil. Sediment grease and oil is determined by the soxhlet extraction method (EPA, 1969, p. 42). Samples for oil and grease determinations were acidified with hydrochloric acid and dried by mixing with anhydrous magnesium sulfate. Dried samples were transferred to cellulose thimbles and covered with glass beads to prevent the carrying over of solid sample into the extraction system. The freon soluble grease and oil was then determined gravimetrically using a soxhlet extraction apparatus. Following the four hour

extraction period, the solvent was removed from the tared boiling flasks using a rotary evaporator. The flasks were dried to constant weight at 22 F (105 C) and then weighed to determine weight-gain, allowing calculation of the grease and oil concentration. Blanks were run to check for impurities in the solvents used.




Figure 6. Shipek Bottom Grab Sampler

Phenols. Phenols, defined as the hydroxy derivatives of benzene were analyzed by the 4-aminoantipyrine colorimetric method (EPA, 1969, p. 77). Fifty gram samples were placed in one liter distillation flasks. Following addition of distilled water, copper sulfate and phosphoric acid, the samples were distilled to produce 500 mls of

aqueous phenols solution that was free of sediments. The phenolics were then treated at pH 10 with 4-aminoantipyrine in the presence of ferricyanide to form the colored antipyrine dye. This dye was concentrated by solvent extraction into chloroform and light absorbance measured at 460 nm. Standards and blanks were also distilled and analyzed to provide standard curves for the calculation of the initial phenolic contents of the sediments.

Trace Metals. All non-volatile metals were analyzed after the sediments had been suitably digested. Duplicate 15-gram aliquots of each sediment sample were digested to dryness in redistilled nitric acid and 20% hydrogen peroxide, mildly ashed and redissolved in a nitric acid-hydrochloric acid-ammonium chloride-calcium matrix. The supernatant and residue were filtered and washed into a volumetric flask and diluted to 250 ml. Each metal was analyzed by atomic absorption spectrometry (AAS) using a Jarrel-Ash Model 810 spectrometer. Background correction using a non-absorbing line or a hydrogen continuum lamp, was used to minimize error due to flame and sample matrix interferences. Matrix-matched standards were analyzed to provide suitable calibration curves. The procedures used were in accordance with EPA procedures (EPA, 1969). When concentrations approached the flame detection limits a flameless atomizer was used in place of the air-acetylene flame. In these cases, replicate analyses were performed and averages determined to minimize error.

Analysis of arsenic was also performed by flameless atomic absorption spectroscopy. Each sample was prepared in a 1000 ppm nickel matrix for the purpose of creating a nickel-arsenic alloy. This provided stabilization of the arsenic so that interferences could be removed through an ashing step prior to atomization.

Mercury concentration was determined from triplicate 1 to 3 g aliquots of each sediment sample which were mildly digested using nitric acid and potassium permanganate in an autoclave at 239 F (115 C) and 15 atmospheres of pressure for 15 minutes. Each digestate was treated with hydroxylamine hydrochloride to reduce any remaining permanganate and manganese dioxide to manganese ion. The amount of mercury in each aliquot was measured by means of the cold vapor AAS technique using a Jarrel-Ash trace metal accessory. Each sample was treated with acidic stannous chloride and stripped of the atomic mercury with air by means of an air stone. This air, containing traces of mercury was passed through a quartz cell where atomic absorption occurred. The peak area gas then related to the quantity of mercury present.

Biochemical Oxygen Demand. Tests for biochemical oxygen demand (BOD) were performed in accordance with interim procedures (EPA, 1969, p. 44), that are still in the process of being revised. This method commonly produces variable results that are dependent upon sample size. A procedural modification recommended by the laboratory group of EPA, Region IX, 1974, was used to lessen the BOD dependency on sample size. Approximately 5-g samples were mixed with 100 ml of dilution water, and the resultant slurry used as the sample. Aliquots of the slurry were transferred to 300-ml BOD bottles, and the bottles were filled

with dilution water (distilled water plus trace nutrients). The dissolved oxygen was measured in the bottles before and after a five-day, 68F (20 C) incubation. The BOD was then calculated as milligrams of oxygen uptake per kilogram of sediment.

### Data Reduction

Water quality data recorded during the field surveys on mylar punched tape were returned to the office and transferred to diskette using a tape reader and minicomputer. Data were then edited and processed by the minicomputer system, wherein salinity and density were computed from measured values of temperature and specific conductivity.

Current meter data recorded on magnetic cassette tapes were processed using a minicomputer. Oscillatory wave motion was filtered out and vectors were averaged to produce hourly values of current speed and direction.

Drogue positions were plotted from miniranger and sextant observations. Mean speed and direction were computed for each drogue release over the tracking period.

## RESULTS

This section of the report presents both a written and graphical representation of the results of the field effort. The physical oceanographic measurement results are presented first, followed by the results of the chemical oceanographic analyses. All physical and chemical parameters measured around the Shell Beta development and pipeline sites were typical of waters within the Southern California Bight and all concentrations and measurements were in agreement with background research data collected by other agencies within the area.

### Physical Oceanographic Measurements

Physical oceanographic measurements included sampling and data collection at 18 stations as shown on Figure 2. One measurement was taken at each of the three proposed platform sites, seven along the pipeline corridor between platform "Elly" and the Long Beach breakwater, and eight within the Long Beach harbor area.

Physical parameters measured during the field effort included: temperature, salinity, density, dissolved oxygen, hydrogen ion concentration (pH), light transmittance, solar irradiance, and current measurements.

Temperature. Temperature profiles were taken at each station during the study period to provide background data for the proposed construction and production operations. All profile data are presented in Table 2 at the end of this section.

Water temperatures at the platform sites ranged from 65.5 F (18.6 C) at the surface to 49.6 F (9.8 C) at a depth of 280 feet (85 m). A thermocline was observed above 80 feet (24 m) with the maximum temperature change (0.25 F/foot) occurring between 33 and 50 feet (10 and 15 m) at Station S2. Surface temperatures showed a diurnal trend with warmest temperatures occurring during late afternoon and coolest temperatures occurring during the early morning hours. Platform area surface temperatures varied 2 to 5 F (1 to 3 C) during the study period.

Temperatures along the proposed pipeline corridor varied between 61.0 and 69.8 F (16.1 and 21.0 C) at the surface and between 50.4 and 56.7 F (10.2 to 13.7 C) at the bottom. Bottom temperature variations were largely a function of fluctuating station depth, while both surface and bottom temperature ranges were influenced by spatial distribution. The observed thermocline was variable and also a function of station depth. Diurnal variation ranged from 2 to 4 F (1 to 2 C).

Temperatures within the harbor area varied between 64.6 and 70.2 F (18.1 and 21.2 C) at the surface and between 57.7 and 62.4 F (14.3 and 16.9 C) at the bottom. Recorded temperatures were generally warmer at Stations T1 and T2 and cooler at Stations T4 and T5.

Salinity. Salinity values during summer months normally exhibit relatively small variations except in the surface waters. Evaporation can cause an increase in surface salinity while rainfall and subsequent stormwater runoff can lower values. Salinity measured throughout the proposed development area was calculated from temperature and conductivity measurements and are presented in Table 2.

The proposed platform site area exhibited salinity values between 32.59 and 33.57 ppt. Lowest salinity values were observed below the surface between 50 and 65 feet (15 and 20 m) of depth, the same depth at which the observed maximum temperature change was observed. Salinity values along the proposed pipeline corridor varied between 32.41 and 34.19 ppt, while values within the harbor area varied between 32.52 and 33.49 ppt. Observed salinity values were slightly lower than average for the Pacific Ocean at local latitudes (34.24 ppt, SCCRWP, 1973), but were within the normal range for oceanic waters.

Density. Density, although not measured directly, was derived using temperature and salinity profile data. Density, an indicator of water column stability, has been reported as sigma-T units in Table 2.

Density values measured at the proposed platform site area varied between 23.72 and 24.08 at the surface and between 25.59 and 25.83 at the bottom. All density profile values increased with depth at all platform and pipeline stations. Values along the proposed pipeline corridor varied between 23.13 and 24.45 at the surface and between 25.06 and 25.88 at the bottom.

Within the harbor area, density values increased with depth at most stations. Several stations, however, exhibited weak density structure, probably

the result of turbulent mixing from strong winds and tidal currents. Density values varied between 22.23 and 23.75 for surface waters and between 24.17 and 24.90 for bottom waters.

Dissolved Oxygen. Dissolved oxygen (DO) concentrations were measured throughout the study area as an indicator of plankton productivity and chemical or biochemical oxygen demand. Generally, dissolved oxygen values were highest at the surface and decreased with depth. All DO values measured during the field study are presented in Table 2.

Concentrations of dissolved oxygen at the proposed platform site varied between 6.7 and 10.3 mg/l at the surface and between 3.5 and 6.3 mg/l at the bottom. Along the proposed pipeline corridor, values varied between 5.7 and 9.1 mg/l at the surface and between 3.3 and 8.9 mg/l at the bottom. Again, the difference observed among bottom DO values was partially due to fluctuating station depth. Values within the harbor area varied between 5.8 and 10.4 mg/l for surface waters and between 4.8 and 8.8 mg/l for bottom waters.

Surface dissolved oxygen concentrations were generally higher during daylight hours than during periods of darkness, while maximum DO values at night were observed 10 to 20 feet (3 to 6 m) below the water's surface. This observation is typical of conditions off southern California where plankton activity plays a dominant role in the spatial distribution of oxygen concentrations.

Hydrogen Ion Concentrations. Hydrogen ion concentrations in seawater are affected by numerous physical and chemical parameters including: temperature and salinity, photosynthesis and respiration, and carbon dioxide concentration and gas exchange in a carbonate/bicarbonate buffer system. Changes in phytoplankton populations are often reflected in pH concentrations. Results of hydrogen ion concentration data collected during the field study are presented as pH units in Table 2.

The pH within the Southern California Bight normally varies between 7.5 and 8.6, with an average of 8.1 (Hancock, 1965). Values measured at the proposed platform site and along the proposed pipeline corridor varied between 7.67 and 8.09, while values within the harbor area were slightly higher than normal, varying between 7.92 and 8.62. Similar high pH values were reported by Hancock (1974) for measurements within Long Beach Harbor.

Light Transmittance. Light transmittance, as a measure of turbidity, may be affected at the proposed platform site area by the discharge of drilling muds and cuttings and along the proposed pipeline corridor within the Long Beach Harbor by dredging operations during construction. Measurements collected during the field study are presented in Table 2 as percent light transmittance.

Light transmittance values at the proposed platform site were generally high, varying between 58 and 76 percent for surface waters. Bottom values usually approached zero percent transmittance prior to reaching the ocean floor.

Maximum values were generally observed just below the thermocline, beyond the productive surface layer. Along the proposed pipeline corridor, transmittance values were lower than those recorded for the platform site. Surface values varied between 04 and 76 percent and were progressively lower with decreasing distance from shore. Maximum values were recorded below the surface layer at most stations. Light transmittance within the harbor area varied between 07 and 23 percent with maximum values occurring most frequently at Stations T5 through T8 at the surface and at Stations T1 and T4 between 10 and 15 feet (3 and 5 m).

Solar Irradiance. Solar irradiance was measured at each station during daylight hours to record the amount of light penetration available for photosynthesis within the water column. Values were recorded at the surface and at 6.5, 20, and 40 feet (2, 6, and 12 m). The amount of initial surface solar irradiance is only important as a reference to calculate the percent of penetration at depth. The greater the percent penetration, the greater the amount of energy available for photosynthesis. Measurement results presented in Table 3 were recorded in the field as millamps then converted to Langeleys per second for the purpose of this report.

The greatest initial surface irradiance (8.9 Langeleys) was measured at Station P2 at 1513 on July 21 while the greatest percent penetration (88%) was measured at Station P1 at 1337 on July 19. The area around the proposed platform development exhibited the greatest percent irradiance penetration with average values ranging from 23 to 43 percent. At stations progressively closer to shore, sunlight penetration decreased dramatically along the proposed pipeline corridor. While the average sunlight penetration at Station P1 was 43 percent, the average near the Long Beach breakwater (P7) was only 04 percent. Although initial surface solar irradiance values recorded within the harbor area were similar to those measured at the platform site, percent penetration was dramatically reduced. The average percent penetration within the harbor area varied between < 1 and 02 percent.

Patterns observed through solar irradiance measurements coincide with the results of the light transmittance measurements, indicating the presence of an increasing turbidity field with decreased distance from shore. The source of turbidity in this area is most probably due to a combination of suspended sediment from wave action and upwelling, and the presence of phytoplankton in the near surface waters.

Current Measurements. Current measurements were determined using both current meters and current drogues.

Current meter measurements for speed and direction were collected at the proposed platform site, midway along the pipeline route (P4), and within the harbor area (T4) as shown on Figure 2. Measurements at the proposed platform site were taken midway between Stations S1 (Elly) and S2 (Ellen) using three electromagnetic current meters. The meters were moored at depths of 3, 120,

Table 2. Water Quality Profile Data

July 18, 1978

July 19, 1978

STATION	P2	P1	S1	P4	P7	P6	P5	P4	P3	P2	P1
TIME	1422	1527	1545	1620	911	1040	1152	1212	1246	1301	1337
DEPTH (METERS)	TEMPERATURE (C)										
0	19.82	20.11	19.66	19.72	20.98	20.13	20.42	20.42	18.41	18.69	18.29
5	18.53	19.46	18.97	19.51	19.72	19.80	19.93	19.84	18.37	18.23	18.30
10	16.43	16.15	16.08	17.90	16.06	17.33	16.65	16.52	16.95	17.00	18.14
15	14.66	13.45	13.88	14.99	13.49	14.35	14.38	14.71	14.11	14.82	16.84
20	12.86	12.68	12.99	13.70	13.36	13.52	13.46	13.46	12.72	13.12	14.42
25	12.38	12.07	12.09				13.46	13.30		12.10	12.52
30	12.23	11.60	11.42						11.92	11.66	11.90
35		11.39	11.14						11.57	11.26	11.25
40		11.05	10.83						11.19	11.06	11.09
45		10.98	10.65						11.01	10.97	11.01
50			10.61							10.76	10.93
55			10.46							10.69	10.78
60			10.25								10.51
65			10.21								10.37
70			10.18								10.29
75			10.15								10.25
80			10.15								10.23
85											10.22

SALINITY (PPT)											
0	33.15	33.31	33.23	33.21	32.94	33.19	33.08	33.18	33.22	31.11	33.38
5	32.80	32.97	32.96	33.16	32.81	33.00	32.98	32.83	33.19	33.02	33.24
10	33.30	33.30	33.22	33.48	33.12	32.67	33.26	33.32	32.97	33.20	33.25
15	33.60	34.19	33.47	33.28	33.37	33.23	33.55	33.81	33.06	33.11	33.20
20	33.43	33.94	33.42	33.26	33.64	33.46	33.25	33.42	33.08	33.19	33.27
25	33.45	33.77	33.58				33.22	33.38	33.18	33.45	33.58
30	33.39	33.60	33.52						33.18	33.51	33.59
35		33.61	33.41						33.21	33.52	33.46
40		33.54	33.39						33.31	33.51	33.69
45		33.43	33.42							33.33	33.72
50			33.47							33.35	33.67
55			33.47							33.42	33.69
60			33.55								33.58
65			33.55								33.61
70			33.58								33.56
75			33.57								33.57
80			33.56								33.58
85											33.56

DENSITY (SIGMAT)											
0	23.42	23.46	23.53	23.49	22.96	23.37	23.21	23.28	23.83	22.15	23.98
5	23.48	23.38	23.49	23.51	23.19	23.31	23.26	23.18	23.82	23.72	23.87
10	24.37	24.42	24.38	24.15	24.31	23.67	24.28	24.36	23.99	24.16	23.92
15	25.05	25.69	25.05	24.67	25.05	24.77	25.01	25.14	24.69	24.57	24.19
20	25.22	25.66	25.19	24.93	25.29	25.12	24.97	25.10	24.99	24.99	24.78
25	25.34	25.64	25.49				24.94	25.10	25.21	25.39	25.41
30	25.31	25.60	25.57						25.28	25.52	25.54
35		25.64	25.53						25.37	25.60	25.56
40		25.65	25.57						25.48	25.62	25.76
45		25.58	25.62							25.50	25.80
50			25.67							25.56	25.77
55			25.70							25.62	25.81
60			25.80								25.78
65			25.81								25.82
70			25.83								25.80
75			25.83								25.81
80			25.82								25.83
85											25.81

Table 2. Water Quality Profile Data (Cont'd)

July 18, 1978

July 19, 1978

STATION P2 TIME	P1 1422	P1 1527	S1 1545	P4 1620	P7 911	P6 1040	P5 1152	P4 1212	P3 1246	P2 1301	P1 1337
DISSOLVED OXYGEN (MG/L)											
0	8.59	7.08	6.84	6.93	9.13	7.01	7.13	7.04	6.82	6.80	6.84
5	8.56	7.07	6.85	7.08	9.03	7.10	7.12	7.02	6.93	6.80	6.89
10	9.01	7.31	7.14	7.18	9.02	7.11	7.27	7.23	6.97	6.93	6.92
15	8.94	7.50	7.42	7.50	9.10	7.28	7.25	7.33	8.45	7.14	6.85
20	18.26	7.38	7.27	7.49	8.94	6.99	7.01	7.23	8.00	7.12	6.99
25	8.74	7.15	6.96				7.00	7.16	7.51	6.79	6.87
30	8.47	6.81	6.54						6.97	6.44	6.61
35		6.52	6.04						6.40	6.01	6.16
40		6.17	5.52						6.11	5.51	5.73
45		6.05	5.10							5.23	5.40
50			4.75							4.91	5.10
55			4.39							4.70	4.83
60			3.99								4.53
65			3.84								4.24
70			3.71								3.98
75			3.62								3.73
80			3.61								3.61
85											3.59

HYDROGEN ION CONCENTRATION (PH)											
0	8.07	8.02	8.01	8.03	8.03	8.09	8.07	8.07	8.01	8.02	8.02
5	8.05	8.00	8.00	8.03	8.01	8.05	8.03	8.04	8.01	8.00	8.01
10	8.04	7.96	7.97	8.03	7.93	7.98	7.96	7.97	7.98	7.99	8.01
15	7.98	7.90	7.94	7.98	7.81	7.86	7.88	7.94	7.98	7.96	8.00
20	7.92	7.87	7.90	7.90	7.79	7.82	7.85	7.85	7.91	7.90	7.94
25	7.87	7.84	7.86				7.85	7.84	7.87	7.85	7.87
30	7.85	7.82	7.81						7.84	7.82	7.83
35		7.81	7.78						7.79	7.75	7.76
40		7.77	7.75						7.77	7.75	7.74
45		7.77	7.74							7.76	7.73
50			7.73							7.73	7.71
55			7.72							7.72	7.70
60			7.71								7.70
65			7.71								7.70
70			7.70								7.69
75			7.70								7.69
80			7.70								7.69
85											7.69

LIGHT TRANSMITTANCE (%)											
0	56	66	76	46	4	6	9	11	73	76	73
5	86	84	82	51	39	39	46	52	73	84	75
10	72	86	85	42	73	62	59	69	75	83	75
15	63	91	89	37	0	1	15	26	79	79	71
20	56	93	94	1	0	0	0	0	78	83	76
25	21	96	10				0	0	73	86	78
30	2	81	3						60	81	83
35		74	34						34	70	84
40		22	79						0	67	85
45		0	90							71	85
50			92							53	86
55			82							0	81
60			96								81
65			87								84
70			85								76
75			17								75
80			0								81
85											82



Table 2. Water Quality Profile Data (Cont'd)

July 19, 1978

July 21, 1978

STATION	S1	S2	S3	S3	S2	S1	P1	P2	P3	P4	P5
TIME	1405	1545	1647	855	927	937	951	1007	1025	1042	1053
DEPTH (METERS)	TEMPERATURE (C)										
0	18.63	18.64	18.64	17.31	16.30	16.30	16.15	16.11	15.78	18.70	18.91
5	18.60	18.57	18.63	17.29	16.24	16.18	16.08	15.85	15.54	15.83	16.86
10	17.34	17.13	17.97	17.09	16.10	16.07	15.92	15.67	14.93	14.60	14.76
15	14.67	14.56	15.57	16.37	15.38	15.43	15.18	14.53	13.32	12.70	13.05
20	13.87	13.58	13.96	14.74	12.85	12.89	12.68	12.70	12.23	12.13	12.20
25	13.20	12.72	12.70	12.76	12.02	11.91	11.69	11.86	11.72	12.04	
30	12.65	12.26	11.91	12.05	11.57	11.50	11.24	11.41	11.32		
35	11.94	11.67	11.37	11.76	11.21	11.19	11.07	11.10	11.27		
40	11.57	11.29	11.18	11.46	11.06	11.01	10.98	11.03			
45	11.22	11.05	11.14	11.28	10.88	10.88	10.80				
50	10.94	10.92	11.03	11.03	10.69	10.63	10.64				
55	10.63	10.44	10.71	10.89	10.58	10.51	10.55				
60	10.34	10.29	10.68	10.75	10.56	10.48	10.54				
65	10.25	10.23	10.27	10.64	10.50	10.45					
70	10.22	10.21	10.04	10.61	10.47	10.42					
75	10.15	10.20	9.94	10.52	10.40	10.41					
80		10.20	9.88	10.49	10.31						
85			9.84	10.49	10.31						

SALINITY (PPT)											
0	33.17	33.18	33.15	32.84	32.88	32.71	32.81	32.82	32.89	32.67	32.77
5	33.19	33.14	33.14	32.83	32.84	32.73	32.79	32.84	32.85	32.80	32.20
10	32.92	33.03	33.08	32.84	32.82	32.75	32.77	32.83	32.93	33.20	32.90
15	33.16	33.31	32.97	32.66	32.59	32.68	32.65	32.41	32.92	33.42	33.03
20	33.29	33.24	33.07	32.59	32.78	33.17	32.74	32.68	33.09	33.29	33.05
25	33.23	33.29	33.20	32.86	33.08	33.47	32.93	32.94	33.09	33.30	
30	33.24	33.24	33.31	33.03	33.07	33.47	33.11	32.96	33.13		
35	33.25	33.20	33.28	33.00	33.06	33.24	33.07	33.13	33.13		
40	33.26	33.25	33.39	33.05	33.00	33.30	33.15	33.27			
45	33.32	33.44	33.43	33.11	33.11	33.41	33.27				
50	33.45	33.46	33.46	33.10	33.13	33.32	33.28				
55	33.44	33.47	33.38	33.17	33.12	33.30	33.27				
60	33.49	33.52	33.41	33.18	33.16	33.28	33.31				
65	33.52	33.53	33.32	33.23	33.23	33.31					
70	33.52	33.52	33.43	33.22	33.23	33.36					
75	33.55	33.52	33.53	33.26	33.24	33.52					
80		33.52	33.46	33.25	33.30						
85			33.45	33.28	33.29						

DENSITY (SIGMA-T)											
0	23.74	23.75	23.72	23.81	24.00	23.94	24.05	24.07	24.20	23.34	23.36
5	23.76	23.73	23.71	23.81	24.06	23.99	24.05	24.14	24.22	24.12	23.42
10	23.86	24.00	23.83	23.86	24.07	24.03	24.08	24.17	24.42	24.69	24.43
15	24.65	24.79	24.30	23.89	24.05	24.11	24.14	24.10	24.74	25.25	24.88
20	24.91	24.93	24.73	24.19	24.73	25.02	24.73	24.68	25.09	25.26	25.06
25	25.00	25.14	25.08	24.81	25.11	25.44	25.06	25.04	25.18	25.29	
30	25.12	25.20	25.31	25.07	25.19	25.51	25.28	25.14	25.29		
35	25.26	25.28	25.39	25.10	25.25	25.40	25.28	25.32	25.29		
40	25.34	25.38	25.51	25.20	25.29	25.47	25.36	25.44			
45	25.45	25.57	25.55	25.20	25.35	25.58	25.49				
50	25.60	25.61	25.59	25.31	25.39	25.55	25.52				
55	25.65	25.70	25.59	25.39	25.41	25.56	25.53				
60	25.74	25.77	25.61	25.43	25.44	25.55	25.56				
65	25.77	25.79	25.62	25.48	25.51	25.58					
70	25.78	25.79	25.74	25.48	25.51	25.63					
75	25.82	25.78	25.84	25.53	25.53	25.75					
80		25.79	25.79	25.53	25.59						
85			25.79	25.55	25.59						

Table 2. Water Quality Profile Data (Cont'd)

July 19, 1978

July 21, 1978

STATION	S1	S2	S3	S3	S2	S1	P1	P2	P3	P4	P5
TIME	1405	1545	1647	855	927	937	951	1007	1025	1042	1053
DISSOLVED OXYGEN (MG/L)											
0	6.67	6.78	6.82	8.38	7.74	7.74	6.64	6.16	5.99	5.80	5.73
5	6.73	6.83	6.91	8.52	7.81	7.80	6.78	6.29	6.13	5.98	5.68
10	6.82	6.93	6.85	8.60	7.89	7.92	6.87	6.50	6.17	5.84	5.83
15	7.07	7.17	6.96	8.61	7.84	7.81	6.74	6.50	6.01	5.69	5.69
20	7.14	7.16	7.20	8.59	7.67	7.62	6.54	6.47	5.76	5.60	5.48
25	7.07	6.97	7.34	8.47	7.30	7.32	5.95	5.54	5.44	5.50	
30	6.94	6.76	7.02	8.12	6.89	6.91	5.25	5.47	4.98		
35	6.70	6.42	6.44	7.72	6.44	6.49	4.72	4.85	4.75		
40	6.35	5.99	6.00	7.28	6.09	6.14	4.35	4.23			
45	5.90	5.55	5.67	6.96	5.65	5.73	3.99				
50	5.22	5.00	5.38	6.61	5.27	5.36	3.64				
55	4.76	4.56	5.11	6.22	5.08	5.14	3.37				
60	4.38	4.14	4.85	5.87	4.97	4.89	3.34				
65	4.00	3.84	4.73	5.57	4.71	4.63					
70	3.80	3.65	4.65	5.43	4.49	4.39					
75	3.66	3.51	4.52	5.32	4.35	4.12					
80		3.47	4.38	5.27	4.23						
85			4.32	5.28	4.23						
HYDROGEN ION CONCENTRATION (PH)											
0	7.98	7.99	7.99	8.00	7.98	8.00	8.01	7.99	7.99	8.04	8.03
5	7.98	7.99	7.99	8.00	7.98	8.01	8.01	7.99	7.98	7.95	7.97
10	7.98	7.98	7.99	8.00	7.98	8.00	8.00	7.99	7.96	7.93	7.93
15	7.94	7.95	7.96	7.97	7.95	7.97	7.97	7.94	7.90	7.85	7.86
20	7.93	7.92	7.95	7.92	7.86	7.87	7.87	7.87	7.85	7.83	7.83
25	7.91	7.89	7.91	7.84	7.81	7.81	7.78	5.00	7.79	7.84	
30	7.89	7.87	7.84	7.79	7.76	7.77	7.73	2.75	7.74		
35	7.85	7.82	7.77	7.76	7.74	7.74	7.72	7.71	7.74		
40	7.82	7.78	7.77	7.74	7.72	7.73	7.71	7.71			
45	7.77	7.73	7.75	7.73	7.70	7.70	7.69				
50	7.72	7.71	7.75	7.70	7.70	7.70	7.68				
55	7.72	7.69	7.73	7.69	7.71	7.70	7.67				
60	7.69	7.68	7.72	7.68	7.70	7.69	7.67				
65	7.69	7.68	7.74	7.67	7.68	7.68					
70	7.69	7.68	7.73	7.67	7.68	7.68					
75	7.68	7.68	7.73	7.67	7.68	7.68					
80		7.68	7.73	7.68	7.68	7.68					
85			7.72	7.68	7.68						
LIGHT TRANSMITTANCE (%)											
0	76	74	71	58	59	59	57	54	44	22	9
5	76	74	72	60	61	64	57	56	55	34	30
10	72	70	69	61	68	69	68	65	66	55	42
15	73	71	64	69	73	75	75	75	43	33	15
20	75	72	66	73	75	78	74	80	50	36	1
25	75	72	69	75	78	77	63	62	61	18	
30	79	78	76	76	72	66	57	68	19		
35	81	80	80	78	78	76	54	68	0		
40	83	80	80	80	68	66	67	17			
45	85	84	82	82	64	71	68				
50	83	81	82	80	69	75	68				
55	80	68	83	83	72	73	16				
60	74	74	83	83	68	70	0				
65	75	71	84	83	67	69					
70	74	62	86	83	68	46					
75	71	58	86	84	70	15					
80		59	86	85	31						
85			87	85	0						

Table 2. Water Quality Profile Data (Cont'd)

July 21, 1978

STATION	P6	P7	S3	S1	P1	P2	P3	P4	P5	P6	P7
TIME	1108	1122	1418	1433	1501	1513	1530	1543	1557	1609	1620
DEPTH (METERS)	TEMPERATURE (C)										
0	18.39	19.66	18.08	18.05	17.47	17.44	16.74	16.80	16.08	16.46	19.21
5	16.19	16.62	17.25	17.40	16.91	16.66	16.09	15.59	15.32	15.52	17.26
10	14.50	13.38	16.72	16.77	16.27	16.00	15.22	14.56	14.91	14.65	14.21
15	13.14	12.52	15.86	16.14	15.81	15.81	13.87	12.61	13.33	13.37	12.63
20	12.39	12.47	14.28	14.71	14.40	15.45	12.21	12.06	12.27	12.39	
25			12.61	12.06	12.14	14.02	11.78	11.99			
30			11.77	11.28	11.34	12.03	11.20				
35			11.35	10.95	11.03	11.20	11.14				
40			11.14	10.74	10.89	11.02					
45			11.00	10.57	10.84						
55			10.85	10.42	10.70						
60			10.70	10.41	10.66						
65			10.53	10.40							
70			10.41	10.37							
75			10.29	10.25							
80			10.22	10.20							
85			10.21								

SALINITY (PPT)											
0	32.95	32.72	33.08	33.01	33.20	33.17	33.26	33.29	33.31	33.03	33.05
5	32.44	32.75	32.93	33.10	32.98	33.16	33.05	33.46	33.21	32.98	32.95
10	33.00	33.62	33.17	33.08	33.16	33.20	33.22	33.22	33.35	33.05	33.16
15	33.37	33.74	33.13	33.04	33.13	33.20	33.04	33.33	33.19	33.17	34.22
20	33.44	33.83	32.92	32.78	32.95	33.10	33.24	33.44	33.55	33.41	
25			33.04	33.04	33.11	32.94	33.35	33.36			
30			33.20	33.48	33.19	33.27	33.40				
35			33.34	33.35	33.39	33.42	33.33				
40			33.35	33.45	33.68	33.61					
45			33.35	33.51	33.48						
50			33.38	33.44	33.46						
55			33.39	33.39	33.50						
60			33.49	33.39	33.42						
65			33.50	33.40							
70			33.46	33.40							
75			33.55	33.43							
80			33.56	33.46							
85			33.57								

DENSITY (SIGMAT)											
0	23.63	23.13	23.80	23.76	24.05	24.03	24.26	24.27	24.45	24.15	23.50
5	23.76	23.89	23.89	23.98	24.01	24.20	24.25	24.67	24.54	24.32	23.90
10	24.56	25.27	24.20	24.12	24.30	24.39	24.58	24.72	24.74	24.57	24.74
15	25.13	25.53	24.36	24.23	24.38	24.43	24.72	25.20	24.94	24.92	25.88
20	25.33	25.61	24.54	24.35	24.54	24.43	25.21	25.39	25.43	25.30	
25			24.97	25.08	25.11	24.61	25.37	25.34			
30			25.25	25.56	25.33	25.26	25.51				
35			25.44	25.52	25.54	25.53	25.47				
40			25.49	25.64	25.78	25.71					
45			25.51	25.71	25.64						
50			25.55	25.66	25.64						
55			25.57	25.65	25.68						
60			25.67	25.65	25.63						
65			25.71	25.66							
70			25.70	25.66							
75			25.79	25.70							
80			25.81	25.74							
85			25.83								

Table 2. Water Quality Profile Data (Cont'd)

July 21, 1978

STATION TIME	P6 1108	P7 1122	S3 1418	S1 1433	P1 1501	P2 1513	P3 1530	P4 1543	P5 1557	P6 1609	P7 1620
DISSOLVED OXYGEN (MG/L)											
0	6.08	5.75	10.34	8.07	7.11	7.05	6.56	6.54	6.43	6.37	6.26
5	6.17	5.76	10.46	8.25	7.33	7.13	6.72	6.76	6.72	6.44	6.46
10	6.39	5.71	10.57	8.48	7.69	7.26	6.80	6.65	6.74	6.60	6.05
15	6.18	5.42	10.44	8.64	7.86	7.35	6.60	6.05	6.34	6.28	5.03
20	5.94	5.31	10.10	8.46	7.87	7.33	6.35	5.70	5.66	5.96	
25			9.80	8.16	7.78	7.13	6.09	5.60			
30			9.29	7.51	7.40	6.81	5.58				
35			8.81	6.91	6.97	6.15	5.48				
40			8.43	6.46	6.60	5.72					
45			8.10	6.21	6.26						
50			7.84	5.98	5.91						
55			7.52	5.66	5.59						
60			7.24	5.48	5.30						
65			7.01	5.40							
70			6.71	5.31							
75			6.53	5.21							
80			6.30	5.18							
85			6.28								
HYDROGEN ION CONCENTRATION (PH)											
0	8.04	8.03	8.00	8.03	7.97	8.01	7.97	7.97	7.97	7.96	8.05
5	7.94	7.95	7.99	8.02	7.97	8.00	7.97	7.96	7.97	7.96	7.99
10	7.92	7.83	7.98	8.01	7.97	8.00	7.95	7.92	7.95	7.94	7.86
15	7.84	7.81	7.96	8.00	7.96	8.00	7.89	7.83	7.86	7.87	7.83
20	7.81	7.81	7.89	7.94	7.90	7.99	7.82	7.82	7.83	7.82	
25			7.82	7.82	7.77	7.93	7.80	7.82			
30			7.75	7.75	7.73	7.83	7.73				
35			7.73	7.73	7.71	7.74	7.73				
40			7.73	7.71	7.71	7.73					
45			7.73	7.73	7.70						
50			7.72	7.72	7.70						
55			7.71	7.70	7.70						
60			7.70	7.70	7.70						
65			7.70	7.70							
70			7.68	7.70							
75			7.67	7.69							
80			7.67	7.68							
85			7.67								
LIGHT TRANSMITTANCE (%)											
0	13	9	67	62	66	62	62	58	58	47	18
5	36	32	61	64	60	59	61	40	53	44	14
10	50	6	66	62	54	59	54	34	42	31	2
15	5	0	69	59	69	63	47	27	4	7	0
20	0	0	69	74	71	64	39	26	4	0	
25			78	67	62	54	54	0			
30			80	58	57	69	17				
35			81	56	63	59	0				
40			82	70	64	0					
45			84	77	62						
50			86	68	61						
55			86	63	25						
60			87	63	0						
65			86	63							
70			88	64							
75			67	24							
80			68	0							
85			69								

Table 2. Water Quality Profile Data (Cont'd)

July 25, 1978		July 26, 1978										
Station Time	T1 2115	T2 2122	T3 2132	T4 2146	T5 2212	T6 2222	T7 2228	T8 2237	T8 0044	T7 0052	T6 0059	T5 0106
Depth (meters)	TEMPERATURE (C)											
0	20.04	20.49	20.00	18.98	19.59	20.01	19.76	19.66	18.10	19.51	19.39	18.69
1	20.05	20.43	20.01	18.97	19.59	20.02	19.73	19.58	18.07	19.49	19.41	18.62
2	19.89	20.18	20.00	18.83	19.35	19.78	19.22	18.66	17.94	19.13	19.46	18.39
3	19.51	19.68	19.88	18.57	18.68	18.66	18.47	17.71	17.62	18.07	19.07	18.22
4	18.97	18.92	19.41	18.11	17.80	17.10	17.66	17.16	17.16	17.09	17.77	17.97
5	18.46	18.07	18.65	17.44	17.04	16.11	16.99	16.86	16.68	16.47	16.48	17.28
6	17.97	17.26	17.63	16.73	16.39	15.72	16.56	16.63	16.25	16.06	15.92	16.27
7	17.33	16.65	16.67	16.14	15.82	15.55	16.23	16.40	15.88	15.82	15.62	15.55
8	16.58	16.17	15.91	15.76	15.33	15.41	15.90	16.21	15.68	15.70	15.38	15.20
9	15.96	15.93	15.44	15.46	14.99	15.26	15.62			15.63	15.13	15.04
10	15.55		15.20	15.15	14.72	15.11					14.89	14.94
11	15.48		15.11	14.82	14.57							14.81
12				14.64	14.49							14.71
13					14.44							
14					14.39							
15					14.34							
16					14.34							
	SALINITY (ppt)											
0	32.82	32.99	32.95	32.96	32.86	32.74	32.81	31.59	33.02	31.56	31.96	32.76
1	32.86	33.03	33.02	33.00	32.90	32.69	32.87	31.80	33.04	31.93	32.37	33.00
2	32.84	32.95	33.06	32.95	32.79	32.67	32.69	32.39	33.03	32.56	32.85	33.09
3	32.78	32.75	32.96	32.84	32.68	32.62	32.57	32.70	33.01	32.77	32.78	33.12
4	32.77	32.62	32.77	32.71	32.68	32.76	32.60	32.89	33.00	32.89	32.63	33.03
5	32.74	32.56	32.62	32.58	32.76	32.96	32.75	33.01	33.04	33.08	32.82	32.90
6	32.69	32.65	32.50	32.62	32.80	33.12	32.92	33.06	33.08	33.18	33.07	32.88
7	32.62	32.78	32.61	32.70	32.87	33.15	33.02	33.09	33.16	33.21	33.17	32.99
8	32.69	32.89	32.73	32.85	32.95	33.11	33.12	33.17	33.22	33.26	33.19	33.14
9	32.86	33.02	32.90	32.85	32.99	33.14	33.32			33.23	33.18	33.19
10	33.09		33.02	32.86	33.05	33.16					33.22	33.21
11	33.24		32.97	32.90	33.10							33.23
12				32.97	33.11							33.35
13					33.11							
14					33.14							
15					33.19							
16					33.26							
	DENSITY (Sigma-t)											
0	23.12	23.12	23.22	23.49	23.26	23.06	23.18	22.27	23.75	22.29	22.63	23.41
1	23.14	23.17	23.27	23.52	23.29	23.02	23.23	22.46	23.78	22.58	22.93	23.61
2	23.16	23.17	23.31	23.52	23.27	23.07	23.23	23.13	23.80	23.15	23.29	23.74
3	23.22	23.15	23.26	23.50	23.35	23.31	23.32	23.61	23.86	23.57	23.33	23.80
4	23.35	23.25	23.23	23.52	23.57	23.79	23.54	23.88	23.97	23.90	23.53	23.79
5	23.45	23.41	23.31	23.58	23.81	24.18	23.81	24.05	24.11	24.18	23.99	23.85
6	23.54	23.67	23.47	23.78	23.99	24.39	24.04	24.13	24.23	24.36	24.31	24.07
7	23.63	23.92	23.78	23.97	24.17	24.44	24.19	24.21	24.38	24.43	24.45	24.33
8	23.87	24.11	24.04	24.17	24.35	24.45	24.35	24.31	24.47	24.50	24.52	24.51
9	24.13	24.27	24.28	24.24	24.45	24.50	24.56			24.49	24.56	24.59
10	24.40		24.42	24.31	24.55	24.55					24.65	24.63
11	24.53		24.40	24.42	24.62							24.67
12				24.51	24.65							24.78
13					24.66							
14					24.69							
15					24.74							
16					24.80							

Table 2. Water Quality Profile Data (Cont'd)

July 25, 1978		July 26, 1978										
Station Time	T1 2115	T2 2122	T3 2132	T4 2146	T5 2212	T6 2222	T7 2228	T8 2237	T8 0044	T7 0052	T6 0059	T5 0106
Depth (meters)	DISSOLVED OXYGEN (mg/l)											
0	10.44	10.20	10.17	10.45	9.89	9.33	9.44	7.34	7.75	7.68	8.31	9.90
1	10.44	10.23	10.25	10.46	9.93	9.38	9.42	7.26	7.73	7.72	8.36	9.94
2	10.43	10.18	10.42	10.40	9.99	9.57	9.29	7.25	7.90	8.03	8.82	9.91
3	10.43	10.06	10.56	10.26	10.01	9.78	9.22	7.48	8.16	8.74	9.27	9.78
4	10.57	9.93	10.59	10.07	10.03	9.94	9.16	7.90	8.39	9.08	9.13	9.58
5	10.74	9.71	10.56	9.79	10.07	9.92	9.07	8.32	8.31	8.76	8.57	9.30
6	10.82	9.40	10.36	9.43	10.01	9.81	8.78	8.49	7.56	8.01	7.95	8.90
7	10.67	9.00	10.08	9.01	9.79	9.77	8.31	8.24	6.11	7.25	7.36	8.38
8	10.24	8.54	9.63	8.58	9.47	9.76	7.69	7.72	4.85	6.72	6.76	7.83
9	9.61	8.28	9.21	8.19	9.16	9.67	7.13			6.36	6.38	7.43
10	8.88		8.82	7.86	8.92	9.54					6.46	7.16
11	8.73		8.63	7.57	8.73							6.97
12				7.38	8.58							6.88
13					8.46							
14					8.34							
15					8.20							
16					8.16							
	LIGHT TRANSMITTANCE (percent)											
0	12	11	14	20	19	17	15	7	11	16	19	23
1	12	9	14	14	20	17	15	8	11	16	19	24
2	15	10	15	13	20	15	14	11	11	16	19	23
3	17	12	16	13	20	10	11	11	10	12	15	21
4	18	14	17	18	19	7	9	10	6	9	13	20
5	18	13	14	18	17	7	7	6	3	7	10	18
6	17	11	10	19	11	5	2	3	1	6	7	14
7	13	4	7	18	4	2	0	0	0	5	5	11
8	8	0	5	16	1	0	0	0	0	0	4	8
9	2	0	2	13	0	0	0	0	0	0	0	6
10	0		0	6	0	0					0	2
11	0		0	0	0							0
12				0	0							0
13					0							
14					0							
15					0							
16					0							

Table 2. Water Quality Profile Data (Cont'd)

July 26, 1978

Station Time	T4 0113	T3 0119	T2 0124	T1 0132	T8 0805	T7 0821	T6 0828	T5 0835	T4 0843	T3 0852	T2 0859	T1 0906
Depth (meters)	TEMPERATURE (C)											
0	18.90	19.77	19.81	19.32	18.82	19.31	20.13	19.59	19.12	19.70	21.24	21.00
1	18.90	19.77	19.76	19.33	18.86	19.40	20.10	19.51	19.17	19.70	21.13	20.92
2	18.77	19.44	19.36	19.19	18.85	19.50	19.88	19.12	18.82	19.36	20.57	20.36
3	18.49	18.80	18.73	18.77	18.61	19.21	19.28	18.47	18.25	18.36	19.71	19.37
4	17.95	17.98	17.97	18.21	18.12	18.41	18.35	17.80	17.62	17.49	18.57	18.31
5	17.28	17.36	17.24	17.68	17.59	17.55	17.38	17.21	17.13	17.06	17.42	17.53
6	16.61	16.82	16.58	17.29	17.22	17.01	16.67	16.56	16.66	16.68	16.59	17.01
7	16.00	16.34	16.02	16.91	16.96	16.70	16.16	15.86	16.15	16.11	16.08	16.53
8	15.50	15.91	15.69	16.51	16.88	16.25	15.78	15.35	15.63	15.53	15.81	16.05
9	15.15	15.62	15.50	16.02		15.80	15.45	15.02	15.25	15.11	15.61	15.69
10	14.90	15.40	15.43	15.54			15.26	14.85	14.99	14.88	15.44	15.41
11	14.73	15.22		15.06				14.68	14.82	14.75		15.25
12	14.56	15.15		14.89				14.66	14.68	14.72		
13	14.52								14.67			
	SALINITY (ppt)											
0	32.85	33.16	32.74	33.11	31.26	32.31	33.16	33.29	32.97	33.16	33.02	33.19
1	33.03	33.16	32.96	33.19	32.11	32.58	33.17	33.21	33.01	33.22	33.02	33.19
2	32.98	32.97	32.94	33.03	32.69	32.94	33.10	33.09	32.78	33.00	32.87	33.02
3	32.83	32.82	32.79	32.89	32.95	33.01	32.95	32.99	32.68	32.87	32.76	32.90
4	32.70	32.78	32.78	32.86	32.92	32.94	32.88	33.04	32.71	33.03	32.79	32.97
5	32.69	32.86	32.79	32.94	33.01	33.01	33.03	33.08	32.79	33.13	32.96	33.13
6	32.67	32.88	32.86	33.04	33.10	33.18	33.12	33.10	32.80	33.06	33.11	33.18
7	32.72	32.92	32.95	33.02	33.16	33.25	33.19	33.12	32.78	33.02	33.26	33.18
8	32.80	32.99	33.08	32.98	33.19	33.17	33.31	33.13	32.83	33.03	33.34	33.15
9	32.90	33.08	33.16	32.94		33.28	33.35	33.26	32.93	33.19	33.40	33.18
10	33.00	33.13	33.22	32.91			33.47	33.33	33.01	33.26	33.43	33.27
11	33.06	33.15		32.95				33.43	33.06	33.25		33.33
12	33.08	33.17		32.88				33.49	33.07	33.18		
13	33.09								33.01			
	DENSITY (Sigma-t)											
0	23.43	23.44	23.11	23.52	22.23	22.91	23.35	23.59	23.46	23.46	22.94	23.14
1	23.56	23.44	23.29	23.57	22.87	23.10	23.37	23.55	23.48	23.51	22.97	23.16
2	23.56	23.38	23.38	23.49	23.32	23.34	23.37	23.55	23.40	23.43	23.01	23.18
3	23.52	23.43	23.42	23.49	23.58	23.47	23.41	23.64	23.46	23.58	23.15	23.34
4	23.55	23.60	23.60	23.61	23.67	23.61	23.59	23.84	23.64	23.91	23.46	23.66
5	23.70	23.81	23.79	23.80	23.87	23.88	23.93	24.02	23.81	24.09	23.87	23.97
6	23.84	23.95	23.99	23.96	24.03	24.14	24.17	24.18	23.93	24.12	24.18	24.14
7	24.02	24.09	24.19	24.04	24.14	24.26	24.34	24.35	24.03	24.22	24.41	24.25
8	24.19	24.24	24.36	24.10	24.17	24.31	24.52	24.48	24.19	24.36	24.53	24.33
9	24.35	24.38	24.47	24.18		24.49	24.62	24.65	24.35	24.57	24.62	24.44
10	24.47	24.46	24.53	24.27			24.76	24.74	24.46	24.67	24.69	24.57
11	24.56	24.52		24.40				24.85	24.54	24.70		24.65
12	24.61	24.55		24.38				24.90	24.57	24.65		
13	24.63								24.53			

Table 2. Water Quality Profile Data (Cont'd)

July 26, 1978

Station Time	T4 0113	T3 0119	T2 0124	T1 0132	T8 0805	T7 0821	T6 0828	T5 0835	T4 0843	T3 0852	T2 0859	T1 0906
Depth (meters)	DISSOLVED OXYGEN (mg/l)											
0	9.85	9.62	9.63	9.86	5.75	8.04	9.26	9.23	9.42	9.49	8.03	8.37
1	9.90	9.68	9.65	9.89	5.84	8.06	9.28	9.28	9.46	9.49	8.04	8.39
2	9.99	9.77	9.60	9.85	6.25	8.28	9.30	9.42	9.53	9.38	8.11	8.47
3	10.09	9.88	9.56	9.75	6.76	8.60	9.26	9.56	9.67	9.17	8.32	8.63
4	10.15	9.93	9.48	9.61	7.25	8.86	9.16	9.64	9.84	8.98	8.69	8.89
5	10.08	9.86	9.28	9.44	7.56	9.06	9.01	9.53	10.01	8.79	9.05	9.08
6	9.89	9.62	8.93	9.26	7.70	9.10	8.64	9.20	10.08	8.52	9.10	9.13
7	9.61	9.25	8.50	9.01	7.68	8.90	7.94	8.74	10.03	8.17	8.94	9.04
8	9.30	8.83	8.04	8.68	7.52	8.26	7.21	8.26	9.86	7.74	8.70	8.91
9	8.97	8.44	7.61	8.25		7.40	6.43	7.81	9.60	7.32	8.44	8.74
10	8.67	8.11	7.33	7.77			5.89	7.45	9.32	6.93	8.24	8.54
11	8.41	7.83		7.31				7.16	9.04	6.64		8.32
12	8.21	7.73		7.18				7.13	8.81	6.55		
13	8.17								8.80			
	LIGHT TRANSMITTANCE (percent)											
0	14	16	13	19	12	11	11	7	16	13	9	5
1	12	14	11	19	13	11	13	11	17	16	12	6
2	10	13	10	19	13	11	15	18	18	17	15	12
3	8	13	11	20	14	13	16	19	17	15	17	16
4	6	11	11	21	11	15	15	18	16	13	18	16
5	8	8	8	21	5	17	11	16	17	12	18	18
6	8	11	5	19	0	8	8	14	15	13	14	23
7	8	8	2	16	0	1	6	13	10	12	9	25
8	7	5	2	13	0	0	2	12	6	8	4	19
9	4	1	0	10		0	0	8	4	4	0	6
10	2	0	0	3			0	1	2	0	0	0
11	0	0		0				0	0	0	0	0
12	0	0		0				0	0	0	0	0
13	0	0		0				0	0	0	0	0



Table 3. Solar Irradiance Data

7/18	Station Time	P2 1422	P1 1527	S1 1545	P4 1620						
	Depth (meters)										
	0	7.63	8.27	8.27	6.05						
	2	5.88	5.58	7.00	4.14						
	6	4.13	3.81	3.50	2.23						
	12	2.86	2.39	2.07	0.95						
	Percent Penetration	37	29	25	16						
7/19	Station Time	P7 0911	P6 1040	P5 1152	P4 1212	P3 1246	P2 1301	P1 1337	S1 1405	S2 1534	S3 1647
	Depth (meters)										
	0	1.08	1.91	2.79	3.50	3.50	3.81	2.52	7.32	5.09	6.36
	2	0.34	0.73	1.75	1.75	2.23	2.07	2.07	5.58	5.09	4.45
	6	0.15	0.34	0.98	1.08	1.83	1.91	1.91	4.61	3.83	3.02
	12	0.07	0.18	0.57	0.64	1.46	1.60	2.23	3.18	2.55	1.60
	Percent Penetration	06	09	20	18	42	42	88	43	50	25
7/21	Station Time	S3 0855	S2 0927	S1 0937	P1 0951	P2 1007	P3 1025	P4 1042	P5 1053	P6 1108	P7 1122
	Depth (meters)										
	0	1.60	1.91	1.83	2.55	2.47	3.34	2.79	4.13	5.24	3.18
	2	1.12	1.30	1.40	1.46	1.68	1.36	1.52	1.75	2.07	1.44
	6	0.92	1.02	0.98	1.20	1.27	1.03	0.84	0.78	0.89	0.95
	12	0.53	0.62	0.62	0.72	0.79	0.64	0.35	0.28	0.37	0.20
	Percent Penetration	33	32	34	28	32	19	12	07	07	06
7/21	Station Time	S3 1418	S2/S1 1433	P1 1501	P2 1513	P3 1530	P4 1543	P5 1557	P6 1609	P7 1620	
	Depth (meters)										
	0	2.79	4.14	8.26	8.90	8.26	7.00	7.00	7.00	5.57	
	2	1.60	3.50	5.72	6.51	6.27	5.64	5.56	5.40	3.18	
	6	1.30	2.07	4.45	4.61	3.66	3.34	3.81	3.34	1.11	
	12	0.35	1.24	2.23	2.23	1.91	1.37	1.68	1.06	0.07	
	Percent Penetration	12	30	27	25	23	20	24	15	01	
7/26	Station Time	T8 0805	T7 0821	T6 0828	T5 0835	T4 0843	T3 0852	T2 0859	T1 0906		
	Depth (meters)										
	0	4.46	4.45	5.40	5.58	6.36	6.05	6.36	6.36		
	2	1.08	0.83	1.60	1.91	2.07	2.23	2.07	1.91		
	6	0.21	0.30	0.35	0.46	0.48	0.48	0.48	0.38		
	12	0.07	0.03	0.05	0.05	0.02	0.02	0.12	0.08		
	Percent Penetration	02	01	01	01	<01	<01	02	01		

and 250 feet (1, 37, and 76 m) to measure surface, mid-depth, and bottom currents. The meters were installed on July 21 for a period of 24 hours. Results of these measurements are presented in Figure 7(a). Due to a malfunction of the bottom current meter, no bottom current data were collected during this period. However, bottom current measurements were obtained on July 25-26 when a new current meter was installed at the site. Results of these measurements are presented in Figure 7(b).

Surface currents at the proposed platform site exhibited a strong tidal influence. Current direction advanced progressively clockwise over the 24-hour recording period, reflecting a progressive tidal wave with a 24 hour period. The selected sampling date coincided with the largest tidal extreme of the year. Current speed varied between 0.34 and 0.64 knots and averaged 0.51 knots.

Tidal influences at mid-depth were weaker with more of an alternating directional pattern. Predominant current direction was toward the northwest during flood tide and toward the southwest during ebb tide. Current speeds varied between 0.12 and 0.46 knots and averaged 0.27 knots.

Bottom currents showed a tidal influence with predominant current direction toward the southwest. During flood tide the current flowed predominantly toward the west, while during ebb tide it shifted toward the south-southwest. Current speeds varied between 0.15 and 0.49 knots and averaged 0.28 knots.

These results are similar to those reported by SCCWRP (1973) for San Pedro Channel, where the tidal current was described as rotary (direction changes progressively clockwise with time) with net resultant current speed of 0.26 to 0.38 knots. However, they do not directly coincide with other sources (OSI, 1978, Hancock, 1965) for net surface current flow during summer months (toward the east or northwest at 0.30 knots). This difference may be due to a combination of reduced wind influence and extreme tidal variation during the survey period, thus allowing for the development of a strong tidal eddy.

Current measurements along the pipeline corridor at Station P4 were collected in a similar manner to those collected at the platform site. Current meters were moored at 3, 36, and 50 foot (1, 11, and 15 m) depths for a period of 25 hours on July 18. Results of these measurements are presented in Figure 8. The bottom current meter was the same one which malfunctioned at the platform site and consequently no bottom data was recovered for this station.

Surface currents were tidally influenced and followed a pattern similar to the one recorded at Station S1. Tidal currents were rotary with a 24-hour period and the current speed varied between 0.12 and 0.43 knots. The predominant current direction for mid-depth currents varied between south and southeast with speed registering between 0.09 and 0.46 knots. The average current speed at this depth was 0.21 knots.





Figure 7. Surface, Mid-Depth, and Bottom Current Data for the Proposed Platform Site Station




Figure 8. Surface and Mid-Depth Current Data for Station P4

In-harbor current measurements were conducted at Station T4, just off the southeast tip of oil island Freeman. A Bendix Q-15 ducted current meter recorded current speed and direction hourly at depths of 3, 5, 10, 15, 20, 30, and 40 feet. Measurements were taken on July 25 and 26 for a 25-hour period and the results are presented on Figure 9.

The in-harbor currents were strongly influenced by tide at all depths. Maximum current velocities occurred during mid-tide and minimum current velocities occurred during slack tide. Current direction was not uniform over the entire water column.

Surface currents were strongly influenced by diurnal northeast winds during daylight hours (0600 to 1800 hours). Winds were averaging from 10 to 15 knots from the southwest and surface currents flowed toward the east-northeast (to the right of the wind) at an average of 0.14 knots. Between 1800 and 0600 hours, winds were calm and surface currents flowed predominantly toward the southwest at an average speed of 0.12 knots.




Figure 9. Wind, Current, and Tide Data for Station T4  
VIII-34




Figure 10. Surface Drogue Tracks for Station S1

Current measurements between the 10 and 20 foot (3 and 6 m) depths were considered to represent mid-depth currents and appeared to follow a clockwise rotary pattern corresponding to the 25-hour tidal cycle. Current speeds ranged from less than 0.05 to 0.23 knots.

Bottom currents (those between 30 and 40 feet) (9 and 12 m) flowed predominantly toward the south-southwest and tidal periodicity was evident. Maximum speeds were measured during high slack tide (0.13 knots) with minimum speeds occurring at mid-tide.

Drogue tracks for the July 21 and 22 releases at Station S1 are shown on Figure 10. The nine drogues released during slack tide before higher-high water (July 21) and during slack tide before lower-low water (July 22) moved in a southwesterly direction (between  $220^{\circ}$  T and  $240^{\circ}$  T) at speeds of 0.43 and 0.45 knots, respectively. During both drogue releases, winds were calm, minimizing possible wind-induced transport and maximizing both tidal and regional current influences.

The drogues released at 0230 on July 22 appeared to shift toward the northwest five hours after release. This shift in direction may have been the result of the tidal influence. The drogue results for Station S1 correspond to the current meter measurements taken during the same period (Figure 7(a)).

Although an equipment malfunction precluded the exact positioning of the P4 station drogue release, visual observations were recorded during each tracking. The first release was made at 1645 hours on July 18, during a period of slack tide before higher-high water. Winds at the time of release were from the west ( $265^{\circ}$  T) at 11 knots. All drogues traveled in a west-southwesterly direction and were retrieved approximately six hours later just inshore of oil platform "Eva". The calculated net current speed was approximately 0.30 knots.

A second drogue release was made during slack tide before lower-low water (0100 hours) on July 19. Drogues released during this period followed a similar course to those previously discussed. An exact visual retrieval location was not recorded for the second drop, so no attempt was made at estimating the speed of transport.

Two drogue releases were made at Station T4, inside Long Beach Harbor, on July 25. Five drogues were released per drop, once during slack tide before higher-high water and once during slack tide before lower-low water. Results of these releases are presented on Figure 11.

Drogues released at 1200 hours moved predominantly toward the southwest at 0.18 to 0.20 knots between 1200 and 1400 hours. Then, between 1400 and 1600 hours, four of the five drogues altered course and moved toward the southeast at 0.07 to 0.20 knots. The remaining drogue continued to travel southwesterly.




Figure 11. Surface Drogue Tracks for Station T1



Three of the five drogues released at 1800 hours moved toward the northwest at 0.30 to 0.40 knots for one hour, altered course toward the southwest at 0.30 knots for two hours, then headed almost due west at 0.12 to 0.16 knots. Net movement was toward the southwest at 0.13 knots. The remaining two drogues followed similar patterns, but net movement was toward the north and northwest.

This complex movement pattern, demonstrated by the in-harbor drogues, was consistent with current meter results previously discussed, and is indicative of the complex tidal and eddy patterns within the harbor area. Similar results have been reported by both the Allan Hancock Foundation (1972) and the U. S. Army Engineers (1975) for this area.

### Chemical Oceanographic Analyses

Chemical oceanographic studies included sampling and analyses of receiving waters and sediments from 18 stations; 1 at each of the 3 platform sites, 7 along the pipeline, and 8 in the terminus area. Surface receiving water chemical parameters measured included: 1) nutrients; ammonia, nitrate, phosphate, and silicate, 2) grease and oil, 3) trace metals; total chromium, barium, iron, lead, aluminum, and zinc, and 4) coliform bacteria. Sediment chemical parameters measured included: 1) organic content; biochemical oxygen demand, grease and oil, and phenol, 2) trace metals; cadmium, copper, lead, silver, zinc, cobalt manganese, arsenic; boron, nickel, mercury, total chromium, barium, iron, and aluminum, and 3) sediment description.

Surface Receiving Waters. All surface receiving water chemical analysis results are presented in Table 4. Nutrients measured are as follows: ammonia nitrogen at the platform sites only and nitrate, phosphate, and silicate at all 18 stations. Samples collected at all stations were analyzed for grease and oil. Samples collected at the 3 platform site stations were analyzed for total chromium, barium, iron, lead, and aluminum. Samples collected at the 7 pipeline stations were analyzed for zinc. Samples were collected at the 3 platform site stations and analyzed for coliform organisms and results are presented in Table 5.

Table 4. Receiving Water Chemical Analysis Results (mg/l)

Station No.	Ammonia Nitrogen	Nitrate Nitrogen	Phosphate Phosphorus	Silicate (Si)	Grease and Oil	Total Chromium	Barium	Iron	Lead	Aluminaum	Zinc
S1	0.003	0.029	0.036	0.11	<5	<0.0005	<0.1	0.0029	0.0065	0.0016	-
S2	0.006	0.032	0.027	0.11	<5	<0.0005	<0.1	0.0057	0.0069	0.0010	-
S3	0.003	0.050	0.022	0.099	<5	<0.0005	<0.1	0.0052	0.0048	0.0024	-
P1	-	0.020	0.030	0.12	<5	-	-	-	-	-	0.0012
P2	-	0.025	0.032	0.13	<5	-	-	-	-	-	0.0015
P3	-	0.026	0.025	0.092	<5	-	-	-	-	-	0.0040
P4	-	0.021	0.030	0.12	<5	-	-	-	-	-	0.0019
P5	-	0.022	0.030	0.16	<5	-	-	-	-	-	0.0008
P6	-	0.019	0.032	0.16	<5	-	-	-	-	-	0.0009
P7	-	0.020	0.025	0.22	<5	-	-	-	-	-	0.0017
T1	-	0.047	0.035	0.31	<5	-	-	-	-	-	-
T2	-	0.048	0.021	0.34	<5	-	-	-	-	-	-
T3	-	0.037	0.035	0.26	<5	-	-	-	-	-	-
T4	-	0.024	0.030	0.24	<5	-	-	-	-	-	-
T5	-	0.034	0.030	0.18	<5	-	-	-	-	-	-
T6	-	0.034	0.031	0.26	<5	-	-	-	-	-	-
T7	-	0.030	0.032	0.26	<5	-	-	-	-	-	-
T8	-	0.045	0.030	0.47	<5	-	-	-	-	-	-

Table 5. Receiving Water Examination for Coliform Organisms

Station No.	Portions Examined		Presumptive Lactose Broth		Confirmed B.G.B.		Coliform Organisms Most Probable Number per 100 ml
	Volume	No.	24 hr.	48 hr.	24 hr.	48 hr.	
S1	10	3	0	0	0	0	<3
	1.0	3	0	0	0	0	
	0.1	3	0	0	0	0	
	0.01	3	0	0	0	0	
S2	10	3	0	0	0	0	<3
	1.0	3	0	0	0	0	
	0.1	3	0	0	0	0	
	0.01	3	0	0	0	0	
S3	10	3	0	0	0	0	<3
	1.0	3	0	0	0	0	
	0.1	3	0	0	0	0	
	0.01	3	0	0	0	0	

Nutrients. Surface receiving water nutrients were measured in order to gain a data base prior to the construction of the platforms and pipeline and to correlate them with the plankton dynamics of the study area.

Ammonia is normally present only in small concentrations in natural waters (SCCWRP, 1973). Samples collected at the 3 platform site stations were analyzed for ammonia in order to obtain the "natural" concentration found in the study area for later comparison with concentrations present after the proposed sanitary waste disposal from the platform. The ammonia concentrations measured ranged from 0.003 to 0.006 mg/l with a mean of 0.004 mg/l and a standard deviation of  $\pm 0.002$  mg/l.

Nitrate is the dominant inorganic nitrogenous nutrient found in natural open ocean waters. Nitrate concentrations at the platform station ranged from 0.029 to 0.050 mg/l with a mean of 0.037 mg/l and a standard deviation of  $\pm 0.011$ . Nitrate concentrations along the pipeline stations ranged from 0.019 to 0.026 mg/l with a mean of 0.022 mg/l and a standard deviation of  $\pm 0.003$  mg/l. Nitrate concentrations in the terminus area stations ranged from 0.024 mg/l to 0.045 mg/l with a mean of 0.037 mg/l and a standard deviation of  $\pm 0.009$  mg/l. The Hancock Foundation found nitrate concentrations in surface waters of the Southern California Bight ranged from 0.01 to 0.16 mg/l (SCCWRP, 1973). Nitrate concentrations observed during the B and C survey are in agreement with concentrations reported by SCCWRP.

Phosphate is considered to be the second most important nutrient next to nitrogen for phytoplankton metabolism. Phosphate concentrations at the platform site stations ranged from 0.022 to 0.036 mg/l with a mean of 0.283 mg/l and a standard deviation of  $\pm 0.007$  mg/l. Along the pipeline stations phosphate concentrations ranged from 0.025 to 0.032 mg/l with a mean of 0.029 mg/l and a

standard deviation of  $\pm 0.003$  mg/l. Phosphate concentrations in the terminus area stations ranged from 0.021 to 0.035 mg/l with a mean of 0.031 mg/l and a standard deviation of  $\pm 0.004$  mg/l. Surface phosphate concentrations of the southern California mainland shelf reported by the Hancock Foundation, 1965, agree with the concentrations observed during the B and C survey.

Silicate is an essential nutrient to diatoms. Diatoms comprise a great deal of the phytoplankton population found in the study area. Silicate concentrations at the platform site station ranged from 0.099 to 0.110 mg/l with a mean of 0.106 mg/l and a standard deviation of  $\pm 0.006$  mg/l. Silicate concentrations along the pipeline stations ranged from 0.092 to 0.220 mg/l with a mean of 0.143 mg/l and a standard deviation of  $\pm 0.042$  mg/l. Silicate concentrations observed in the terminus area stations ranged from 0.180 to 0.470 mg/l with a mean of 0.290 mg/l and a standard deviation of  $\pm 0.087$  mg/l. The surface silicate concentrations observed during the B and C survey were lower than those reported by the Hancock Foundation, 1965. Silicate concentrations are greatly dependent upon upwelling and phytoplankton assimilation. During the B and C survey phytoplankton uptake of silicate, without replenishment from upwelling, lowered concentrations below those reported by the Hancock Foundation, 1965.

Grease and Oil. Grease and oil samples were analyzed because of their close association to potential oil seepage and spills. Throughout the entire study area, platform site stations, pipeline stations, and terminus area,  $<5$  mg/l of grease and oil was detected in the surface receiving waters.

Trace Metals. Total chromium, barium, iron, and lead were analyzed because of their importance during construction and production as components of suspended compounds at the platform site stations. Analysis for aluminum was included because aluminum will be used as a sacrificial anode on the proposed platforms. Zinc was analyzed in surface receiving waters along the pipeline stations because it will be used as anodes on the pipeline. All measured total chromium concentrations were  $<0.1$  mg/l. Iron concentrations ranged from 0.0028 to 0.0057 mg/l with a mean of 0.0046 mg/l and a standard deviation of  $\pm 0.0016$  mg/l. Lead concentrations ranged from 0.0048 to 0.0069 mg/l with a mean of 0.0061 mg/l and a standard deviation of  $\pm 0.0011$  mg/l. Aluminum concentrations ranged from 0.0010 to 0.0024 mg/l with a mean of 0.0017 mg/l and a standard deviation of  $\pm 0.0007$  mg/l. Surface zinc concentrations along the pipeline stations ranged from 0.0008 to 0.0040 mg/l with a mean of 0.0017 mg/l and a standard deviation of  $\pm 0.0011$  mg/l.

Coliform Organisms. Most probable number of coliform organisms per 100 ml were analyzed in surface receiving waters at the platform site stations as background information for the proposed sanitary waste disposal from the platform. At the 3 platform site stations  $<3$  most probable number of coliform bacteria per 100 ml were observed.

Sediment Chemistry. The sediment chemistry analysis results are presented in Table 6. There are three areas of interest in the sediment analysis: 1) organic content; biochemical oxygen demand (BOD), grease and oil, and phenolic compounds

Table 6. Sediment Chemical Analysis Results (mg/kg)

Station No.	Biochemical Oxygen Demand (BOD <sub>5</sub> )	Grease and Oil	Phenol (C <sub>6</sub> H <sub>5</sub> OH)	Cadmium	Copper	Lead	Silver	Zinc	Cobalt	Manganese
S1	54	<50	0.87	0.30	14	8.7	0.47	77	53	150
S2	660	<50	5.5	0.22	12	8.2	0.37	67	53	150
S3	740	<50	0.13	0.38	15	2.2	0.35	76	57	150
P1	900	1300	-	-	-	-	-	71	-	-
P2	510	220	-	-	-	-	-	47	40	-
P3	220	<50	-	-	-	-	-	30	-	-
P4	210	<50	-	-	-	-	-	21	19	-
P5	57	<50	-	-	-	-	-	15	-	-
P6	180	<50	-	-	-	-	-	14	16	-
P7	140	<50	-	-	-	-	-	46	-	-
T1	390	390	0.62	0.40	22	52	0.43	95	69	160
T2	380	540	4.4	0.34	25	44	0.54	94	74	180
T3	450	490	1.4	1.2	40	80	0.59	140	95	210
T4	480	90	0.23	1.1	41	100	0.62	150	100	190
T5	330	<50	0.24	1.2	31	83	0.46	130	90	180
T6	340	<50	0.26	1.3	33	55	0.99	110	100	190
T7	340	76	0.004	1.0	42	68	0.56	130	110	240
T8	700	340	0.26	0.79	40	100	0.46	140	100	260

Station No.	Arsenic	Boron	Nickel	Mercury	Total Chromium	Barium	Iron	Aluminum	Sediment Description
S1	47	45	20	0.28	35	1.8	21,000	68	coarse silt
S2	51	40	20	0.33	34	1.8	20,000	63	coarse silt
S3	30	42	23	0.22	41	2.0	22,000	82	coarse silt
P1	-	-	-	-	-	-	-	-	very fine sand
P2	-	-	-	-	-	-	-	33	fine sand
P3	-	-	-	-	-	-	-	-	medium sand
P4	-	-	-	-	-	-	-	18	medium sand
P5	-	-	-	-	-	-	-	-	coarse sand
P6	-	-	-	-	-	-	-	12	coarse sand
P7	-	-	-	-	-	-	-	37	fine sand
T1	24	63	20	0.23	24	2.0	23,000	-	coarse silt
T2	33	70	22	0.24	26	1.8	24,000	-	coarse silt
T3	78	52	32	0.35	35	2.3	31,000	110	coarse silt
T4	110	58	32	0.40	34	2.2	33,000	-	medium silt
T5	67	53	30	0.23	28	1.9	29,000	110	coarse silt
T6	77	50	31	0.44	30	2.3	32,000	-	coarse silt
T7	79	53	33	0.41	30	2.2	36,000	140	medium silt
T8	100	60	29	0.30	28	2.2	33,000	-	coarse silt

were analyzed because of their close association to potential oil seepage, spills, and refinery processes, 2) trace metals; cadmium, copper, lead, silver, zinc, arsenic, boron, nickel, and mercury samples were analyzed because of their importance during construction and drilling as components of the suspended sediment. Cobalt and manganese were included because of their inclusion and discussion on other oil platform EIRs. Total chromium, barium, and iron were analyzed because they are major components in drilling muds. Aluminum and zinc were analyzed because they will be used as anodes on the platforms and along the pipeline, respectively, 3) sediment description as grain size was analyzed to complement all other sediment characteristics and to provide an indication of the indigenous benthic biota.

**Organic Content.** Biochemical oxygen demand (BOD), grease and oil, and phenol were analyzed as a representative of some of the organic content of the sediments. Sediment samples were collected at all the stations and analyzed for biochemical oxygen demand (BOD) and grease and oil. Samples were collected at the platform site stations and at the terminus area stations and analyzed for phenol content.

Concentrations of biochemical oxygen demand (BOD) at the platform site stations ranged from 54 to 740 mg/kg with a mean of 485 mg/kg and a standard deviation of  $\pm 573$  mg/kg. BOD concentrations along the pipeline stations ranged from 57 to 900 mg/kg with a mean of 317 mg/kg and a standard deviation of  $\pm 293$  mg/kg. BOD concentrations in the terminus area stations ranged from 330 to 700 mg/kg with a mean of 426 mg/kg and a standard deviation of  $\pm 123$  mg/kg. Sediment concentrations of grease and oil at the platform site stations were  $< 50$  mg/kg. Sediment concentrations of grease and oil along the pipeline were all  $< 50$  mg/kg, except at Stations P1 and P2 where concentrations were 1300 and 220 mg/kg respectively. Stations P1 and P2 were located along the proposed pipeline route near the platform sites and near an exploratory drilling platform. Concentrations observed at these two stations may have been due to the drilling platform or possible natural seepage within the area. Grease and oil concentrations varied greatly among the terminus area stations. In this area sediment grease and oil concentration patterns are the result of shipping traffic and industrial discharges. In the terminus area grease and oil ranged from  $< 50$  to 540 mg/kg.

Phenol concentrations at the platform site stations ranged from 0.13 to 5.50 mg/kg with a mean of 2.17 mg/kg and a standard deviation of  $\pm 2.91$  mg/kg. Phenol concentrations in the terminus area stations ranged from 0.004 to 4.40 mg/kg with a mean of 0.927 mg/kg and a standard deviation of  $\pm 1.47$  mg/kg.

**Trace Metals.** Trace metals sediment analyses included cadmium, copper, lead, silver, zinc, cobalt, manganese, arsenic, boron, nickel, mercury, chromium, barium, iron, and aluminum. Samples collected throughout the study area were analyzed for zinc, cobalt, and aluminum. Samples collected at the platform site stations and the terminus stations were analyzed for all trace metals.

Cadmium concentrations at the platform site stations ranged from 0.22 to 0.38 mg/kg with a mean of 0.30 mg/kg and a standard deviation of  $\pm 0.08$  mg/kg. In the terminus area sediment concentrations of cadmium ranged from 0.34 to 1.30 mg/kg with a mean of 0.92 mg/kg and a standard deviation of  $\pm 0.37$  mg/kg.

Copper concentrations at the platform site stations ranged from 12.0 to 15.0 mg/kg with a mean of 13.6 mg/kg and a standard deviation of  $\pm 1.53$  mg/kg. In the terminus area sediment copper concentrations ranged from 22.0 to 42.0 mg/kg with a mean of 34.3 mg/kg and a standard deviation of  $\pm 7.74$  mg/kg.

Lead concentrations at the platform site stations ranged from 2.2 to 8.7 mg/kg with a mean of 6.4 mg/kg and a standard deviation of  $\pm 3.6$  mg/kg. Lead concentrations in the terminus area stations ranged from 44.0 to 100 mg/kg with a mean of 72.8 and a standard deviation of  $\pm 21.5$  mg/kg.

Silver concentrations at the platform site stations ranged from 0.35 to 0.47 mg/kg with a mean of 0.40 mg/kg and a standard deviation of  $\pm 0.06$  mg/kg. In the terminus area silver concentrations ranged from 0.43 to 0.99 mg/kg with a mean of 0.58 mg/kg and a standard deviation of  $\pm 0.18$  mg/kg.

Zinc concentrations at the platform site stations ranged from 67.0 to 77.0 mg/kg with a mean of 73.3 mg/kg and a standard deviation of  $\pm 5.51$  mg/kg. Along the pipeline stations concentrations of zinc ranged from 14.0 to 71.0 mg/kg with a mean of 34.9 mg/kg and a standard deviation of  $\pm 20.9$  mg/kg. Zinc concentrations in the terminus area ranged from 94.0 to 150 mg/kg with a mean of 123 mg/kg and a standard deviation of  $\pm 21.3$  mg/kg.

Cobalt concentrations at the platform site stations ranged from 53 to 57 mg/kg with a mean of 54 mg/kg and a standard deviation of  $\pm 2.3$  mg/kg. Measured cobalt concentrations along the pipeline stations ranged from 26 to 40 mg/kg with a mean of 25 mg/kg and a standard deviation of  $\pm 13$  mg/kg. Cobalt concentrations in the terminus area ranged from 69 to 110 mg/kg with a mean of 92 mg/kg and a standard deviation of  $\pm 14$  mg/kg.

Manganese concentrations at the platform site stations were all 150 mg/kg. In the terminus area manganese concentrations ranged from 160 to 260 mg/kg with a mean of 201 mg/kg and a standard deviation of  $\pm 34$  mg/kg.

Arsenic concentrations at the platform site stations ranged from 30 to 51 mg/kg with a mean of 43 mg/kg and a standard deviation of  $\pm 11$  mg/kg. Arsenic concentrations in the terminus area ranged from 24 to 110 mg/kg with a mean of 71 mg/kg and a standard deviation of  $\pm 29$  mg/kg.

Boron concentrations at the platform site stations ranged from 40 to 45 mg/kg with a mean of 42 mg/kg and a standard deviation of  $\pm 2.5$  mg/kg. In the terminus area boron concentrations ranged from 50 to 70 mg/kg with a mean of 46 mg/kg and a standard deviation of  $\pm 7$  mg/kg.

Nickel concentrations at the platform site stations ranged from 20 to 23 mg/kg with a mean of 21 mg/kg and a standard deviation of  $\pm 1.7$  mg/kg. In the terminus area nickel concentrations ranged from 20 to 33 mg/kg with a mean of 29 mg/kg and a standard deviation of  $\pm 4.8$  mg/kg.

Mercury concentrations at the platform site stations ranged from 0.22 to 0.33 mg/kg with a mean of 0.28 mg/kg and a standard deviation of  $\pm 0.055$  mg/kg. In the terminus area mercury concentrations ranged from 0.23 to 0.44 mg/kg with a mean of 0.33 mg/kg and a standard deviation of  $\pm 0.09$  mg/kg.

Total chromium concentrations at the platform site stations ranged from 34 to 41 mg/kg with a mean of 37 mg/kg and a standard deviation of  $\pm 3.8$  mg/kg. In the terminus area chromium concentrations ranged from 24 to 35 mg/kg with a mean of 29 mg/kg and a standard deviation of  $\pm 3.7$  mg/kg.

Barium concentrations at the platform site stations ranged from 1.8 to 2.0 mg/kg with a mean of 1.9 mg/kg and a standard deviation of  $\pm 0.1$  mg/kg. In the terminus area barium concentrations ranged from 1.8 to 2.3 mg/kg with a mean of 2.1 mg/kg and a standard deviation of  $\pm 0.2$  mg/kg.

Iron concentrations at the platform site stations ranged from 20,000 to 22,000 mg/kg with a mean of 21,000 mg/kg and a standard deviation of  $\pm 1,000$  mg/kg. In the terminus area iron concentrations ranged from 23,000 to 36,000 mg/kg with a mean of 30,000 mg/kg and a standard deviation of  $\pm 4,500$  mg/kg.

Aluminum concentrations at the platform site stations ranged from 63 to 82 mg/kg with a mean of 71 mg/kg and a standard deviation of  $\pm 9.8$  mg/kg. Measured aluminum concentrations at pipeline stations ranged from 12 to 37 mg/kg with a mean of 25 mg/kg and a standard deviation of  $\pm 12$  mg/kg. In the terminus area measured aluminum concentrations ranged from 110 to 140 mg/kg with a mean of 120 mg/kg and a standard deviation of  $\pm 17$  mg/kg.

Sediment Description. The sediment at all platform site stations was observed to be coarse silt. Along the pipeline, sediments ranged from very fine sand to coarse sand. Very fine sand was found at the pipeline stations farthest offshore near the platform site stations, medium to coarse sands were found at stations along the middle of the proposed pipeline (outside the terminus area), and fine sand was found at the station closest to the Long Beach Breakwater. In the harbor area, medium and coarse silt sediments were observed with no apparent spatial patterns.